Settlement Agreement Report

Fourth Quarter October - December 2012

Prepared for the Technical Oversight Committee

Revised on May 9, 2013 (Original on February 13, 2013)



Final Shark River Slough flow data for Water Year 2012 (covering the first nine months of this reporting period) were not available in time for publication of the initial report (February 13, 2013). The Shark River Slough Section was updated using the final flow data for October 2011 to September 2012 and provisional flow data for October 2012 to December 2012 in this version.

Prepared by:

Cheol Mo, Violeta Ciuca, and Stuart Van Horn

Compliance Assessment and Reporting Section
Water Quality Bureau
South Florida Water Management District
3301 Gun Club Road
West Palm Beach, FL 33406

PURPOSE

The South Florida Water Management District has prepared this report to provide a quarterly update to the Technical Oversight Committee on the compliance status with total phosphorus levels or limits defined in the 1991 Settlement Agreement entered as a Consent Decree in 1992 and modified in 1995. The areas of interest in this report include the interior marsh stations in Arthur R. Marshall Loxahatchee National Wildlife Refuge and two discharges to Everglades National Park: inflows to Shark River Slough and inflows to Taylor Slough and Coastal Basins.

CONTENTS

	mmary					
Arthur R. Marshall Loxahatchee National Wildlife Refuge						
0	.d					
	Period Update					
Everglades N	lational Park	7				
Shark Rive	r Slough	7				
Backgro	und	7				
	ng Period Update					
Taylor Slou	ugh and Coastal Basins	16				
Backgro	und	16				
Reportir	ng Period Update	16				
	APPENDICES					
Appendix A	Monthly Total Phosphorus Concentration Data for the Arthur R. Marshall					
	Loxahatchee National Wildlife Refuge	A-1				
Appendix B	Weekly Grab Total Phosphorus Concentration Data for Shark River Slough.	B-1				
Appendix C	Weekly Grab Total Phosphorus Concentration Data for Taylor Slough and					
	Coastal Basins	C-1				
Appendix D	Calculation Methods	D - 1				
Appendix E	Document Revisions	E-1				
	TABLES					
Table 1.	Total phosphorus compliance, fourth quarter 2012	1				
Table 2.	Arthur R. Marshall Loxahatchee National Wildlife Refuge TP compliance					
	tracking.	6				
Table 3.	Shark River Slough total phosphorus compliance tracking	11				
Table 4.	Taylor Slough and Coastal Basins total phosphorus compliance tracking					
	FIGURES					
Figure 1.	Areas of interest.	2				
Figure 2.	Arthur R. Marshall Loxahatchee National Wildlife Refuge water quality samp					
	and stage measurement sites					
Figure 3.	(A) Monthly total phosphorus geometric mean concentrations for the Arthur					
0	Marshall Loxahatchee National Wildlife Refuge compared to calculated long					
	term levels, which are adjusted for fluctuations in stage. (B) Deviation of more					
	geometric mean total phosphorus concentrations with calculated long-term	J				
	levels.	5				
Figure 4.	Everglades National Park flow structures.					
Figure 5.	The 12-month flow-weighted mean total phosphorus concentrations at inflow					

	the Everglades National Park through Shark River Slough at the end of each
	water year compared to the interim and long-term total phosphorus
Eigene 6	limits
Figure 6.	National Park through Shark River Slough at the end of each month and the
	flow-weighted mean concentration for each sampling event
Figure 7.	Daily flows at Shark River Slough structures as a stacked sum of five inflows13
Figure 8.	Daily flows at structures S333 and S334.
Figure 9.	Daily flows at individual inflow structures to Shark River Slough. This figure
C	includes most of the data illustrated in Figures 7 and 8
Figure 10.	Flow at Shark River Slough structures on the day of sampling and the
_	corresponding flow-weighted mean total phosphorus concentrations for
	individual sampling events15
Figure 11.	The 12-month flow-weighted mean total phosphorus concentrations in inflows to
	Everglades National Park through Taylor Slough and Coastal Basins at the end
	of each water year compared to the 11 ppb long-term total phosphorus limit 18
Figure 12.	The 12-month flow-weighted mean total phosphorus concentrations in inflows to
	Everglades National Park through Taylor Slough and Coastal Basins at the end
	of each month and the flow-weighted mean total phosphorus concentration for
	each sampling event
Figure 13.	Daily flows measured in cubic feet per second into Everglades National Park as a
	stacked sum of Taylor Slough (structures S332D + S174) and Coastal Basins
	(structure S18C)
Figure 14.	Daily flows at individual Coastal Basins (S18C) and Taylor Slough (S332D +
	S174) structures into the Everglades National Park19
Figure 15.	Flow from Taylor Slough and Coastal Basins structures (S332D + S174 and S18C)
	on the day of sampling and the corresponding flow-weighted mean total
	phosphorus concentrations for individual sampling events21

ACRONYMS AND ABBREVIATIONS

cfs cubic feet per second ENP Everglades National Park

kac-ft thousand acre feet

NGVD 29 National Geodetic Vertical Datum of 1929

OFW Outstanding Florida Waters

ppb parts per billion

Refuge Arthur R. Marshall Loxahatchee National Wildlife Refuge

SFWMD South Florida Water Management District

TOC Technical Oversight Committee

 $\begin{array}{ll} TP & total \ phosphorus \\ \mu g/L & micrograms \ per \ liter \end{array}$

USACE United States Army Corps of Engineers

WCA Water Conservation Area

EXECUTIVE SUMMARY

This report fulfills the South Florida Water Management District's reporting requirements under the 1991 Settlement Agreement, entered as a Consent Decree in 1992 and modified in 1995, for the fourth quarter of 2012 (October – December 2012). Total phosphorus (TP) compliance highlights for this period are summarized below for the Arthur R. Marshall Loxahatchee National Wildlife Refuge (Refuge) and Everglades National Park, including Shark River Slough, and Taylor Slough and Coastal Basins (**Table 1** and **Figure 1**):

- **Refuge:** The geometric mean TP concentration was above the long-term level in October and below the long-term levels in November and December 2012.
- Shark River Slough: The provisional 12-month flow-weighted mean TP concentrations were below the 12-month moving long-term limit, calculated using provisional flow data, during the quarter.
- **Taylor Slough and Coastal Basins:** The 12-month flow-weighted mean TP concentrations were below the 12-month moving long-term limit during the quarter.

Table 1. Total phosphorus compliance, fourth quarter 2012.

Mont	h	Geometric Mean TP Concentration (ppb)		Long-term Level (ppb)		Mean Stage (feet NGVD 29)			Number of Samples	
Arthur R.	Arthur R. Marshall Loxahatchee National Wildlife Refuge									
Oct 20:	12		7.6	7.	2	1	17.20		14	
Nov 20	12		6.1	7.	3	1	17.11		14	
Dec 20	12		6.7	8.	3	1	16.82		14	
12-Month Period	Period Total Flow		12-Month Flow-weighted Mean		Long-term Limi		n Limit Grea		of Sampling Events ater than 10 ppb	
Ending	(ka	c-ft)	TP Concentration (ppb)		(ppb)		Guideline	е	Observed	
Everglade	s Nationa	al Park – S	hark River Slo	ugh						
Oct 2012	90	9.8	8.3		8.4	1	43.6		28.0	
Nov 2012	88	32.4	7.9		8.5	5	44.3		28.0	
Dec 2012	2012 873.1		3.1 7.8		8.5	5	44.5		28.0	
Everglades National Park — Taylor Slough and Coastal Basins										
Oct 2012	28	33.6	5.3	3	11.	0	53.1		0.0	
Nov 2012	27	73.0	5.1	[11.	0	53.1		0.0	
Dec 2012	27	79.6	5.1		11.	0	53.1		0.0	

- ppb = parts per billion. Values are actually in μg/L (micrograms per liter), which, for the purposes of this report, is equivalent to ppb.
- feet NGVD 29 = elevation in feet related to the National Geodetic Vertical Datum of 1929.
- kac-ft = thousand acre feet.
- Compliance for inflows to Everglades National Park (Shark River Slough, Taylor Slough and Coastal Basins) is evaluated annually based on the 12-month flow-weighted mean TP concentration for the federal water year ending on September 30.




Figure 1. Areas of interest.

ARTHUR R. MARSHALL LOXAHATCHEE NATIONAL WILDLIFE REFUGE

Background

The 1991 Settlement Agreement ended the Everglades lawsuit and was entered into by the federal government, the State of Florida, and the South Florida Water Management District. The subsequent Consent Decree, as modified in 1995, specified that interim and long-term total phosphorus (TP) concentration levels for the Arthur R. Marshall Loxahatchee National Wildlife Refuge (Refuge) must be met by February 1, 1999, and December 31, 2006, respectively. Both the interim and long-term concentration levels vary monthly because they are calculated as a function of water stage measured at gauging stations 1-7, 1-8C, and 1-9 within the Refuge. The stage range within which the interim and long-term concentration levels are applicable is 15.42 to 17.14 feet relative to the National Geodetic Vertical Datum of 1929 (NGVD 29). The monthly TP concentrations are determined from water samples collected at 14 interior marsh stations, LOX3 through LOX16 (Figure 2). As required in the Consent Decree, the concentrations are converted to a geometric mean, which is compared to the long-term concentration level. Monthly TP data for each station for the past 36 months are provided in Appendix A. The calculation methods specified in the Consent Decree are provided in Appendix D.

Reporting Period Update

Average sampling day stages in the Refuge were 17.20, 17.11, and 16.82 feet in October, November, and December 2012, respectively (**Figure 3** and **Table 2**). The geometric means, calculated from TP concentrations measured in water samples collected in October, November, and December 2012, were 7.6, 6.1, and 6.7 parts per billion (ppb), respectively. The geometric mean TP concentration was above the long-term level for the month of October 2012, and below the long-term levels for the months of November and December 2012.

TP samples were collected at all of the 14 stations for the months of October, November, and December 2012.




Figure 2. Arthur R. Marshall Loxahatchee National Wildlife Refuge water quality sampling and stage measurement sites.




Figure 3. (A) Monthly total phosphorus geometric mean concentrations for the Arthur R. Marshall Loxahatchee National Wildlife Refuge compared to calculated long-term levels, which are adjusted for fluctuations in stage. The long-term level was not applicable for May 2011 through July 2011 because the average stage was less than 15.42 feet. The geometric mean TP concentration was above the long-term level in October 2012. **(B)** Deviation of monthly geometric mean total phosphorus concentrations with calculated long-term levels. Values smaller than zero indicate that the geometric mean was lower than the long-term level.

Table 2. Arthur R. Marshall Loxahatchee National Wildlife Refuge TP compliance tracking.

Month	Geometric Mean TP Concentration (ppb)	Long-Term Level (ppb) Effective 12/31/2006	Average Stage ^a (ft NGVD 29)	Number of Samples
Jan-2010	5.2	8.9	16.68	14
Feb-2010	6.7	9.1	16.62	14
Mar-2010	5.9	9.0	16.65	14
Apr-2010	7.2	9.6	16.51	14
May-2010	7.8	10.4	16.35	14
Jun-2010	7.3	10.9	16.26	14
Jul-2010	6.9	10.7	16.29	14
Aug-2010	6.1	11.6	16.14	10
Sep-2010	6.5	8.5	16.76	14
Oct-2010	6.2	7.7	16.99	14
Nov-2010	6.6	8.2	16.83	14
Dec-2010	4.0	9.6	16.52	13
Jan-2011	5.1	10.7	16.29	10
Feb-2011	6.0	11.3	16.19	10
Mar-2011	8.1	14.0	15.79	8
Apr-2011	11.4	16.3	15.53	7
May-2011	no data	N/A ^b	14.87	0
Jun-2011	no data	N/A ^b	14.30	0
Jul-2011	20.2	N/A ^b	15.11	4
Aug-2011	11.5	16.6	15.50	8
Sep-2011	9.8	14.0	15.80	11
Oct-2011	7.4	12.1	16.06	11
Nov-2011	7.8	9.2	16.59	14
Dec-2011	6.2	9.2	16.59	7
Jan-2012	6.0	10.0	16.43	14
Feb-2012	7.5	11.1	16.23	13
Mar-2012	6.3	12.8	15.96	10
Apr-2012	8.1	13.1	15.92	10
May-2012	7.2	11.7	16.12	12
Jun-2012	7.4	11.6	16.14	14
Jul-2012	6.0	12.5	16.00	14
Aug-2012	5.5	11.3	16.18	13
Sep-2012	7.0	7.2	17.38	14
Oct-2012	7.6*	7.2	17.20	14
Nov-2012	6.1	7.3	17.11	14
Dec-2012	6.7	8.3	16.82	14

- ppb = parts per billion. Values are actually in µg/L (micrograms per liter), which, for the purposes of this report, is equivalent to ppb.
- ft NGVD 29 = elevation in feet related to the National Geodetic Vertical Datum of 1929.
- Highlighted rows with bold, italicized text indicate when an excursion over the long-term level occurred.
- The seven "J" flag qualified data collected on December 13, 2011, were excluded from the December 2011 geometric mean calculation following the TOC's decision at the May 30, 2012 quarterly meeting.
- ^a Average stage is calculated using stage elevations at stations 1-7, 1-8C, and 1-9 for a given sampling date.
- ^b N/A denotes that the level was not applicable because the average stage was less than 15.42 feet.

^{*}The geometric-mean was greater than the long-term level

EVERGLADES NATIONAL PARK

Shark River Slough

Background

The Settlement Agreement/Consent Decree (1995) specified that interim and long-term TP concentration limits for discharges into the Everglades National Park (ENP) (Figure 4) through Shark River Slough be met by October 1, 2003, and December 31, 2006, respectively. It was specified that the TP concentrations be presented as 12-month flow-weighted means. Only the TP concentrations for the water year ending September 30 are evaluated for compliance with the Consent Decree limits (Appendix D). The long-term TP concentration limit for inflows to Shark River Slough through structures S12A, S12B, S12C, S12D, and S333 represents the concentrations delivered during the Outstanding Florida Waters baseline period of March 1, 1978, to March 1, 1979, and is adjusted for variations in flow. Inflow concentrations of TP through Shark River Slough are compared to the interim and long-term limits at the end of each water year (October 1 through September 30) from 1991 to 2012 (Figure 5). The 12-month flow-weighted mean TP concentration (8.9 ppb) was higher than the long-term limit (8.8 ppb) for Water Year 2012 ending on September 30, 2012.

Reporting Period Update

Table 3 presents the 12-month flow-weighted mean concentrations for each month with the corresponding long-term TP concentration limits calculated using the 12-month period flow. Routine monitoring was changed to weekly for all Shark River Slough sites beginning in August 2007. In accordance with Appendix A of the Consent Decree, only the every-other-week grab concentration data were used for the flow-weighted mean calculations from October 2007 forward¹. Weekly TP data for each station for the past 12 months are provided in **Appendix B**. For the 12-month periods ending in October, November and December 2012, the provisional 12-month flow-weighted mean TP concentrations were 8.3, 7.9, and 7.8 ppb, respectively. The provisional 12-month long-term limits, based on the total flow into Shark River Slough and calculated using provisional flow data for the reporting quarter, were 8.4, 8.5, and 8.5 ppb, respectively for the periods.

The Consent Decree stipulates that the percentage of flow-weighted mean TP concentrations greater than 10 ppb from each sampling event in any 12-month period must not exceed a guideline value based on flow into Shark River Slough for the same 12-month period. For the 12-month periods ending October, November and December 2012, the sampling event TP concentration greater than 10 ppb was 28.0 percent. The observed percentages of the sampling event flow-weighted mean TP concentrations greater than 10 ppb were lower than the guideline for the 12-month periods ending in October, November and December 2012 (**Table 3**).

¹ S12A and S333 are sampled weekly if flowing, otherwise monthly. S12B, S12C, and S12D are sampled weekly if flowing.

7

The 12-month flow-weighted mean TP concentrations and the flow-weighted mean TP concentrations for individual sampling events are presented in **Figure 6**.

The daily flows through the individual Shark River Slough structures from January 2010 through December 2012 are presented in Figures 7 and 9. The stage in Water Conservation Area 3A (WCA-3A) remained in Zone A (flood releases) throughout the reporting quarter.

Based on provisional flow data, a total of 375,509 acre-feet of water was discharged through the S12 structures and only 32,372 acre-feet of water was discharged through S333 during the quarter. About 74 percent (24,108 acre-feet) of the water through S333 was discharged through S334 during the quarter (**Figure 8**).

For additional information on the WCA-3A regulation schedule, please refer to the U.S. Army Corps of Engineers (USACE) – Jacksonville District's website².

The relationship between the sum of the daily flows at Shark River Slough structures and corresponding flow-weighted mean TP concentrations for individual sampling events is presented in **Figure 10**. The average of the flow-weighted mean TP concentrations was 6.1 ppb in the fourth quarter (**Figure 10**).

-

² http://w3.saj.usace.army.mil/h2o/plots.htm




Figure 4. Everglades National Park flow structures.




Figure 5. The 12-month flow-weighted mean total phosphorus concentrations at inflows to the Everglades National Park through Shark River Slough at the end of each water year compared to the interim and long-term total phosphorus limits. The 12-month flow-weighted mean TP concentration for Water Year 2012 exceeded the long-term limit when a re-sampled datum was not included in the 12-month flow-weighted mean calculation. The exceedance is currently being evaluated by the TOC. The Water Year 2008 concentration was above the long-term limit but the TOC at the March 1, 2011, quarterly meeting determined substantial evidence indicates this exceedance was due to error.

Table 3. Shark River Slough total phosphorus compliance tracking

12-Month	Total How	How-Weighted Mean TP	Long-Term Limit (ppb)		ampling Events han 10 ppb
Period	(kac-ft)	Concentration (ppb)	<i>Effective</i> 12/31/2006	Guideline	Observed
Feb 2009 - Jan 2010	656.3	9.9	9.7	50.4	30.4
Mar 2009 - Feb 2010	682.1	9.8	9.5	49.6	34.8
Apr 2009 - Mar 2010	733.9	9.7	9.3	48.2	34.8
May 2009 - Apr 2010	790.9	9.9	9.0	46.6	37.5
Jun 2009 - May 2010	869.0	9.9	8.6	44.6	42.3
Jul 2009 - Jun 2010	861.2	9.0	8.6	44.8	42.3
Aug 2009 - July 2010	859.2	8.8	8.6	44.8	42.3
Sep 2009 - Aug 2010	842.5	8.8	8.7	45.3	48.1ª
Oct 2009 - Sep 2010	809.9	8.9	8.9	46.1	50.0 °
Nov 2009 - Oct 2010	757.3	9.0	9.1	47.5	50.0 ^a
Dec 2009 - Nov 2010	742.5	9.0	9.2	47.9	46.2
Jan 2010 - Dec 2010	739.1	9.0	9.2	48.0	46.2
Feb 2010 - Jan 2011	730.5	9.1	9.3	48.3	48.0
Mar 2010 - Feb 2011	695.2	9.1	9.5	49.3	45.8
Apr 2010 - Mar 2011	645.6	9.2	9.7	50.7	54.2ª
May 2010 - Apr 2011	585.0	9.0	10.1	52.5	54.2ª
Jun 2010 - May 2011	526.0	8.8	10.4	54.4	54.2
Jul 2010 - Jun 2011	484.1	8.6	10.6	55.8	52.2
Aug 2010 - Jul 2011	399.7	8.0	11.1	58.6	47.6
Sep 2010 - Aug 2011	311.9	7.4	11.6	61.7	47.4
Oct 2010 - Sep 2011	247.0	9.2	12.0	64.1	57.9
Nov 2010 - Oct 2011	235.6	11.3	12.1	64.6	68.4ª
Dec 2010 - Nov 2011	373.8	10.0	11.3	59.5	68.4ª
Jan 2011 - Dec 2011	479.2	9.3	10.7	55.9	68.4ª
Feb 2011 - Jan 2012	506.3	9.2	10.5	55.0	61.9ª
Mar 2011 - Feb 2012	513.2	9.3	10.5	54.8	63.6ª
Apr 2011 - Mar 2012	503.4	9.3	10.5	55.1	57.1 ^a
May 2011 - Apr 2012	498.5	9.2	10.6	55.3	55.0
Jun 2011 - May 2012	508.6	9.5	10.5	55.0	50.0
Jul 2011 - Jun 2012	561.7	9.6	10.2	53.3	52.4
Aug 2011 - July 2012	644.9	9.3	9.7	50.7	45.8
Sep 2011 - Aug 2012	731.1	9.2	9.3	48.2	40.0
Oct 2011 - Sep 2012	818.3	8.9 (8.8*)	8.8	45.9	36.0
Nov 2011 - Oct 2012	909.8	8.3	8.4	43.6	28.0
Dec 2011 - Nov 2012	882.4	7.9	8.5	44.3	28.0
Jan 2011 - Dec 2012	873.1	7.8	8.5	44.5	28.0

- kac-ft = thousand acre feet.
- ppb = parts per billion. Values are actually in μg/L (micrograms per liter), which, for the purposes of this report, is equivalent to ppb.
- Compliance is evaluated annually based on the 12-month flow-weighted mean (FWM) TP concentration for the federal water year ending on September 30. The compliance periods are shown as highlighted rows with bold, italicized text.
- ^a Value exceeded the guideline percentage.
- * 12-month FWM calculation including the re-sampled datum (7 ppb) at S12D on December 8, 2011 (see Table B-1) is presented in parentheses as additional information. The exceedance is currently being evaluated by the TOC.




Figure 6. The 12-month flow-weighted mean TP concentrations in inflows to Everglades National Park through Shark River Slough at the end of each month and the flow-weighted mean concentration for each sampling event. There are no sampling event values for some months because there was little or no flow in those periods.




Figure 7. Daily flows at Shark River Slough structures as a stacked sum of five inflows.




Figure 8. Daily flows at structures S333 and S334.




Figure 9. Daily flows at individual inflow structures to Shark River Slough. This figure includes most of the data illustrated in Figures 7 and 8.




Figure 10. Flow at Shark River Slough structures on the day of sampling and the corresponding flow-weighted mean total phosphorus concentrations for individual sampling events.

Taylor Slough and Coastal Basins

Background

Under the Consent Decree, a single TP long-term limit of 11 ppb, to be met by December 31, 2006, was set for the two points of inflow to Taylor Slough (S332 and S175) and the inflow point to the Coastal Basins (S18C) (see **Appendix C**). The 12-month flow-weighted mean concentrations have consistently been lower than the long-term limit of 11 ppb.

Inflow TP concentrations to the ENP through Taylor Slough and Coastal Basins are compared to the 11 ppb limit at the end of each water year using data from both the old (S175, S332, S18C) and new (S174, S332D, S18C) combinations of structures (**Figure 11**). The narrow bars in **Figure 11** represent the 12-month flow-weighted mean TP concentrations from S332, S175, and S18C for water years 1991 through 2002. The wider bars for water years 1999 through 2012 represent the new combination of structures.

TP and flow data from both sets of structures presented in prior editions of this report through December 2001 (April 2002 report) showed that, beginning in October 2000, the 12-month moving total flow for S332D, S174, and S18C was consistently greater than flow at S332, S175, and S18C. There was also a shift in flow-weighted mean TP concentration data whereby S332D, S174, and S18C concentrations became equal to and then consistently lower than the concentrations at S332, S175, and S18C. These changes reflected the switch from S332 to S332D for water delivery to Taylor Slough between July 3 and July 5, 2000. Consequently, as of the July 2002 report, only S332D, S174, and S18C data are presented for monthly tracking of data in **Figure 11**. However, almost no flow passed through S174 from March 2006 to September 2007. The site was plugged in September 2007, preventing any additional flow.

The flow-weighted mean TP concentration was below the long-term limit for the 12-month period ending on September 30, 2012. Therefore, Taylor Slough and Coastal Basins TP concentration was in compliance for Water Year 2012.

Reporting Period Update

Figure 12 presents the 12-month and individual sampling event flow-weighted mean TP concentrations at the S332D and S18C structures. All TP grab sample concentrations taken on positive flow days reported for surface water monitoring at the sites were used for the compliance calculations.

The daily flows into ENP through S332D, S174, and S18C are presented in Figures 13 and 14.

For the combined flow through S332D and S18C, the 12-month flow-weighted mean TP concentrations for the periods ending in October, November, and December 2012 were 5.3, 5.1, and 5.1 ppb, respectively (**Table 4**).

The Consent Decree stipulates that the percent of flow-weighted mean TP concentrations greater than 10 ppb from each sampling event in any 12-month period must not exceed a fixed

guideline of 53.1 percent. There were no sampling event TP concentrations greater than 10 ppb for the 12-month periods ending October, November, and December 2012.

Figure 15 shows the relationship between the daily inflows and the corresponding flow-weighted mean TP concentrations for each sampling event. The sampling event flow-weighted mean concentrations generally remained very low. The average of the flow-weighted mean TP concentrations was 4.6 ppb in the fourth quarter.

The USACE authorized the C-111 project in 1995 to restore more natural hydrologic conditions in Taylor Slough and to maintain flood protection to the east of the L31N and C-111 canals.

The original project facilities consisted of pump stations (S332B, S332C, and S332D), detention cells (Cell 1 through Cell 5), a connector cell between Cell 2 and Cell 3, a flow-way cell originating at Berm 3 of Cell 5, and four diversion structures (DS1 through DS4). Upon completion of a USACE construction project in 2009, an interconnected detention system now exists, starting at S332B west discharge and continuing to the S332D high head cell.




Figure 11. The 12-month flow-weighted mean total phosphorus concentrations in inflows to Everglades National Park through Taylor Slough and Coastal Basins at the end of each water year compared to the 11 ppb long-term total phosphorus limit.




Figure 12. The 12-month flow-weighted mean total phosphorus concentrations in inflows to Everglades National Park through Taylor Slough and Coastal Basins at the end of each month and the flow-weighted mean total phosphorus concentration for each sampling event.




Figure 13. Daily flows measured in cubic feet per second into Everglades National Park as a stacked sum of Taylor Slough (structures S332D + S174) and Coastal Basins (structure S18C). Structure S174 was plugged in September 2007, and is no longer used.




Figure 14. Daily flows at individual Coastal Basins (S18C) and Taylor Slough (S332D + S174) structures into the Everglades National Park.

Table 4. Taylor Slough and Coastal Basins total phosphorus compliance tracking.

12-Month	Total How	How-Weighted Mean TP	Long-Term Limit (ppb)		ampling Events han 10 ppb
Period	(kac-ft)	Concentration (ppb)	Effective 12/31/2006	Guideline	Observed
Feb 2009 - Jan 2010	395.0	6.2	11.0	53.1	2.3
Mar 2009 - Feb 2010	414.5	6.2	11.0	53.1	2.1
Apr 2009 - Mar 2010	418.5	6.2	11.0	53.1	2.1
May 2009 - Apr 2010	430.6	6.2	11.0	53.1	2.0
Jun 2009 - May 2010	441.7	5.6	11.0	53.1	0.0
Jul 2009 - Jun 2010	428.2	5.5	11.0	53.1	0.0
Aug 2009 - Jul 2010	413.2	5.5	11.0	53.1	0.0
Sep 2009 - Aug 2010	404.8	5.4	11.0	53.1	0.0
Oct 2009 - Sep 2010	377.5	5.4	11.0	53.1	0.0
Nov 2009 - Oct 2010	349.1	5.5	11.0	53.1	0.0
Dec 2009 - Nov 2010	328.9	5.5	11.0	53.1	0.0
Jan 2010 - Dec 2010	283.3	5.4	11.0	53.1	0.0
Feb 2010 - Jan 2011	273.0	5.4	11.0	53.1	0.0
Mar 2010 - Feb 2011	253.5	5.4	11.0	53.1	0.0
Apr 2010 - Mar 2011	246.6	5.3	11.0	53.1	0.0
May 2010 - Apr 2011	235.2	5.3	11.0	53.1	0.0
Jun 2010 - May 2011	215.0	5.2	11.0	53.1	7.1
Jul 2010 - Jun 2011	183.4	5.2	11.0	53.1	8.1
Aug 2010 - Jul 2011	156.3	5.3	11.0	53.1	8.8
Sep 2010 - Aug 2011	126.4	5.5	11.0	53.1	9.4
Oct 2010 - Sep 2011	111.4	5.6	11.0	53.1	9.4
Nov 2010 - Oct 2011	134.6	6.3	11.0	53.1	12.1
Dec 2010 - Nov 2011	157.9	6.4	11.0	53.1	12.5
Jan 2011 - Dec 2011	170.2	6.3	11.0	53.1	11.1
Feb 2011 - Jan 2012	169.5	6.4	11.0	53.1	10.0
Mar 2011 - Feb 2012	170.2	6.4	11.0	53.1	9.3
Apr 2011 - Mar 2012	169.5	6.4	11.0	53.1	9.5
May 2011 - Apr 2012	170.3	6.3	11.0	53.1	10.0
Jun 2011 - May 2012	193.5	6.1	11.0	53.1	2.5
Jul 2011 - Jun 2012	224.6	6.0	11.0	53.1	2.3
Aug 2011 - July 2012	255.1	5.8	11.0	53.1	2.1
Sep 2011 - Aug 2012	273.0	5.8	11.0	53.1	2.0
Oct 2011 - Sep 2012	290.7	5.7	11.0	53.1	2.0
Nov 2011 - Oct 2012	283.6	5.3	11.0	53.1	0.0
Dec 2011 - Nov 2012	273.0	5.1	11.0	53.1	0.0
Jan 2012 - Dec 2012	279.6	5.1	11.0	53.1	0.0

- kac-ft = thousand acre feet.
- ppb = parts per billion. Values are actually in μ g/L (micrograms per liter), which, for the purposes of this report, is equivalent to ppb.
- Compliance is evaluated annually based on the 12-month flow-weighted mean TP concentration for the federal water year ending on September 30. The compliance periods are shown as highlighted rows with bold, italicized text.




Figure 15. Flow from Taylor Slough and Coastal Basins structures (S332D + S174 and S18C) on the day of sampling and the corresponding flow-weighted mean total phosphorus concentrations for individual sampling events.

APPENDIX A

MONTHLY TOTAL PHOSPHORUS CONCENTRATION DATA FOR THE ARTHUR R. MARSHALL LOXAHATCHEE NATIONAL WILDLIFE REFUGE

Total phosphorus (TP) concentration data used in this report can be directly retrieved from the SFWMD DBHYDRO database by copying and pasting the following link it to the address field of a web browser:

 $http://www.sfwmd.gov/dbhydroplsql/water_quality_data.report_full?v_where_clause=where e+station_id+like+('LOX%25')+and+station_id+not+like+('LOXA%25')+and+test_number+=+25+and+date_collected+>=+'01-OCT-2012'+and+date_collected+<+'01-JAN-2013'+and+sample_type_new+=+'SAMP'&v_target_code=file_csv$

The link above generates only data that have not been qualified. Qualified water quality data must be retrieved interactively via the DBHYDRO Browser.

Stage data for stations 1-7, 1-8C, and 1-9 from the reporting quarter can be retrieved by copying and pasting the following link it to the address field of a web browser:

http://www.sfwmd.gov/dbhydroplsql/web_io.report_process?v_period=uspec&v_start_date =20121001&v_end_date=20121231&v_report_type=format7&v_target_code=file_csv&v_run_m ode=onLine&v_js_flag=Y&v_dbkey=FE775/FE776/FE777

Settlement Agreement Report October-December 2012

Table A-1. Refuge monthly TP data (ppb).

Month-Year	LOX3	LOX4	LOX5	LOX6	LOX7	LOX8	LOX9	LOX10	LOX11	LOX12	LOX13	LOX14	LOX15	LOX16
Jan-2010	7	5	7	4	5	8	5	5	5	5	5	4	5	4
Feb-2010	10	8	11	5	8	8	6	7	5	6	5	6	6	6
Mar-2010	11	7	10	2	7	8	6	6	4	6	5	5	6	6
Apr-2010	9	8	10	5	9	10	6	6	6	6	6	7	6	9
May-2010	9	7	14	7	8	8	6	5	9	8	7	8	7	10
Jun-2010	8	8	7	9	7	7	5	5	7	7	6	7	10	12
Jul-2010	9	9	9	8	7	6	5	7	5	6	6	6	7	8
Aug-2010				6	7	6	5	6	6	6		5	7	7
Sep-2010	6	10	6	5	6	6	6	6	6	8	6	6	7	8
Oct-2010	5	17	5	6	5	7	5	7	6	7	5	5	6	7
Nov-2010	5	11	6	6	7	9	5	6	7	7	7	7	5	6
Dec-2010	4	7 (J)	5	3	4	5	4	4	3	5	4	4	4	4
Jan-2011		8		5	6	8			4	5	4	4	4	5
Feb-2011		9		5	7	8			5	6	6	5	4	7
Mar-2011					12	15			7	6	7	7	7	7
Apr-2011						48			10	7	7	12	8	11
May-2011														
Jun-2011														
Jul-2011		30			12					20			23	
Aug-2011				12	13	17			8	11		14	8	12
Sep-2011		13		10	6	10	12	10	8	11		9	8	13
Oct-2011		11		6	5	8	6	10	7	8		7	7	8
Nov-2011	7	18	7	6	7	10	7	8	9	7	9	6	6	7
Dec-2011	7 (J)	10 (J)	6 (J)	5	7 (J)	9 (J)	8 (J)	8 (J)	7	6	8	6	5	7
Jan-2012	5	9	6	5	6	8	3	8	7	6	6	6	6	6
Feb-2012		10	9	5	9	10	9	11	7	6	6	7	5	7
Mar-2012		6		5	8	10			6	5	7	6	5	7
Apr-2012		11		5	10	13			8	8	7	7	6	9
May-2012		9		5	8	8	9	9	7	6	7	6	6	8
Jun-2012	10	9	9	6	9	10	10	7	7	6	7	5	5	6
Jul-2012	5	7	6	5	5	7	6	6	5	6	8	5	7	7
Aug-2012	5	6	7	4	5	5	6	6	5	7		4	6	6
Sep-2012	8	15	8	6	8	7	7	7	5	6	5	7	6	7
Oct-2012	5	7	7	8	9	6	9	8	8	9	8	8	8	8
Nov-2012	7	7	7	5	7	8	6	7	7	6	6	4	5	5
Dec-2012	6	6	8	6	8	7	8	8	6	7	6	5	8	6

- --- indicates sample was not collected due to insufficient water depth.
- (J) indicates analyte detected in field blank and/or associated sample.

APPENDIX B

WEEKLY GRAB TOTAL PHOSPHORUS CONCENTRATION DATA FOR SHARK RIVER SLOUGH

Total phosphorus (TP) concentration data used in this report can be directly retrieved from the SFWMD DBHYDRO database by copying and pasting the following link it to the address field of a web browser:

http://www.sfwmd.gov/dbhydroplsql/water_quality_data.report_full?v_where_clause=wher e+station_id+in+('S12A','S12B','S12C','S12D','S333')+and+test_number+=+25+and+collect_meth od+=+'G'+and+date_collected+>=+'01-OCT-2012'+and+date_collected+<+'01-JAN-2013'+and+sample_type_new+=+'SAMP'&v_target_code=file_csv

The link generates only data that have not been flagged. Flagged water quality data must be retrieved interactively via DBHYDRO Browser.

Flow data data used in this report for the reporting quarter for stations at Shark River Slough, S12A, S12B, S12C, S12D, S333, and S334, can be retrieved by copying and pasting the following link it to the address field of a web browser:

http://www.sfwmd.gov/dbhydroplsql/web_io.report_process?v_period=uspec&v_start_date =20121001&v_end_date=20121231&v_report_type=format7&v_target_code=file_csv&v_run_m ode=onLine&v_js_flag=Y&v_dbkey=FE771/03626/03632/FE774/15042/FB752

Table B-1. Weekly Grab TP Concentration Data for Shark River Slough (ppb).

Date	S12A	S12B	S12C	S12D	S333	Remarks
	11	3126	5	10	9	Compliance data
01/04/2012						
01/11/2012	10			8 7	8	N/A
01/18/2012	12				8	Compliance data
01/24/2012	8			7	8	N/A
01/31/2012	14			8	9	Compliance data
02/07/2012	14			7	11	N/A
02/14/2012	11			8	8	Compliance data
02/22/2012	12			9	9	N/A
02/28/2012	14			28	10	Compliance data
03/06/2012	12			9	11	N/A
03/13/2012	18			9	11	Compliance data
03/20/2012	18			10	8	N/A
03/27/2012	22			13	10	Compliance data
04/04/2012	16 (J)				10 (J)	N/A
04/13/2012	20				12	Compliance data
04/17/2012	19				12	N/A
04/24/2012	18				11	Compliance data
05/01/2012	16				12	N/A
05/09/2012	32			9	9	Compliance data
05/15/2012	14			12	12	N/A
05/22/2012	12			15	17	Compliance data
05/30/2012	26			**	13	N/A
06/05/2012	21			9	13	Compliance data
06/12/2012	18			9	19	N/A
	16			8	17	Compliance data
06/19/2012	20			8	13	•
06/26/2012						N/A
07/02/2012	20			8	10	Compliance data
07/11/2012	14				9	N/A
07/17/2012	7	5	7	8	8	Compliance data
07/24/2012	6	6	9	9	10	N/A
07/31/2012	6	7	9	9	9	Compliance data
08/07/2012	6	7	10	12	10	N/A
08/14/2012	7	8	9	***	9	Compliance data
08/21/2012	9	8	8	***	8	N/A
08/29/2012	5	7	8	12	11	Compliance data
09/04/2012	8	10	8	10	9	N/A
09/11/2012	7	6	9	17	7	Compliance data
09/18/2012	7	7	8	14	9	N/A
09/25/2012	5	5	6	9	7	Compliance data
10/01/2012	6	6	7	9	8	N/A
10/08/2012	6	5	7	9	9	Compliance data
10/15/2012	4	4	6	7	8	N/A
10/22/2012	6	4	6	8	7	Compliance data
10/29/2012	5	4	5	8	8	N/A
11/05/2012	9	5	4	7	6	Compliance data
11/13/2012	7	5	3	6	6	N/A
11/19/2012	7	4	4	7	5	Compliance data
11/26/2012	8	5	4	7	6	N/A
12/03/2012	8	4	4	8	7	Compliance data
12/10/2012	11	6	4	7	7	N/A
	9	7	5	8	7	Compliance data
12/17/2012						'
12/26/2012	7	5	3	6	7	N/A

⁻⁻⁻ indicates water sample was not collected because the spillway gates were closed at the time of the site visit.

⁽J) indicates analyte detected in field blank and/or associated sample.

[&]quot;Compliance data" indicates bi-weekly sampling data used for consent decree calculation.

[&]quot;N/A" indicates bi-weekly sampling data presented for informational purposes only and not used for consent decree calculation.

^{**} Water sample taken at S12D on May 30, 2012, was misplaced and could not be analyzed.

^{***} Water samples were not collected at S12D on the August 14 and August 21, 2012, trips because the site was not accessible due to structure maintenance activities being conducted by the USACE.

APPENDIX C

WEEKLY GRAB TOTAL PHOSPHORUS CONCENTRATION DATA FOR TAYLOR SLOUGH AND COASTAL BASINS

Total phosphorus (TP) concentration data used in this report can be directly retrieved from the SFWMD DBHYDRO database by copying and pasting the following link it to the address field of a web browser:

 $http://www.sfwmd.gov/dbhydroplsql/water_quality_data.report_full?v_where_clause=where + station_id+in+('S332DX','S18C','S174')+and+test_number+=+25+and+collect_method+=+'G'+and+date_collected+>=+'01-OCT-2012'+and+date_collected+<+'01-JAN-2013'+and+sample_type_new+=+'SAMP'&v_target_code=file_csv$

The link generates only data that have not been flagged. Flagged water quality data must be retrieved interactively via DBHYDRO Browser.

Daily mean flow data for stations at Taylor Slough, S332D and S174, and Coastal Basins, S18C, during the reporting quarter can be retrieved by copying and pasting the following link to the address field of a web browser:

http://www.sfwmd.gov/dbhydroplsql/web_io.report_process?v_period=uspec&v_start_date =20121001&v_end_date=20121231&v_report_type=format7&v_target_code=file_csv&v_run_m ode=onLine&v_js_flag=Y&v_dbkey=15760/TA413/15769

Table C-1. Weekly Grab TP Concentration Data for Taylor Slough and Coastal Basins (ppb).

Date	S332DX	S18C
01/03/2012	7	6
01/09/2012		4
01/10/2012	5	
01/17/2012	6	5
01/23/2012	5	4
01/30/2012	6	4
02/06/2012	6	3
02/13/2012	5	4
02/21/2012	6	5
02/27/2012	5	4
03/05/2012	7	4
03/12/2012	5	3
03/19/2012	6	4
03/26/2012	6	3
04/02/2012		4
04/03/2012	9	
04/09/2012	8	4
04/16/2012	10	4
04/23/2012	7	7
04/30/2012	6	6
05/08/2012	6	4
05/14/2012	5	3
05/21/2012	6	5
05/29/2012	6	4
06/04/2012	6	4
06/11/2012	6	4
06/18/2012	6	6
06/25/2012	7	6
07/02/2012	5	9
07/10/2012	6	
07/11/2012		3
07/16/2012	6	4
07/23/2012	6	4
07/30/2012	6	5

Date	S332DX	S18C
08/06/2012	6	3
08/13/2012	5	5
08/20/2012	5	6
08/28/2012	5	6
09/04/2012	5	5
09/10/2012	7	4
09/17/2012	5	6
09/24/2012	6	4
10/01/2012	-	4
10/02/2012	6	
10/08/2012	-	4
10/09/2012	6	
10/15/2012		3
10/16/2012	6	
10/22/2012		4
10/23/2012	6	
10/29/2012		4
10/30/2012	6	
11/05/2012		4
11/06/2012	6	
11/13/2012	4	3
11/19/2012		3
11/20/2012	6	
11/26/2012		3
11/27/2012	5	
12/03/2012		4
12/04/2012	6	
12/10/2012	5	
12/11/2012		3
12/17/2012	5	
12/18/2012		3
12/27/2012		3

Note: -- indicates water sample was not collected.

APPENDIX D

CALCULATION METHODS

Long Term Marsh Concentration Levels for Loxahatchee National Wildlife Refuge

Long Term Marsh Concentration Levels:

$$C = 10.7172 - 0.541156S + 1.372\sqrt{7.5819 - 0.9310S + 0.02902216S^2}$$

Terms:

C = the natural log of the geometric mean total phosphorus concentration across 14 marsh stations.

S = average stage measured at gauges CA1-9, CA1-7, and CA1-8C on sampling date (feet).

This equation is applicable over a stage range of 15.42 to 17.14 feet. If the stage on any sampling date exceeds 17.14 feet, a stage of 17.14 feet should be used in calculating the long term concentration levels. The equation shall not apply to dates when the average stage is less than 15.42 feet.

(1991 Settlement Agreement entered as a Consent Decree in 1992 and modified in 1995, Exhibit B, Appendix B, Attachment II, page B-7)

Discharge Limits and OFW Standards for Shark River Slough

Interim Discharge Limit:

$$C = 11.16 - 0.00465Q + 1.397\sqrt{6.377 - 0.00591Q + 0.00000436Q^2}$$

Long-Term Discharge Limit & OFW Standard:

$$C = 11.38 - 0.00538Q + 1.397\sqrt{2.493 - 0.00231Q + 0.00000170Q^2}$$

Frequency Exceedance:

$$F = 48.411 - 0.02896Q + 1.397\sqrt{330.1 - 0.3071Q + 0.0002254Q^2}$$

Terms:

Water Year = October through September

Q = total inflow to Shark River Slough for water year, S-12s + S-333 + any additional inflow from the WCAs established in the future, thousand acre-ft/yr (Kac-ft/yr).

C = limit on maximum flow-weighted-mean inflow concentration for any Water Year, composite of all inflows to Shark Slough (ppb).

F = exceedance for maximum frequency (percent) of inflow concentrations exceeding 10 ppb, computed from the time series of concentrations composited across all inflow structures on each sampling date with positive flow in a given Water Year.

The range of flow (Q) used in deriving the limits is 117 to 1061 Kac-ft/yr. If the total flow for any water year exceeds 1061 Kac-ft/yr, a flow of 1061 Kac-ft/yr should be used in calculating the discharge limits.

(1991 Settlement Agreement entered as a Consent Decree in 1992 and modified in 1995, Exhibit B, Appendix A, Attachment I, page A-5)

Note: Additional inflows, currently, comprises the discharges through S334, S355A and S335B, and S356. The latest TOC approved methodology to incorporate these additional inflows was documented in the Shark River Slough section of the Settlement Agreement July – September 2006 Report (dated November 9, 2011).

Discharge Limits and OFW Standards for Taylor Slough and Coastal Basins

Long-Term Flow-Weighted Discharge Limit & OFW Standard = 11.0 ppb

Frequency Exceedance:

Frequency of values > 10 ppb must be less than 53.1%.

Terms:

Limits are defined on a Water Year basis, October through September.

Basin flow is the total flow through structures S-332, S-175, S-18C, plus any new release points from this basin established in the future, thousand acre-ft/yr (Kac-ft/yr).

Limits apply to the flow-weighted-mean concentration for any Water Year, composite of all inflows to Taylor Slough (S-332) and Coastal Basin (S-18C).

Frequency exceedance is the exceedance for maximum frequency (percent) of inflow concentrations exceeding 10 ppb, computed from the time series of concentrations composited across all inflow structures on each sampling date with positive flow in a given Water Year.

(1991 Settlement Agreement entered as a Consent Decree in 1992 and modified in 1995, Exhibit B, Appendix A, Attachment II, page A-6)

APPENDIX E DOCUMENT REVISIONS

Table E-1. Revisions to this report since initial publication.

Page/Date	Original	Revision/Update
	The finalized flow data for the	
	first nine months for Shark	
Table of Contents	River Slough, was pending at	The Shark River Slough values using the
	the time of the initial publication	final flow data for October 2011 to
Shark River	(Report published on Feb. 13,	September 2012 and provisional flow
Slough Section	2013).	data for October 2012 to December 2012
(pages 7-15, Table	The Shark River Slough Section	is presented in this version as an update
3, Figures 5-10)	of the report did not present any	for Shark River Slough Section of the
	data, figures or tables for the	report.
Appendix B	fourth quarter of Water Year	
(pageB-1 and B-2)	2012. Only TP concentration	Updated for flow DBKEYs.
	data were presented in	
	Table B-1.	