Settlement Agreement Report

Third Quarter July - September 2009

Revisions were made to this document on December 16, 2009. (See Figure 7 on page 13.)

Prepared for the Technical Oversight Committee December 08, 2009



Prepared by:

Cheol Mo, Violeta Ciuca, and Pamela Lehr Restoration Sciences Department South Florida Water Management District 3301 Gun Club Road West Palm Beach, FL 33406

PURPOSE

The South Florida Water Management District has prepared this report to provide a quarterly update to the Technical Oversight Committee on the compliance status with total phosphorus levels or limits defined in the 1991 Settlement Agreement entered as a Consent Decree in 1992 and modified in 1995. The areas of interest in this report include the interior marsh stations in Arthur R. Marshall Loxahatchee National Wildlife Refuge and two discharges to Everglades National Park: inflows to Shark River Slough and inflows to Taylor Slough and Coastal Basins.

CONTENTS

Executive Sur	nmary	1
	rshall Loxahatchee National Wildlife Refuge	
	d	
Reporting l	Period Update	3
Everglades N	ational Park	7
	r Slough	
U	ınd	
-	g Period Update	
	igh and Coastal Basins	
	and	
Reportin	g Period Update	16
	APPENDICES	
Appendix A.	Monthly Total Phosphorus Concentration Data for the Arthur R. Marshall Loxahatchee National Wildlife Refuge	۸ 1
Appendix B.	Weekly Grab Total Phosphorus Concentration Data for Shark	A-1
Appendix b.	River Slough	B-1
Appendix C.		
11	Coastal Basins	C-1
Appendix D.	Calculation Methods	D-1
	TABLES	
Table 1.	Total phosphorus compliance, third quarter 2009	1
Table 2.	Loxahatchee National Wildlife Refuge total phosphorus compliance tracking	
Table 3.	Shark River Slough total phosphorus compliance tracking	
Table 4.	Taylor Slough and Coastal Basins total phosphorus compliance tracking	21
	FIGURES	
Figure 1.	Areas of interest	2
Figure 2.	Arthur R. Marshall Loxahatchee National Wildlife Refuge water quality	
	sampling and stage measurement sites	
Figure 3.	(A) Monthly total phosphorus geometric mean concentrations for the Arthur	r
	R. Marshall Loxahatchee National Wildlife Refuge	
	(B) Deviation of monthly geometric mean total phosphorus concentrations	F
	with calculated long-term levels	5

Figure 4.	Everglades National Park flow structures	9
Figure 5.	The 12-month flow-weighted mean total phosphorus concentrations at	
<u> </u>	inflows to the Everglades National Park through Shark River Slough at	
	the end of each water year	10
Figure 6.	The 12-month flow-weighted mean TP concentrations in inflows to Everglade	es
	National Park through Shark River Slough at the end of each month and the	
	flow-weighted mean concentration for each sampling event	12
Figure 7.	Daily flows into Shark River Slough as a stacked sum of five structures	13
Figure 8.	Daily flows at structures S333 and S334.	13
Figure 9.	Daily flows at individual inflow structures to Shark River Slough	14
Figure 10.	Total daily flow at Shark River Slough structures and the corresponding	
_	flow-weighted mean total phosphorus concentrations for individual	
	sampling events	15
Figure 11.	The 12-month flow-weighted mean total phosphorus concentrations in inflow	
	to Everglades National Park through Taylor Slough and Coastal Basins	18
Figure 12.	The 12-month flow-weighted mean total phosphorus concentrations in inflow	7S
	to Everglades National Park through Taylor Slough and Coastal Basins at	
	the end of each month and the flow-weighted mean total phosphorus	
	concentration for each sampling event	18
Figure 13.	Daily flows into Everglades National Park through Taylor Slough as a stacked	1
	total of structures S332D + S174 and S18C	19
Figure 14.	Daily flows at individual Taylor Slough structures (S332S + S174 and S18C)	
	into Everglades National Park.	20
Figure 15.	Total daily flow from Taylor Slough structures (S332D + S174 and S18C) and	
	the corresponding flow-weighted mean total phosphorus concentrations for	
	individual sampling events.	. 212

ACRONYMS AND ABBREVIATIONS

ENP Everglades National Park

FDEP Florida Department of Environmental Protection

IOP Interim Operational Plan

kac-ft thousand acre-feet

NGVD 29 National Geodetic Vertical Datum of 1929

OFW Outstanding Florida Waters

ppb parts per billion

Refuge Arthur R. Marshall Loxahatchee National Wildlife Refuge

SFWMD South Florida Water Management District

STA Stormwater Treatment Area TOC Technical Oversight Committee

TP total phosphorus μg/L micrograms per liter

USACE United States Army Corps of Engineers

WCA Water Conservation Area

EXECUTIVE SUMMARY

This report fulfills the South Florida Water Management District's reporting requirements under the 1991 Settlement Agreement, entered as a Consent Decree in 1992 and modified in 1995, for the third quarter of 2009 (July – September 2009). Total phosphorus (TP) compliance highlights for this period are summarized below for the Arthur R. Marshall Loxahatchee National Wildlife Refuge (Refuge) and Everglades National Park, including Shark River Slough, Taylor Slough, and Coastal Basins (**Table 1** and **Figure 1**):

- **Refuge:** The geometric mean TP concentrations were below the long-term levels in July, August, and September 2009.
- **Shark River Slough:** The flow-weighted mean TP concentration was the same as the long-term limit for the 12-month period ending on September 30, 2009. Therefore, Shark River Slough TP concentration was in compliance for federal water year 2009.
- Taylor Slough and Coastal Basins: The flow-weighted mean TP concentration was below
 the long-term limit for the 12-month period ending on September 30, 2009. Therefore, the TP
 concentration for Taylor Slough and Coastal Basins was in compliance for federal water year
 2009.

Table 1. Total phosphorus compliance, third quarter 2009.

Month	n		etric Mean ntration (ppb)	Long-ter (pp			nn Stage IGVD 29)		Number of Samples
Arthur R. M	larshall L	oxahatche	National Wildli	ife Refuge					
Jul-200)9		7.4	9.	8		16.47		14
Aug-200	09		8.8	11	.5		16.16		14
Sep-200	09		8.1	8.	9		16.68		14
12-Month Period		l Flow	12-Mo Flow-weigh		Long-teri				npling Events an 10 ppb
Ending	(ка	c-ft)	TP Concentra	ation (ppb)	(ppl	D)	Guideline	0	Observed
Everglades	National	Park – Sha	rk River Slough						
Jul-2009	10	47.8	8.3	3	7.7	7	40.4		36.4
Aug-2009	10:	22.2	8.1		7.8	3	41.0		27.3
Sep-2009	94	18.0	8.2	2	8.2	2	42.7		26.1
Everglades	National	Park – Tay	lor Slough and	Coastal Basin	s				
Jul-2009	38	88.8	6.5	5	11.	0	53.1		1.9
Aug-2009	39	8.6	6.3	3	11.	0	53.1		1.9
Sep-2009	41	11.4	6.2	2	11.	0	53.1	·	2.2

Notes:

- ppb = parts per billion. Values are actually in μg/L (micrograms per liter), which, for the purposes of this report, is equivalent to ppb
- ft NGVD 29 = elevation in feet related to the National Geodetic Vertical Datum of 1929.
- kac-ft = thousand acre feet.
- Compliance for inflows to Everglades National Park (Shark River Slough, Taylor Slough and Coastal Basins) is evaluated annually based on the 12-month flow-weighted mean TP concentration for the federal water year ending in September 30 (highlighted rows with bold, italicized text).




Figure 1. Areas of interest.

ARTHUR R. MARSHALL LOXAHATCHEE NATIONAL WILDLIFE REFUGE

Background

The 1991 Settlement Agreement ended the Everglades lawsuit and was entered into by the federal government, the State of Florida, and the South Florida Water Management District. The subsequent Consent Decree, as modified in 1995, specified that interim and long-term total phosphorus (TP) concentration levels for the Arthur R. Marshall Loxahatchee National Wildlife Refuge (Refuge) must be met by February 1, 1999, and December 31, 2006, respectively. Both the interim and long-term concentration levels vary monthly because they are calculated as a function of water stage measured at gauging stations 1-7, 1-8C, and 1-9 within the Refuge. The stage range within which the interim and long-term concentration levels are applicable is 15.42 to 17.14 feet relative to the National Geodetic Vertical Datum of 1929 (NGVD 29). The monthly TP concentrations are determined from water samples collected at 14 interior marsh stations, LOX3 through LOX16 (Figure 2). As required in the Consent Decree, the concentrations are converted to a geometric mean, which is compared to the interim and long-term concentration levels. Monthly TP data for each station for the past 36 months are provided in Appendix A. The calculation methods specified in the Consent Decree are provided in Appendix D.

Reporting Period Update

Average stages in the Refuge were 16.47, 16.16, and 16.68 feet in July, August, and September 2009, respectively (**Figure 3** and **Table 2**). The geometric means, calculated from TP concentrations measured in water samples collected in July, August, and September 2009 were 7.4, 8.8, and 8.1 parts per billion (ppb), respectively. The geometric mean TP concentrations were below the long-term level in July, August, and September 2009. Water quality samples were collected at all of the 14 interior marsh stations in each month.





Figure 2. Arthur R. Marshall Loxahatchee National Wildlife Refuge water quality sampling and stage measurement sites.






Figure 3. (A) Monthly total phosphorus geometric mean concentrations for the Arthur R. Marshall Loxahatchee National Wildlife Refuge compared to calculated long-term levels. These calculated long-term total phosphorus levels are adjusted for fluctuations in stage. The long-term levels were not applicable for April 2007 through June 2007 because the average stage was less than 15.42 feet. The geometric mean was greater than the long-term level in October 2007, November 2008, and June 2009.

(B) Deviation of monthly geometric mean total phosphorus concentrations with calculated long-term levels. Values greater than zero suggest that the geometric mean was greater than the long-term level.

Table 2. Loxahatchee National Wildlife Refuge total phosphorus compliance tracking.

Month	Geometric Mean TP Concentration (ppb)	Interim Level (ppb) 2/1/1999-12/30/2006	Long-Term Level (ppb) Effective 12/31/2006	Average Stage ^a (ft NGVD 29)	Number of Samples
Oct-2006	7.5	10.1		16.74	14
Nov-2006	7.4	11.1		16.56	14
Dec-2006	5.6	13.4		16.23	11
Jan-2007	6.9		9.3	16.57	14
Feb-2007	6.2		10.4	16.34	13
Mar-2007	7.3		11.8	16.10	10
Apr-2007	8.0		N/A ^b	15.30	3
May-2007			N/A ^b	14.68	0
Jun-2007	13.6		N/A ^b	15.37	4
Jul-2007	10.4		13.5	15.86	13
Aug-2007	10.1		10.5	16.33	13
Sep-2007	8.5		9.2	16.59	14
Oct-2007	8.4		7.2	17.26	14
Nov-2007	7.0		7.2	17.19	14
Dec-2007	7.9		8.2	16.84	14
Jan-2008	6.3		8.9	16.68	14
Feb-2008	7.6		9.4	16.55	14
Mar-2008	6.7		9.4	16.54	14
Apr-2008	5.8		10.3	16.36	14
May-2008	9.1		14.3	15.76	9
Jun-2008	8.9		14.9	15.68	8
Jul-2008	9.2		10.3	16.37	14
Aug-2008	8.6		10.2	16.39	14
Sep-2008	7.7		8.3	16.81	14
Oct-2008	7.2		7.2	17.42	14
Nov-2008	7.4		7.2	17.22	14
Dec-2008	6.3		7.8	16.95	14
Jan-2009	7.0		8.8	16.68	14
Feb-2009	4.7		10.4	16.35	12
Mar-2009	7.5		13.5	15.86	9
Apr-2009	9.7		14.9	15.69	8
May-2009	11.0		N/A ^b	15.01	1
Jun-2009 ^c	13.2		12.1	16.05	12
Jul-2009	7.4		9.8	16.47	14
Aug-2009	8.8		11.5	16.16	14
Sep-2009	8.1		8.9	16.68	14

Notes:

- ppb = parts per billion. Values are actually in μ g/L (micrograms per liter), which, for the purposes of this report, is equivalent to ppb.
- ft NGVD 29 = elevation in feet related to the National Geodetic Vertical Datum of 1929.
- Highlighted rows with bold, italicized text indicate when an excursion over the long-term level occurred.
- ^a Average stage is calculated using stage elevations at stations 1-7, 1-8C, and 1-9 for a given sampling date.
- ^b N/A denotes that the level was not applicable because the average stage was less than 15.42 feet.
- ^c June 1 and 3, 2009 compliance sampling data only.

EVERGLADES NATIONAL PARK

Shark River Slough

Background

The Settlement Agreement/Consent Decree (1995) specified that interim and long-term TP concentration limits for discharges into the Everglades National Park (ENP) (Figure 4) through Shark River Slough be met by October 1, 2003, and December 31, 2006, respectively. It was specified that the TP concentrations be presented as 12-month flow-weighted means. Only the TP concentrations for the water year ending September 30 are evaluated for compliance with the Consent Decree limits (Appendix D). The long-term TP concentration limit for inflows to Shark River Slough through structures S12A, S12B, S12C, S12D, and S333 represents the concentrations delivered during the Outstanding Florida Waters baseline period of March 1, 1978, to March 1, 1979, and is adjusted for variations in flow. Inflow concentrations of TP through Shark River Slough are compared to the interim and long-term limits at the end of each water year (October 1 through September 30) from 1991 to 2009 (Figure 5).

Reporting Period Update

Table 3 presents the 12-month flow-weighted mean concentrations for each month with the corresponding long-term TP concentration limits calculated using the 12-month period flow. Supplemental water quality samples were collected at some of the Shark River Slough monitoring sites from January 2006 through September 2007, but only the bi-weekly compliance monitoring grab concentration data were used for flow-weighted mean TP concentration calculations. Routine monitoring was changed to weekly for all Shark River Slough sites beginning in October 2007. In accordance with Appendix A of the Consent Decree, only the every-other-week grab concentration data were used for the flow-weighted mean calculations from October 2007 forward¹. For the 12-month periods ending in July, August, and September 2009, the 12-month flow-weighted mean TP concentrations were 8.3, 8.1, and 8.2 ppb, respectively. The long-term limits were 7.7, 7.8, and 8.2 ppb, respectively.

The flow-weighted mean TP concentration was the same as the long-term limit for the 12-month period ending on September 30, 2009. Therefore, Shark River Slough TP concentration was in compliance for federal water year 2009.

-

¹ S12A and S333 are sampled weekly if flowing, otherwise monthly. S12B, S12C, and S12D are sampled weekly if flowing.

The Consent Decree stipulates that the percentage of flow-weighted mean TP concentrations greater than 10 ppb from each sampling event in any 12-month period must not exceed a guideline value based on flow into Shark River Slough for the same 12-month period. For the 12-month periods ending July, August, and September 2009, the sampling event TP concentrations greater than 10 ppb were 36.4, 27.3, and 26.1 percent, respectively. The observed percentages of the sampling event flow-weighted mean TP concentrations greater than 10 ppb were lower than the guidelines for July, August, and September 2009 (**Table 3**). The 12-month flow-weighted mean TP concentrations and the flow-weighted mean TP concentrations for individual sampling events are presented in **Figure 6**.

The daily flows through the individual Shark River Slough structures from October 2006 through September 2009 are presented in **Figures 7** and **9**.

The stage levels at Water Conservation Area (WCA) 3A were in Zone A for the most of reporting period. S12A, S12B and S12C remained closed until July 15, 2009, in accordance with the Interim Operational Plan for Protection of the Cape Sable Seaside Sparrow. S12D and S333 were open during the quarter.

A total of 102,650 acre-feet of water were discharged through S333; 53,259 acre-feet was diverted to S334 during the quarter (**Figure 8**).

For additional information on the WCA-3A regulation schedule, please refer to the U.S. Army Corps of Engineers (USACE) – Jacksonville District's web site http://www.saj.usace.army.mil/h2o/plots/wca3ahp.pdf.

The relationship between the sum of the daily flows at Shark River Slough structures and corresponding flow-weighted mean TP concentrations for individual sampling events is presented in **Figure 10**. Flow and TP concentrations for waters entering the ENP through Shark River Slough had been following an inverse relationship in previous periods (**Figure 10**).




Figure 4. Everglades National Park flow structures.




Figure 5. The 12-month flow-weighted mean total phosphorus concentrations at inflows to the Everglades National Park through Shark River Slough at the end of each water year compared to the interim and long-term total phosphorus limits. The 12-month flow-weighted mean concentrations have met the interim limits since they became effective on October 1, 2003. The 12-month flow-weighted mean TP concentration for the compliance year through September 30, 2009, was 8.2 ppb and was equal to the long-term limit.

Table 3. Shark River Slough total phosphorus compliance tracking.

12-Month	Total Flow	Flow-Weighted Mean TP	Interim Limit (ppb)	Long-Term Limit (ppb)		impling Events han 10 ppb
Period	(kac-ft)	Concentration (ppb)	10/1/2003- 12/30/2006	Effective 12/31/2006	Guideline	Observed
Nov 2005 - Oct 2006	779.6 ^{a,b}	9.5	10.5		46.9	50.0 ^c
Dec 2005 - Nov 2006	642.8 ^{a,b}	10.0	11.1		50.8	50.0
Jan 2006 - Dec 2006	507.9 ^{a,b}	10.3	11.8		55.0	50.0
Feb 2006 - Jan 2007	446.1 ^{a,b}	10.7		10.9	57.0	55.0
Mar 2006 - Feb 2007	442.4 ^b	10.8		10.9	57.2	52.4
Apr 2006 - Mar 2007	456.0 ^b	10.8		10.8	56.7	57.1 ^c
May 2006 - Apr 2007	449.5 ^b	10.8		10.8	56.9	52.6
Jun 2006 - May 2007	445.5 ^b	10.7		10.9	57.1	47.1
Jul 2006 - Jun 2007	447.6 ^b	10.8		10.8	57.0	50.0
Aug 2006 - Jul 2007	444.6 ^b	10.6		10.9	57.1	47.4
Sep 2006 - Aug 2007	401.0	10.3		11.1	58.6	44.4
Oct 2006 - Sep 2007	289.7	9.8		11.8	62.6	37.5
Nov 2006 - Oct 2007	147.9	11.7		12.7	67.9	50.0
Dec 2006 - Nov 2007	121.2	12.6		12.8	68.9	62.5
Jan 2007 - Dec 2007	118.2	12.8		12.8	69.1	71.4 ^c
Feb 2007 - Jan 2008	115.8	12.9		12.9	69.2	69.2
Mar 2007 - Feb 2008	106.8	12.9		12.9	69.5	81.8 ^c
Apr 2007 - Mar 2008	88.9	12.8		13.0	70.2	75.0 ^c
May 2007 - Apr 2008	110.9	12.4		12.9	69.4	78.6 ^c
Jun 2007 - May 2008	142.3	12.2		12.7	68.1	75.0 ^c
Jul 2007 - Jun 2008	153.6	11.8		12.6	67.7	70.6 ^c
Aug 2007 - Jul 2008	227.4	12.4		12.2	64.9	76.5 ^c
Sep 2007 - Aug 2008	356.8	12.0		11.4	60.1	77.8 ^c
Oct 2007 - Sep 2008	562.0	10.2		10.2	53.3	70.0°
Nov 2007 - Oct 2008	775.9	8.0		9.0	47.0	55.0 ^c
Dec 2007 - Nov 2008	935.4	7.6		8.2	43.0	45.0°
Jan 2008 - Dec 2008	1003.1	7.6		7.9	41.4	42.9 ^c
Feb 2008 - Jan 2009	1007.1	7.6		7.9	41.3	40.9
Mar 2008 - Feb 2009	1021.5	7.6		7.8	41.0	37.5
Apr 2008 - Mar 2009	1030.3	7.6		7.8	40.8	36.0
May 2008 - Apr 2009	1024.2	7.5		7.8	40.9	33.3
Jun 2008 - May 2009	992.8	7.4		8.0	41.6	31.8
Jul 2008 - Jun 2009	1035.1	8.4		7.8	40.7	36.4
Aug 2008 - Jul 2009	1047.8	8.3		7.7	40.4	36.4
Sep 2008 - Aug 2009	1022.2	8.1		7.8	41.0	27.3
Oct 2008 - Sep 2009	948.0	8.16		8.2	42.7	26.1

Notes:

- kac-ft = thousand acre feet.
- ppb = parts per billion. Values are actually in μ g/L (micrograms per liter), which, for the purposes of this report, is equivalent to ppb.
- Compliance is evaluated annually based on the 12-month flow-weighted mean TP concentration for the federal water year ending on September 30. The compliance periods are shown as highlighted rows with bold, italicized text.

^a Flow of 1.82 kac-ft in February 2006 at S355A and S355B was included for the 12-month total flows.

^b Flow of 3.33 kac-ft in August 2006 at S356 structure was included for the 12-month total flows.

^c Value exceeded the guideline percentage.




Figure 6. The 12-month flow-weighted mean TP concentrations in inflows to Everglades National Park through Shark River Slough at the end of each month and the flow-weighted mean concentration for each sampling event. There are no sampling event values for some months because there was little or no flow in those periods.




Figure 7. Daily flows at Shark River Slough structures as a stacked sum of five inflows.²




Figure 8. Daily flows at structures S333 and S334.

13

² Figure 7 was modified on December 16, 2009.




Figure 9. Daily flows at individual inflow structures to Shark River Slough. This figure includes most of the data illustrated in **Figures 7** and **8**.




Figure 10. Total daily flow at Shark River Slough structures and the corresponding flow-weighted mean total phosphorus concentrations for individual sampling events.

Taylor Slough and Coastal Basins

Background

Under the Consent Decree, a single TP long-term limit of 11 ppb, to be met by December 31, 2006, was set for the two points of inflow to Taylor Slough (S332 and S175) and the inflow point to the Coastal Basins (S18C)(see **Appendix C**). The 12-month flow-weighted mean concentrations have consistently been lower than the long-term limit of 11 ppb.

Inflow TP concentrations to the ENP through Taylor Slough and Coastal Basins are compared to the 11 ppb limit at the end of each water year using data from both the old (S175, S332, S18C) and new (S174, S332D, S18C) combinations of structures (**Figure 11**). The narrow bars in **Figure 11** represent the 12-month flow-weighted mean TP concentrations from S332, S175, and S18C for water years 1989 through 2002. The wider bars for water years 1999 through 2009 represent the new combination of structures.

TP and flow data from both sets of structures presented in prior editions of this report through December 2001 (April 2002 report) showed that, beginning in October 2000, the 12-month moving total flow for S332D, S174, and S18C was consistently greater than flow at S332, S175, and S18C. There was also a shift in flow-weighted mean TP concentration data whereby S332D, S174, and S18C concentrations became equal to and then consistently lower than the concentrations at S332, S175, and S18C. These changes reflected the switch from S332 to S332D for water delivery to Taylor Slough between July 3 and July 5, 2000. Consequently, as of the July 2002 report, only S332D, S174, and S18C data are presented for monthly tracking of data in **Figure 11**. However, almost no flow passed through S174 from March 2006 to September 2007. The site was plugged in September 2007, preventing any additional flow.

Reporting Period Update

Figure 12 presents the 12-month and individual sampling event flow-weighted mean TP concentrations at the S332D and S18C structures. All TP grab sample concentrations taken on positive flow days reported for surface water monitoring at the sites were used for the compliance calculations.

The daily flows into ENP through S332D, S174, and S18C are presented in **Figures 13** and **14**.

For the combined flow through S332D and S18C, the 12-month flow-weighted mean TP concentrations for the 12-month periods ending in July, August, and September 2009 were 6.5, 6.3, and 6.2 ppb, respectively (**Table 4**).

The flow-weighted mean TP concentration was below the long-term limit for the 12-month period ending on September 30, 2009. Therefore, Taylor Slough and Coastal Basins TP concentration was in compliance for the federal water year 2009.

The Consent Decree stipulates that the percent of flow-weighted mean TP concentrations greater than 10 ppb from each sampling event in any 12-month period must not exceed a fixed guideline of 53.1 percent. The sampling event flow-weighted mean concentrations generally remained very low. There was no sampling event flow-weighted mean TP concentration greater than 11 ppb since December 2006 except 23 ppb at S18C on July 14, 2008. **Figure 15** shows the relationship between the daily inflows and the corresponding flow-weighted mean TP concentrations for each sampling event.

The USACE authorized the C-111 project in 1995 to restore more natural hydrologic conditions in Taylor Slough and to maintain flood protection to the east of the L31N and C-111 canals.

The original project facilities consisted of pump stations (S332B, S332C, and S332D), detention cells (Cell 1 through Cell 5), a connector cell between Cell 2 and Cell 3, a flow-way cell originating at Berm 3 of Cell 5, and four diversion structures (DS1 through DS4). Upon completion of a USACE construction project in 2009, an interconnected detention system now exists, starting at S-332B west discharge and continuing to the S-332D high head cell.




Figure 11. The 12-month flow-weighted mean total phosphorus concentrations in inflows to Everglades National Park through Taylor Slough and Coastal Basins at the end of each water year compared to the 11 ppb long-term total phosphorus limit.




Figure 12. The 12-month flow-weighted mean total phosphorus concentrations in inflows to Everglades National Park through Taylor Slough and Coastal Basins at the end of each month and the flow-weighted mean total phosphorus concentration for each sampling event.




Figure 13. Daily flows into Everglades National Park through Taylor Slough (structure S332D + S174) and Coastal Basins (structure S18C). Structure S174 was plugged in September 2007 and is no longer used.




Figure 14. Daily flows at individual Coastal Basins (S18C) and Taylor Slough (S332D + S174) structures into the Everglades National Park.

Table 4. Taylor Slough and Coastal Basins total phosphorus compliance tracking.

12-Month	Total Flow	Flow-Weighted Mean TP	Long-Term Limit (ppb)		ampling Events han 10 ppb
Period	(kac-ft)	Concentration (ppb)	Effective 12/31/2006	Guideline	Observed
Nov 2005 - Oct 2006	179.5	5.6	11.0	53.1	3.9
Dec 2005 - Nov 2006	158.8	5.3	11.0	53.1	3.9
Jan 2006 - Dec 2006	137.4	5.2	11.0	53.1	6.0
Feb 2006 - Jan 2007	127.6	5.2	11.0	53.1	6.4
Mar 2006 - Feb 2007	125.4	5.2	11.0	53.1	7.0
Apr 2006 - Mar 2007	123.8	5.2	11.0	53.1	10.0
May 2006 - Apr 2007	125.4	5.2	11.0	53.1	10.0
Jun 2006 - May 2007	126.0	5.2	11.0	53.1	7.5
Jul 2006 - Jun 2007	153.0	5.1	11.0	53.1	6.5
Aug 2006 - Jul 2007	153.4	5.1	11.0	53.1	6.4
Sep 2006 - Aug 2007	143.5	5.0	11.0	53.1	6.5
Oct 2006 - Sep 2007	120.7	4.8	11.0	53.1	4.7
Nov 2006 - Oct 2007	150.3	4.8	11.0	53.1	4.3
Dec 2006 - Nov 2007	155.7	4.8	11.0	53.1	4.4
Jan 2007 - Dec 2007	154.6	4.7	11.0	53.1	2.3
Feb 2007 - Jan 2008	155.5	4.7	11.0	53.1	2.3
Mar 2007 - Feb 2008	156.1	4.7	11.0	53.1	2.3
Apr 2007 - Mar 2008	155.4	4.7	11.0	53.1	0.0
May 2007 - Apr 2008	157.1	4.6	11.0	53.1	0.0
Jun 2007 - May 2008	155.9	4.6	11.0	53.1	0.0
Jul 2007 - Jun 2008	145.0	4.8	11.0	53.1	0.0
Aug 2007 - Jul 2008	130.0	5.0	11.0	53.1	2.6
Sep 2007 - Aug 2008	165.6	5.5	11.0	53.1	2.5
Oct 2007 - Sep 2008	207.7	5.6	11.0	53.1	2.2
Nov 2007 - Oct 2008	234.8	5.7	11.0	53.1	2.3
Dec 2007 - Nov 2008	273.0	5.7	11.0	53.1	2.2
Jan 2008 - Dec 2008	308.8	5.6	11.0	53.1	2.0
Feb 2008 - Jan 2009	317.1	5.6	11.0	53.1	1.9
Mar 2008 - Feb 2009	316.5	5.6	11.0	53.1	1.9
Apr 2008 - Mar 2009	320.0	5.6	11.0	53.1	1.8
May 2008 - Apr 2009	317.6	5.7	11.0	53.1	1.9
Jun 2008 - May 2009	329.2	6.6	11.0	53.1	3.7
Jul 2008 - Jun 2009	358.3	6.5	11.0	53.1	3.5
Aug 2008 - Jul 2009	388.8	6.5	11.0	53.1	1.9
Sep 2008 - Aug 2009	398.6	6.3	11.0	53.1	1.9
Oct 2008 - Sep 2009	411.4	6.2	11.0	53.1	2.2

Notes:

- kac-ft = thousand acre feet.
- ppb = parts per billion. Values are actually in µg/L (micrograms per liter), which, for the purposes of this report, is equivalent to npb
- Compliance is evaluated annually based on the 12-month flow-weighted mean TP concentration for the federal water year ending in September. The compliance periods are shown as highlighted rows with bold, italicized text.




Figure 15. Total daily flow from Taylor Slough and Coastal Basins structures (S332D + S174 and S18C) and the corresponding flow-weighted mean total phosphorus concentrations for individual sampling events.

APPENDIX A

MONTHLY TOTAL PHOSPHORUS CONCENTRATION DATA FOR THE ARTHUR R. MARSHALL LOXAHATCHEE NATIONAL WILDLIFE REFUGE

Settlement Agreement Report July – September 2009

Table A-1. Refuge monthly TP data (ppb).

Month-Year	LOX3	LOX4	LOX5	LOX6	LOX7	LOX8	LOX9	LOX10	LOX11	LOX12	LOX13	LOX14	LOX15	LOX16
Oct-2006	9	8	8	4	7	8	7	8	7	7	8	11	8	7
Nov-2006	16	9	10	6	7	8	8	7	6	6	8	5	8	5
Dec-2006				6	7	7	7	7	4	4	6	5	5	5
Jan-2007	9	7	8	6	7	7	5	6	6	7	7	7	8	8
Feb-2007		12	9	6	7	7	6	6	4	5	6	4	7	5
Mar-2007		12		6	10	10			6	6	7	6	7	6
Apr-2007									8	8	8			
May-2007		1	-	-	-					J5	-			
Jun-2007						15			13		16	11		
Jul-2007	10	13	11	10	9	9	11		9	13	9	9	11	13
Aug-2007	9	17	8	10	10	8	8	15	10	12	-	9	8	11
Sep-2007	8	17	8	6	8	8	10	9	6.5 ^a	10	7	7	8	11
Oct-2007	8	11	7	6	9	10	8	8	10	9	8	8	8	9
Nov-2007	9	7	7	4	8	9	7	7	7	6	7	6	7	8
Dec-2007	11	9	8	5	9	11	8	8	7	9	7	7	8	6
Jan-2008	8	8	7	4	6	9	7	7	5	6	6	5	6	6
Feb-2008	10	8	8	5	7	11	8	8	7	8	7	7	7	7
Mar-2008	8	6	8	5	7	10	6	7	7	6	6	6	7	6
Apr-2008	7	7	8	4	6	13	8	8	4	4	4	4	6	4
May-2008				12	10	14			8	9	8	6	9	8
Jun-2008					10	10			8	6	10	9	10	9
Jul-2008	10	21	8	7	8	18	7	10	8	9	8	8	6	10
Aug-2008	7	11	7	18	8	12	6	9	7	6	8	7	11	10
Sep-2008	9	6	9	5	7	9	7	10	7	9	8	8	7	8
Oct-2008	8	15	7	6	9	9	6	8	7	7	5	6	5	7
Nov-2008	8	7	7	7	6	10	6	8	8	6	9	7	8	8
Dec-2008	8	9	6	4	7	8	4	7	6	7	6	6	6	6
Jan-2009	9	10	9	6	7	10	4	7	6	7	7	6	6	7
Feb-2009		5		4	5	9	4	6	5	4	5	3	4	5
Mar-2009				22	11	13			5	6	5	5	5	6
Apr-2009					18	18			7	7	7	8	8	11
May-2009										11				
Jun-2009 ^b		25		14	10	20	8	15	14	14	12	12	11	11
Jul-2009	6	11	6	6	8	10	7	5	6	17	5	8	7	8
Aug-2009	9	8	7	8	8	12	10	11	7	11	7	10	8	9
Sep-2009	8	20	7	7	8	9	7	7	8	8	8	6	7	9

June 17 and 19, 2009 values are as follows:

Month-Year	LOX3	LOX4	LOX5	LOX6	LOX7	LOX8	LOX9	LOX10	LOX11	LOX12	LOX13	LOX14	LOX15	LOX16
Jun-2009	11	12	13	10	9	14	9	9	11	6	7	11	7	10

Notes:

--- indicates no sample was collected due to insufficient water depth.

^a Average of two values: 6 and 7 ppb

^b June 1 and 2, 2009, compliance sampling data values.

APPENDIX B

WEEKLY GRAB TOTAL PHOSPHORUS CONCENTRATION DATA FOR SHARK RIVER SLOUGH

Table B-1. Weekly Grab TP Concentration Data for Shark River Slough.

Month-Year	S12A	S12B	S12C	S12D	S333	Remarks
10/01/2008	6	5	5	8	7	Compliance data
10/08/2008	6	5	5	8	8	μ
10/16/2008	4	4	5	8	7	Compliance data
10/22/2008	5	4	5	7	8	2000 p. 1000 0000
10/29/2008	5	4	5	7	8	Compliance data
11/05/2008	8	6	5	6	7	
11/12/2008		6	4	7	7	Compliance data
11/19/2008		6	5	8	8	o ompilarios data
11/25/2008	11	6	4	6	8	Compliance data
12/02/2008	13	7	6	6	6	o ompilarios data
12/09/2008	8	10	5	7	6	Compliance data
	9	7	7	8	8	compliance data
12/17/2008 12/23/2008	8				9	Compliance data
	10		9	12	9	compliance data
12/29/2008	10				9	Compliance data
01/06/2009	10				8	Compliance data
01/14/2009						Compliance data
01/21/2009	6 11				7	Compliance data
01/27/2009						Campullanaa data
02/03/2009	11				8	Compliance data
02/10/2009	8				5	0 1 - 1 - 1 - 1
02/18/2009	12				7	Compliance data
02/24/2009	13				8	
03/03/2009	14				8	Compliance data
03/11/2009	19				9	
03/17/2009	27				11	Compliance data
03/24/2009	30				13	
03/31/2009	28				10	Compliance data
04/08/2009	28				14	
04/14/2009	30				13	Compliance data
04/21/2009	48				25	
04/28/2009	53				22	Compliance data
05/05/2009	65				24	
05/12/2009	60				21	Compliance data
05/19/2009	59				23	
05/27/2009	59				30	Compliance data
06/02/2009	79				22	
06/09/2009	41			16	22	Compliance data
06/16/2009	28			13	26	
06/23/2009	31			15	32	Compliance data
06/30/2009	32			16	22	
07/07/2009	13			15	15	Compliance data
07/14/2009	14			6	33	
07/21/2009	7	9	12	13	26	Compliance data
07/28/2009	7	7	9	12	20	
08/04/2009	8	7	9	8	12	Compliance data
08/11/2009	8	7	9	9	11	
08/18/2009	8	8	9	11	10	Compliance data
08/25/2009	7	3	9	10	10	zzp.ia.ioo data
09/01/2009	7	7	10	9	9	Compliance data
09/01/2009	6	7	7	9	9	Compliance data
09/09/2009	6	6	7	9	10	Compliance data
	6	7	6	8	8	Compilative data
09/22/2009		+	7	7		Compliance data
09/29/2009	7	5	/	/	7	Compliance data

Notes:

--- indicates water sample was not collected because the spillway gates were closed at the time of the site visit.

APPENDIX C

WEEKLY GRAB TOTAL PHOSPHORUS CONCENTRATION DATA FOR TAYLOR SLOUGH AND COASTAL BASINS

Table C-1. Weekly Grab TP Concentration Data for Taylor Slough and coastal Basins (ppb).

Month-Year	S332DX	S18C
10/06/2008		5
10/07/2008	6	
10/14/2008		5
10/15/2008	7	
10/20/2008		3
10/21/2008	7	
10/28/2008	6	
11/03/2008		5
11/04/2008	6	
11/12/2008	6	3
11/17/2008		3
11/18/2008	7	
11/24/2008	8	4
12/01/2008	7	5
12/08/2008	6	3
12/15/2008	6	
12/16/2008		5
12/22/2008	5	5
12/29/2008	5	5
01/05/2009	6	4
01/12/2009		4
01/13/2009	5	
01/20/2009	5	3
01/26/2009	5	5
02/02/2009	9	3
02/09/2009	7	3
02/17/2009	8	5
02/23/2009	6	3
03/02/2009	9	7
03/09/2009	6	8

APPENDIX D

CALCULATION METHODS

Long Term Marsh Concentration Levels for Loxahatchee National Wildlife Refuge

Long Term Marsh Concentration Levels:

$$C = 10.7172 - 0.541156S + 1.372\sqrt{7.5819 - 0.9310S + 0.02902216S^2}$$

Terms:

C = the natural log of the geometric mean total phosphorus concentration across 14 marsh stations.

S = average stage measured at gauges CA1-9, CA1-7, and CA1-8C on sampling date (feet).

This equation is applicable over a stage range of 15.42 to 17.14 feet. If the stage on any sampling date exceeds 17.14 feet, a stage of 17.14 feet should be used in calculating the long term concentration levels. The equation shall not apply to dates when the average stage is less than 15.42 feet.

(1991 Settlement Agreement entered as a Consent Decree in 1992 and modified in 1995, Exhibit B, Appendix B, Attachment II, page B-7)

Discharge Limits and OFW Standards for Shark River Slough

Interim Discharge Limit:

$$C = 11.16 - 0.00465Q + 1.397\sqrt{6.377 - 0.00591Q + 0.00000436Q^2}$$

Long-Term Discharge Limit & OFW Standard:

$$C = 11.38 - 0.00538Q + 1.397\sqrt{2.493 - 0.00231Q + 0.00000170Q^2}$$

Frequency Exceedance:

$$F = 48.411 - 0.02896Q + 1.397\sqrt{330.1 - 0.3071Q + 0.0002254Q^2}$$

Terms:

Water Year = October through September

Q = total inflow to Shark River Slough for water year, S-12s + S-333 + any additional inflow from the WCAs established in the future, thousand acre-ft/yr (Kac-ft/yr).

C = limit on maximum flow-weighted-mean inflow concentration for any Water Year, composite of all inflows to Shark Slough (ppb).

F = exceedance for maximum frequency (percent) of inflow concentrations exceeding 10 ppb, computed from the time series of concentrations composited across all inflow structures on each sampling date with positive flow in a given Water Year.

The range of flow (Q) used in deriving the limits is 117 to 1061 Kac-ft/yr. If the total flow for any water year exceeds 1061 Kac-ft/yr, a flow of 1061 Kac-ft/yr should be used in calculating the discharge limits.

(1991 Settlement Agreement entered as a Consent Decree in 1992 and modified in 1995, Exhibit B, Appendix A, Attachment I, page A-5)

Discharge Limits and OFW Standards for Taylor Slough and Coastal Basins

Long-Term Flow-Weighted Discharge Limit & OFW Standard = 11.0 ppb

Frequency Exceedance:

Frequency of values > 10 ppb must be less than 53.1%.

Terms:

Limits are defined on a Water Year basis, October through September.

Basin flow is the total flow through structures S-332, S-175, S-18C, plus any new release points from this basin established in the future, thousand acre-ft/yr (Kac-ft/yr).

Limits apply to the flow-weighted-mean concentration for any Water Year, composite of all inflows to Taylor Slough (S-332) and Coastal Basin (S-18C).

Frequency exceedance is the exceedance for maximum frequency (percent) of inflow concentrations exceeding 10 ppb, computed from the time series of concentrations composited across all inflow structures on each sampling date with positive flow in a given Water Year.

(1991 Settlement Agreement entered as a Consent Decree in 1992 and modified in 1995, Exhibit B, Appendix A, Attachment II, page A-6)