Settlement Agreement Report

Third Quarter July - September 2011

Prepared for the Technical Oversight Committee

November 21, 2011 (Appendix A, Table A-1 was revised on December 19, 2011)



Prepared by:

Cheol Mo, Violeta Ciuca, and Stuart Van Horn

Compliance Assessment and Reporting Section Water Quality Bureau South Florida Water Management District 3301 Gun Club Road West Palm Beach, FL 33406

PURPOSE

The South Florida Water Management District has prepared this report to provide a quarterly update to the Technical Oversight Committee on the compliance status with total phosphorus levels or limits defined in the 1991 Settlement Agreement entered as a Consent Decree in 1992 and modified in 1995. The areas of interest in this report include the interior marsh stations in Arthur R. Marshall Loxahatchee National Wildlife Refuge and two discharges to Everglades National Park: inflows to Shark River Slough and inflows to Taylor Slough and Coastal Basins.

CONTENTS

APPENDICES

Appendix A	Monthly Total Phosphorus Concentration Data for the Arthur R. Marshall	
	Loxahatchee National Wildlife Refuge	A-1
Appendix B	Weekly Grab Total Phosphorus Concentration Data for Shark	
	River Slough	B - 1
Appendix C	Weekly Grab Total Phosphorus Concentration Data for Taylor Slough and	
	Coastal Basins	C-1
Appendix D	Calculation Methods	D - 1
Appendix E	Document revisions	E-1

TABLES

Table 1. Total phosphorus compliance, third quarter 2011	1
Table 2. Loxahatchee National Wildlife Refuge total phosphorus compliance tracking	
Table 3. Shark River Slough total phosphorus compliance tracking.	11
Table 4. Taylor Slough and Coastal Basins total phosphorus compliance tracking	21

FIGURES

Figure 1.	Areas of interest	.2
Figure 2.	Arthur R. Marshall Loxahatchee National Wildlife Refuge water quality	
-	sampling and stage measurement sites	.4
Figure 3.	(A) Monthly total phosphorus geometric mean concentrations for the Arthur	
	R. Marshall Loxahatchee National Wildlife Refuge	.5

i

	(B) Deviation of monthly geometric mean total phosphorus concentrations
	with calculated long-term levels
Figure 4.	Everglades National Park flow structures9
Figure 5.	The 12-month flow-weighted mean total phosphorus concentrations at
	inflows to the Everglades National Park through Shark River Slough at
	the end of each water year10
Figure 6.	The 12-month flow-weighted mean TP concentrations in inflows to Everglades
	National Park through Shark River Slough at the end of each month and the
	flow-weighted mean concentration for each sampling event
Figure 7.	Daily flows into Shark River Slough as a stacked sum of five structures13
Figure 8.	Daily flows at structures S333 and S33413
Figure 9.	Daily flows at individual inflow structures to Shark River Slough14
Figure 10.	Flow at Shark River Slough structures on the day of sampling and the
	corresponding flow-weighted mean total phosphorus concentrations for
	individual sampling events
Figure 11.	The 12-month flow-weighted mean total phosphorus concentrations in inflows
	to Everglades National Park through Taylor Slough and Coastal Basins
Figure 12.	The 12-month flow-weighted mean total phosphorus concentrations in inflows
	to Everglades National Park through Taylor Slough and Coastal Basins at
	the end of each month and the flow-weighted mean total phosphorus
	concentration for each sampling event18
Figure 13.	Daily flows into Everglades National Park through Taylor Slough as a stacked
	total of structures (S332D + S174) and Coastal Basins (S18C)
Figure 14.	Daily flows at individual Coastal Basins (S18C) and Taylor Slough structures
	(S332S + S174) into Everglades National Park20
Figure 15.	Flow from Taylor Slough and Coastal Basins structures (S332D + S174 and S18C)
	on the day of sampling and the corresponding flow-weighted mean total
	phosphorus concentrations for individual sampling events

ACRONYMS AND ABBREVIATIONS

ENP	Everglades National Park
kac-ft	thousand acre feet
NGVD 29	National Geodetic Vertical Datum of 1929
OFW	Outstanding Florida Waters
ppb	parts per billion
Refuge	Arthur R. Marshall Loxahatchee National Wildlife Refuge
SFWMD	South Florida Water Management District
TOC	Technical Oversight Committee
TP	total phosphorus
µg/L	micrograms per liter
USACE	United States Army Corps of Engineers
WCA	Water Conservation Area

EXECUTIVE SUMMARY

This report fulfills the South Florida Water Management District's reporting requirements under the 1991 Settlement Agreement, entered as a Consent Decree in 1992 and modified in 1995, for the third quarter of 2011 (July - September 2011). Total phosphorus (TP) compliance highlights for this period are summarized below for the Arthur R. Marshall Loxahatchee National Wildlife Refuge (Refuge) and Everglades National Park, including Shark River Slough, Taylor Slough, and Coastal Basins (**Table 1** and **Figure 1**):

- **Refuge:** The geometric mean TP concentrations were below the long-term levels in August and September, 2011. The long-term level was not applicable in July 2011.
- Shark River Slough: The 12-month flow-weighted mean TP concentration was below the 12-month moving long-term limit during the third quarter. Shark River Slough was in compliance for Water Year 2011 ending on September 30, 2011.
- **Taylor Slough and Coastal Basins:** The 12-month flow-weighted mean TP concentration was below the 12-month moving long-term limit during the third quarter. Taylor Slough and Coastal Basins was in compliance for Water Year 2011 ending on September 30, 2011.

Month		Geometric Mean TP Concentration (ppb)		Long-term Level (ppb)		Mean Stage (ft NGVD 29)			Number of Samples
Arthur R. I	Marshall	Loxahatch	ee National W	ildlife Refug	е				
Jul 201	1		20.2	N/	'A ^a		15.11		4
Aug 20	11		11.5	16	.6		15.50		8
Sep 207	11		9.8	14	.0		15.80		11
12-Month Period		I Flow	12-Mo Flow-weigh		Long-ter			of Sampling Events ter than 10 ppb	
Ending	(ka	c-ft)	TP Concentra		(рр	(ppb) G		е	Observed
Everglade	s Nationa	al Park – S	hark River Slo	ugh					
Jul 2011	39	9.7	8.0)	11.	1	58.6		47.6
Aug 2011	Aug 2011 31		7.4		11.	11.6			47.4
Sep 2011 ^b	24	7.0	9.2		12.	0	64.1	57.9	
Everglades	s Nationa	al Park – T	aylor Slough a	nd Coastal B	asins				
Jul 2011	15	6.3	5.3	5.3 11.0 53.1			8.8		
Aug 2011	12	26.4	5.5	5	11.	0	53.1		9.4
Sep 2011 ^b	11	1.4	5.6 (5	.5*)	11.	0	53.1		9.4 (11.4 [*])

Table 1. Total phosphorus compliance, third quarter 2011.

Notes:

 ppb = parts per billion. Values are actually in µg/L (micrograms per liter), which, for the purposes of this report, is equivalent to ppb.

• ft NGVD 29 = elevation in feet related to the National Geodetic Vertical Datum of 1929.

• kac-ft = thousand acre feet.

^b Water Year 2011 (October 1, 2010 – September 30, 2011) compliance evaluation period for inflows to Everglades National Park (Shark River Slough, Taylor Slough, and Coastal Basins).

* Including flagged data.

^a N/A denotes that the level was not applicable because the average stage was less than 15.42 feet.




Figure 1. Areas of interest.

ARTHUR R. MARSHALL LOXAHATCHEE NATIONAL WILDLIFE REFUGE

Background

The 1991 Settlement Agreement ended the Everglades lawsuit and was entered into by the federal government, the State of Florida, and the South Florida Water Management District. The subsequent Consent Decree, as modified in 1995, specified that interim and long-term total phosphorus (TP) concentration levels for the Arthur R. Marshall Loxahatchee National Wildlife Refuge (Refuge) must be met by February 1, 1999, and December 31, 2006, respectively. Both the interim and long-term concentration levels vary monthly because they are calculated as a function of water stage measured at gauging stations 1-7, 1-8C, and 1-9 within the Refuge. The stage range within which the interim and long-term concentration levels are applicable is 15.42 to 17.14 feet relative to the National Geodetic Vertical Datum of 1929 (NGVD 29). The monthly TP concentrations are determined from water samples collected at 14 interior marsh stations, LOX3 through LOX16 (**Figure 2**). As required in the Consent Decree, the concentrations are converted to a geometric mean, which is compared to the interim and long-term concentration levels. Monthly TP data for each station for the past 36 months are provided in **Appendix A**. The calculation methods specified in the Consent Decree are provided in **Appendix D**.

Reporting Period Update

Average sampling day stages in the Refuge were 15.11, 15.50, and 15.80 feet in July, August, and September 2011, respectively (**Figure 3** and **Table 2**). The geometric mean, calculated from TP concentrations measured in water samples collected in July, August, and September 2011, were 20.2, 11.5, and 9.8 parts per billion (ppb), respectively. The long-term level was not applicable in July 2011 because the average stages of the scheduled sampling days were below 15.42 feet; the geometric mean TP concentration was below the long-term level for the months of August and September 2011.

TP samples were not collected at stations LOX3, LOX5, LOX6, LOX8, LOX9, LOX10, LOX11, LOX13, LOX14, and LOX16 for the month of July, at stations LOX3, LOX4, LOX5, LOX9, LOX10, and LOX13 for August, and at LOX3, LOX5, and LOX13 for the month of September, because the water depth was less than 0.1 meters at the sites.




Figure 2. Arthur R. Marshall Loxahatchee National Wildlife Refuge water quality sampling and stage measurement sites.

4




Figure 3. (A) Monthly total phosphorus geometric mean concentrations for the Arthur R. Marshall Loxahatchee National Wildlife Refuge compared to calculated long-term levels. These calculated long-term total phosphorus levels are adjusted for fluctuations in stage. The long-term level was not applicable for May 2009 and May through July 2011 because the average stage was less than 15.42 feet. The geometric mean was greater than the long-term level in November 2008 and June 2009.

(**B**) Deviation of monthly geometric mean total phosphorus concentrations with calculated long-term levels. Values greater than zero indicate that the geometric mean was greater than the long-term level.

Month	Geometric Mean TP Concentration (ppb)	Long-Term Level (ppb) Effective 12/31/2006	Average Stage ^a (ft NGVD 29)	Number of Samples
Oct-2008	7.2	7.2	17.42	14
Nov-2008	7.4	7.2	17.22	14
Dec-2008	6.3	7.8	16.95	14
Jan-2009	7.0	8.8	16.68	14
Feb-2009	4.7	10.4	16.35	12
Mar-2009	7.5	13.5	15.86	9
Apr-2009	9.7	14.9	15.69	8
May-2009	11.0	N/A ^b	15.01	1
<i>Jun-2009</i> ്	13.2	12.1	16.05	12
Jul-2009	7.4	9.8	16.47	14
Aug-2009	8.8	11.5	16.16	14
Sep-2009	8.1	8.9	16.68	14
Oct-2009	7.0	8.2	16.86	14
Nov-2009	6.9	9.4	16.55	14
Dec-2009	5.8	10.4	16.36	12
Jan-2010	5.2	8.9	16.68	14
Feb-2010	6.7	9.1	16.62	14
Mar-2010	5.9	9.0	16.65	14
Apr-2010	7.2	9.6	16.51	14
May-2010	7.8	10.4	16.35	14
Jun-2010	7.3	10.9	16.26	14
Jul-2010	6.9	10.7	16.29	14
Aug-2010	6.1	11.6	16.14	10
Sep-2010	6.5	8.5	16.75	14
Oct-2010	6.2	7.7	16.99	14
Nov-2010	6.6	8.2	16.83	14
Dec-2010	4.0	9.6	16.52	13
Jan-2011	5.1	10.7	16.29	10
Feb-2011	6.0	11.3	16.19	10
Mar-2011	8.1	14.0	15.79	8
Apr-2011	11.4	16.3	15.53	7
May-2011	no data	N/A ^b	14.87	0
Jun-2011	no data	N/A ^b	14.30	0
Jul-2011	20.2	N/A ^b	15.11	4
Aug-2011	11.5	16.6	15.50	8
Sep-2011	9.8	14.0	15.80	11

Table 2. Loxahatchee National Wildlife Refuge total phosphorus compliance tracking.

Notes:

• ppb = parts per billion. Values are actually in μ g/L (micrograms per liter), which, for the purposes of this report, is equivalent to ppb.

• ft NGVD 29 = elevation in feet related to the National Geodetic Vertical Datum of 1929.

• Highlighted rows with bold, italicized text indicate when an excursion over the long-term level occurred.

^a Average stage is calculated using stage elevations at stations 1-7, 1-8C, and 1-9 for a given sampling date.

^b N/A denotes that the level was not applicable because the average stage was less than 15.42 feet.

^c June 1 and 2, 2009 compliance sampling data only.

EVERGLADES NATIONAL PARK

Shark River Slough

Background

The Settlement Agreement/Consent Decree (1995) specified that interim and long-term TP concentration limits for discharges into the Everglades National Park (ENP) (Figure 4) through Shark River Slough be met by October 1, 2003, and December 31, 2006, respectively. It was specified that the TP concentrations be presented as 12-month flow-weighted means. Only the TP concentrations for the water year ending September 30 are evaluated for compliance with the Consent Decree limits (**Appendix D**). The long-term TP concentration limit for inflows to Shark River Slough through structures S12A, S12B, S12C, S12D, and S333 represents the concentrations delivered during the Outstanding Florida Waters baseline period of March 1, 1978, to March 1, 1979, and is adjusted for variations in flow. Inflow concentrations of TP through Shark River Slough are compared to the interim and long-term limits at the end of each water year (October 1 through September 30) from 1991 to 2011 (Figure 5). The flow-weighted mean TP concentration was below the long-term limit of 12.0 ppb for the 12-month period ending on September 30, 2011. Therefore, Shark River Slough TP concentration was in compliance for federal water year 2011.

Reporting Period Update

Table 3 presents the 12-month flow-weighted mean concentrations for each month with the corresponding long-term TP concentration limits calculated using the 12-month period flow. Routine monitoring was changed to weekly for all Shark River Slough sites beginning in August 2007. In accordance with Appendix A of the Consent Decree, only the every-other-week grab concentration data were used for the flow-weighted mean calculations from October 2007 forward¹. For the 12-month periods ending in July, August and September 2011, the 12-month flow-weighted mean TP concentrations were 8.0, 7.4, and 9.2 ppb, respectively. The long-term limits were 11.1, 11.6, and 12.0 ppb, respectively for the periods.

The Consent Decree stipulates that the percentage of flow-weighted mean TP concentrations greater than 10 ppb from each sampling event in any 12-month period must not exceed a guideline value based on flow into Shark River Slough for the same 12-month period. For the 12-month periods ending July, August and September 2011, the sampling event TP concentrations greater than 10 ppb were 58.6, 61.7, and 64.1 percent, respectively.

The observed percentages of the sampling event flow-weighted mean TP concentrations greater than 10 ppb were lower than the guideline for the 12-month period ending in July, August and

¹ S12A and S333 are sampled weekly if flowing, otherwise monthly. S12B, S12C, and S12D are sampled weekly if flowing.

September 2011 (**Table 3**). The 12-month flow-weighted mean TP concentrations and the flow-weighted mean TP concentrations for individual sampling events are presented in **Figure 6**.

The daily flows through the individual Shark River Slough structures from October 2008 through September 2011 are presented in **Figures 7** and **9**. The stage in Water Conservation Area 3A (WCA-3A) had been very low and remained in Zone E of the Regulation Schedule since November 2010 but it rose quickly to Zone D in the third quarter. There was no flow through the S12A, B, and C structures during the quarter. A total of 17,886 acre-feet of water was discharged through S333; only 57 acre-feet of water was discharged through S334 during the quarter (**Figure 8**).

For additional information on the WCA-3A regulation schedule, please refer to the U.S. Army Corps of Engineers (USACE) – Jacksonville District's website: <u>http://www.saj.usace.army.mil/h2o/plots/wca3ahp.pdf</u>.

The relationship between the sum of the daily flows at Shark River Slough structures and corresponding flow-weighted mean TP concentrations for individual sampling events is presented in **Figure 10**. Flow and TP concentrations for waters entering the ENP through Shark River Slough have been following an inverse relationship (**Figure 10**).




Figure 4. Everglades National Park flow structures.




Figure 5. The 12-month flow-weighted mean total phosphorus concentrations at inflows to the Everglades National Park through Shark River Slough at the end of each water year compared to the interim and long-term total phosphorus limits. Water Year 2011 concentration was below the long-term limit. Water Year 2008 concentration exceeded the long-term limit but the TOC at the March 1, 2011 quarterly meeting determined substantial evidence indicates this exceedance was due to error.

12-Month	Total Flow	Flow-Weighted Mean TP	Long-Term Limit (ppb)		mpling Events nan 10 ppb
Period	(kac-ft)	Concentration (ppb)	Effective 12/31/2006	Guideline	Observed
Nov 2007 - Oct 2008	775.9	8.0	9.0	47.0	57.9 ^a
Dec 2007 - Nov 2008	935.4	7.6	8.2	43.0	47.4 ^a
Jan 2008 - Dec 2008	1003.1	7.5	7.9	41.4	45.0 ^a
Feb 2008 - Jan 2009	1007.1	7.5	7.9	41.3	42.9 ^a
Mar 2008 - Feb 2009	1021.5	7.5	7.8	41.0	39.1
Apr 2008 - Mar 2009	1030.3	7.5	7.8	40.8	37.5
May 2008 - Apr 2009	1024.2	7.4	7.8	40.9	34.8
Jun 2008 - May 2009	992.8	7.3	8.0	41.6	33.3
Jul 2008 - Jun 2009	1035.1	8.4	7.7	40.7	38.1
Aug 2008 - Jul 2009	1045.1	8.4	7.7	40.4	38.1
Sep 2008 - Aug 2009	1019.4	8.1	7.8	41.0	28.6
Oct 2008 - Sep 2009	945.3	8.2	<i>8.2</i>	42.7	26.1
Nov 2008 - Oct 2009	847.5	9.1	8.7	45.1	27.3
Dec 2008 - Nov 2009	708.3	9.7	9.4	48.9	31.8
Jan 2009 - Dec 2009	647.6	9.9	9.7	50.7	30.4
Feb 2009 - Jan 2010	656.3	9.9	9.7	50.4	30.4
Mar 2009 - Feb 2010	682.1	9.8	9.5	49.6	34.8
Apr 2009 - Mar 2010	733.9	9.7	9.3	48.2	34.8
May 2009 - Apr 2010	790.9	9.9	9.0	46.6	37.5
Jun 2009 - May 2010	869.0	9.9	8.6	44.6	42.3
Jul 2009 - Jun 2010	861.2	9.0	8.6	44.8	42.3
Aug 2009 - July 2010	859.2	8.8	8.6	44.8	42.3
Sep 2009 - Aug 2010	842.5	8.8	8.7	45.3	48.1 ^a
Oct 2009 - Sep 2010	809.9	8.9	8.9	46.1	50.0 ª
Nov 2009 - Oct 2010	757.3	9.0	9.1	47.5	50.0 ^a
Dec 2009 - Nov 2010	742.5	9.0	9.2	47.9	46.2
Jan 2010 - Dec 2010	739.1	9.0	9.2	48.0	46.2
Feb 2010 - Jan 2011	730.5	9.1	9.3	48.3	48.0
Mar 2010 - Feb 2011	695.2	9.1	9.5	49.3	45.8
Apr 2010 - Mar 2011	645.6	9.2	9.7	50.7	54.2 ^a
May 2010 - Apr 2011	585.0	9.0	10.1	52.5	54.2 ^a
Jun 2010 - May 2011	526.0	8.8	10.4	54.4	54.2
Jul 2010 - Jun 2011	484.1	8.6	10.6	55.8	52.2
Aug 2010 - Jul 2011	399.7	8.0	11.1	58.6	47.6
Sep 2010 - Aug 2011	311.9	7.4	11.6	61.7	47.4
<i>Oct 2010 - Sep 2011</i>	247.0	9.2	12.0	64.1	57.9

Table 3. Shark River Slough total phosphorus compliance tracking.

Notes:

• kac-ft = thousand acre feet.

• ppb = parts per billion. Values are actually in $\mu g/L$ (micrograms per liter), which, for the purposes of this report, is equivalent to ppb.

• Compliance is evaluated annually based on the 12-month flow-weighted mean TP concentration for the federal water year ending on September 30. The compliance periods are shown as highlighted rows with bold, italicized text.

^a Value exceeded the guideline percentage.




Figure 6. The 12-month flow-weighted mean TP concentrations in inflows to Everglades National Park through Shark River Slough at the end of each month and the flow-weighted mean concentration for each sampling event. There are no sampling event values for some months because there was little or no flow in those periods.




Figure 7. Daily flows at Shark River Slough structures as a stacked sum of five inflows.




Figure 8. Daily flows at structures S333 and S334.




Figure 9. Daily flows at individual inflow structures to Shark River Slough. This figure includes most of the data illustrated in **Figures 7** and **8**.




Figure 10. Flow at Shark River Slough structures on the day of sampling and the corresponding flow-weighted mean total phosphorus concentrations for individual sampling events.

Taylor Slough and Coastal Basins

Background

Under the Consent Decree, a single TP long-term limit of 11 ppb, to be met by December 31, 2006, was set for the two points of inflow to Taylor Slough (S332 and S175) and the inflow point to the Coastal Basins (S18C) (see **Appendix C**). The 12-month flow-weighted mean concentrations have consistently been lower than the long-term limit of 11 ppb.

Inflow TP concentrations to the ENP through Taylor Slough and Coastal Basins are compared to the 11 ppb limit at the end of each water year using data from both the old (S175, S332, S18C) and new (S174, S332D, S18C) combinations of structures (**Figure 11**). The narrow bars in **Figure 11** represent the 12-month flow-weighted mean TP concentrations from S332, S175, and S18C for water years 1991 through 2002. The wider bars for water years 1999 through 2011 represent the new combination of structures.

TP and flow data from both sets of structures presented in prior editions of this report through December 2001 (April 2002 report) showed that, beginning in October 2000, the 12-month moving total flow for S332D, S174, and S18C was consistently greater than flow at S332, S175, and S18C. There was also a shift in flow-weighted mean TP concentration data whereby S332D, S174, and S18C concentrations became equal to and then consistently lower than the concentrations at S332, S175, and S18C. These changes reflected the switch from S332 to S332D for water delivery to Taylor Slough between July 3 and July 5, 2000. Consequently, as of the July 2002 report, only S332D, S174, and S18C data are presented for monthly tracking of data in **Figure 11**. However, almost no flow passed through S174 from March 2006 to September 2007. The site was plugged in September 2007, preventing any additional flow. The flow-weighted mean TP concentration was below the long-term limit for the 12-month period ending on September 30, 2011. Therefore, Taylor Slough and Coastal Basins TP concentration was in compliance for the federal water year 2011.

Reporting Period Update

Figure 12 presents the 12-month and individual sampling event flow-weighted mean TP concentrations at the S332D and S18C structures. All TP grab sample concentrations taken on positive flow days reported for surface water monitoring at the sites were used for the compliance calculations.

The daily flows into ENP through S332D, S174, and S18C are presented in Figures 13 and 14.

For the combined flow through S332D and S18C, the 12-month flow-weighted mean TP concentration for the periods ending in July, August and September 2011 was 5.3, 5.5, and 5.6² ppb, respectively (**Table 4**).

² 5.4, 5.5, and 5.5 ppb, respectively, if flagged data were included for the flow-weighted mean calculation.


The Consent Decree stipulates that the percent of flow-weighted mean TP concentrations greater than 10 ppb from each sampling event in any 12-month period must not exceed a fixed guideline of 53.1 percent. For the 12 month periods ending July, August, and September 2011, the sampling event TP concentrations greater than 10 ppb were 8.8, 9.4, and 9.4³ percent, respectively.

Figure 15 shows the relationship between the daily inflows and the corresponding flow-weighted mean TP concentrations for each sampling event. The sampling event flow-weighted mean concentrations generally remained very low regardless of flow condition. There had been no sampling event flow-weighted mean TP concentration greater than 11 ppb since December 2006 except 23 ppb on July 14, 2008, and 34 ppb on May 26, 2009, at S18C. TP concentrations were between 4 and 7 ppb in the reporting quarter.

The USACE authorized the C-111 project in 1995 to restore more natural hydrologic conditions in Taylor Slough and to maintain flood protection to the east of the L31N and C-111 canals.

The original project facilities consisted of pump stations (S332B, S332C, and S332D), detention cells (Cell 1 through Cell 5), a connector cell between Cell 2 and Cell 3, a flow-way cell originating at Berm 3 of Cell 5, and four diversion structures (DS1 through DS4). Upon completion of a USACE construction project in 2009, an interconnected detention system now exists, starting at S332B west discharge and continuing to the S332D high head cell.

³ 11.1, 11.4, and 11.4 percent, respectively, if flagged data were included.



Water Year (October 1 - September 30)

Figure 11. The 12-month flow-weighted mean total phosphorus concentrations in inflows to Everglades National Park through Taylor Slough and Coastal Basins at the end of each water year compared to the 11 ppb long-term total phosphorus limit.




Figure 12. The 12-month flow-weighted mean total phosphorus concentrations in inflows to Everglades National Park through Taylor Slough and Coastal Basins at the end of each month and the flow-weighted mean total phosphorus concentration for each sampling event.




Figure 13. Daily flows into Everglades National Park as a stacked sum of Taylor Slough (structures S332D + S174) and Coastal Basins (structure S18C). Structure S174 was plugged in September 2007 and is no longer used.




Figure 14. Daily flows at individual Coastal Basins (S18C) and Taylor Slough (S332D + S174) structures into the Everglades National Park.

12-Month	Total Flow	Flow-Weighted Mean TP	Long-Term Limit (ppb)		Percent of Sampling Events Greater than 10 ppb		
Period	(kac-ft)	Concentration (ppb)	Effective 12/31/2006	Guideline	Observed		
Nov 2007 - Oct 2008	234.8	5.7	11.0	53.1	2.3		
Dec 2007 - Nov 2008	273.0	5.7	11.0	53.1	2.2		
Jan 2008 - Dec 2008	308.8	5.6	11.0	53.1	2.0		
Feb 2008 - Jan 2009	317.1	5.6	11.0	53.1	1.9		
Mar 2008 - Feb 2009	316.5	5.6	11.0	53.1	1.9		
Apr 2008 - Mar 2009	320.0	5.6	11.0	53.1	1.8		
May 2008 - Apr 2009	317.6	5.7	11.0	53.1	1.9		
Jun 2008 - May 2009	329.2	6.6	11.0	53.1	3.7		
Jul 2008 - Jun 2009	358.3	6.5	11.0	53.1	3.5		
Aug 2008 - Jul 2009	388.8	6.5	11.0	53.1	1.9		
Sep 2008 - Aug 2009	398.6	6.3	11.0	53.1	1.9		
Oct 2008 - Sep 2009	411.4	6.2	11.0	53.1	2.2		
Nov 2008 - Oct 2009	399.3	6.1	11.0	53.1	2.3		
Dec 2008 - Nov 2009	383.5	6.2	11.0	53.1	2.3		
Jan 2009 - Dec 2009	391.5	6.3	11.0	53.1	2.4		
Feb 2009 - Jan 2010	395.0	6.2	11.0	53.1	2.3		
Mar 2009 - Feb 2010	414.5	6.2	11.0	53.1	2.1		
Apr 2009 - Mar 2010	418.5	6.2	11.0	53.1	2.1		
May 2009 - Apr 2010	430.6	6.2	11.0	53.1	2.0		
Jun 2009 - May 2010	441.7	5.6	11.0	53.1	0.0		
Jul 2009 - Jun 2010	428.2	5.5	11.0	53.1	0.0		
Aug 2009 - Jul 2010	413.2	5.5	11.0	53.1	0.0		
Sep 2009 - Aug 2010	404.8	5.4	11.0	53.1	0.0		
Oct 2009 - Sep 2010	377.5	5.4	11.0	53.1	0.0		
Nov 2009 - Oct 2010	349.1	5.5	11.0	53.1	0.0		
Dec 2009 - Nov 2010	328.9	5.5	11.0	53.1	0.0		
Jan 2010 - Dec 2010	283.3	5.4	11.0	53.1	0.0		
Feb 2010 - Jan 2011	273.0	5.4	11.0	53.1	0.0		
Mar 2010 - Feb 2011	253.5	5.4	11.0	53.1	0.0		
Apr 2010 - Mar 2011	246.6	5.3	11.0	53.1	0.0		
May 2010 - Apr 2011	235.2	5.3	11.0	53.1	0.0		
Jun 2010 - May 2011	215.0	5.2	11.0	53.1	7.1		
Jul 2010 - Jun 2011	183.4	5.2	11.0	53.1	8.1		
Aug 2010 - Jul 2011	156.3	5.3	11.0	53.1	8.8		
Sep 2010 - Aug 2011	126.4	5.5	11.0	53.1	9.4		
Oct 2010 - Sep 2011	111.4	5.6 (5.5 [*])	11.0	53.1	9.4 (11.4*)		

Table 4. Taylor Slough and	Coastal Basins total	phosphorus com	pliance tracking.

Notes:

• kac-ft = thousand acre feet.

• ppb = parts per billion. Values are actually in µg/L (micrograms per liter), which, for the purposes of this report, is

equivalent to ppb.Compliance is evaluated annually based on the 12-month flow-weighted mean TP concentration for the federal water year ending on September 30. The compliance periods are shown as highlighted rows with bold, italicized text. Including flagged data.




Figure 15. Flow from Taylor Slough and Coastal Basins structures (S332D + S174 and S18C) on the day of sampling and the corresponding flow-weighted mean total phosphorus concentrations for individual sampling events.

APPENDIX A

MONTHLY TOTAL PHOSPHORUS CONCENTRATION DATA FOR THE ARTHUR R. MARSHALL LOXAHATCHEE NATIONAL WILDLIFE REFUGE

Month-Year	LOX3	LOX4	LOX5	LOX6	LOX7	LOX8	LOX9	LOX10	LOX11	LOX12	LOX13	LOX14	LOX15	LOX16
Oct-2008	8	15	7	6	9	9	6	8	7	7	5	6	5	7
Nov-2008	8	7	7	7	6	10	6	8	8	6	9	7	8	8
Dec-2008	8	9	6	4	7	8	4	7	6	7	6	6	6	6
Jan-2009	9	10	9	6	7	10	4	7	6	7	7	6	6	7
Feb-2009		5		4	5	9	4	6	5	4	5	3	4	5
Mar-2009				22	11	13			5	6	5	5	5	6
Apr-2009					18	18			7	7	7	8	8	11
May-2009										11				
Jun-2009 ^a		25		14	10	20	8	15	14	14	12	12	11	11
Jul-2009	6	11	6	6	8	10	7	5	6	17	5	8	7	8
Aug-2009	9	8	7	8	8	12	10	11	7	11	7	10	8	9
Sep-2009	8	20	7	7	8	9	7	7	8	8	8	6	7	9
Oct-2009	7	10	7	4	8	9	7	7	7	9	6	7	5	7
Nov-2009	7	16	6	5	6	9	6	5	7	9	6	6	7	7
Dec-2009		8		4	7	9	6	5	4	5	7	5	6	6
Jan-2010	7	5	7	4	5	8	5	5	5	5	5	4	5	4
Feb-2010	10	8	11	5	8	8	6	7	5	6	5	6	6	6
Mar-2010	11	7	10	2	7	8	6	6	4	6	5	5	6	6
Apr-2010	9	8	10	5	9	10	6	6	6	6	6	7	6	9
May-2010	9	7	14	7	8	8	6	5	9	8	7	8	7	10
Jun-2010	8	8	7	9	7	7	5	5	7	7	6	7	10	12
Jul-2010	9	9	9	8	7	6	5	7	5	6	6	6	7	8
Aug-2010				6	7	6	5	6	6	6		5	7	7
Sep-2010	6	10	6	5	6	6	6	6	6	8	6	6	7	8
Oct-2010	5	17	5	6	5	7	5	7	6	7	5	5	6	7
Nov-2010	5	11	6	6	7	9	5	6	7	7	7	7	5	6
Dec-2010	4	7 (J)	5	3	4	5	4	4	3	5	4	4	4	4
Jan-2011		8		5	6	8			4	5	4	4	4	5
Feb-2011		9		5	7	8			5	6	6	5	4	7
Mar-2011					12	15			7	6	7	7	7	7
Apr-2011						48			10	7	7	12	8	11
May-2011														
Jun-2011														
Jul-2011		30			12					20			23	
Aug-2011				12	13	17			8	11		14	8	12
Sep-2011		13		10	6	10	12	10	8	11		9	8	13

Table A-1. Refuge monthly TP data (ppb).

June 17 and 19, 2009 values are as follows:

Month-Year	LOX3	LOX4	LOX5	LOX6	LOX7	LOX8	LOX9	LOX10	LOX11	LOX12	LOX13	LOX14	LOX15	LOX16
Jun-2009	11	12	13	10	9	14	9	9	11	6	7	11	7	10

Notes:

--- Indicates sample was not collected due to insufficient water depth.

^a June 1 and 2, 2009, compliance sampling data values.

J indicates analyte detected in field blank and/or associated sample. Data qualifier was removed for LOX7 and LOX8 collected in March 2011.

APPENDIX B

WEEKLY GRAB TOTAL PHOSPHORUS CONCENTRATION DATA FOR SHARK RIVER SLOUGH

Date	S12A	S12B	S12C	S12D	S333	Remarks	
10/06/2010	5	6	7	8	8	N/A	
10/13/2010	6	5	6	7	8	Compliance data	
10/19/2010	7	6	7	8	9	N/A	
10/26/2010	6	5	7	9	8	Compliance data	
11/02/2010	19	8	6	10	9	N/A	
11/09/2010	9		6	7	7	Compliance data	
11/16/2010	9	9	7	7	8	N/A	
11/23/2010	12	13	9	9	8	Compliance data	
11/30/2010	9	10	9	8	9	N/A	
12/07/2010	7	7	6	6	7	Compliance data	
12/14/2010	6 (J)				6 (J)	N/A	
12/21/2010	7				6	Compliance data	
12/28/2010	9				39	N/A	
01/05/2011	10				6	Compliance data	
01/01/2011	15				7	N/A	
01/19/2011	15				6	Compliance data	
01/25/2011	12				6	N/A	
	12				7	Compliance data	
02/01/2011	9				19	N/A	
02/08/2011	18				9	Compliance data	
02/15/2011	28				13	N/A	
02/23/2011							
03/01/2011	34				15	Compliance data	
03/08/2011	32				17	N/A	
03/15/2011	66				27	Compliance data	
03/22/2011	74				36	N/A	
03/29/2011	50				31	Compliance data	
04/06/2011	45				31	N/A	
04/12/2011	67				27	Compliance data	
04/19/2011	53				26	N/A	
04/26/2011	64				31	Compliance data	
05/03/2011	56				43	N/A	
05/10/2011	44				50	Compliance data	
05/17/2011	43				29	N/A	
05/24/2011	59				30	Compliance data	
06/01/2011	74				39	N/A	
06/07/2011	78				30	Compliance data	
06/14/2011	62				49	N/A	
06/21/2011	78				35	Compliance data	
06/28/2011	83				30	N/A	
07/06/2011	52				25	Compliance data	
07/13/2011	53				20	N/A	
07/19/2011	93				20	Compliance data	
07/26/2011	47				19	N/A	
08/02/2011	28				17	Compliance data	
08/09/2011	29				24	N/A	
08/16/2011	21				18	Compliance data	
08/23/2011	56				25	N/A	
08/30/2011	19			15	17	Compliance data	
09/07/2011	16			15	13	N/A	
09/13/2011	28			15	13	Compliance data	
	20			13	13	N/A	
09/20/2011 09/27/2011	20			14	13	Compliance data	

Table B-1. Weekly Grab TP Concentration Data for Shark River Slough (ppb).

Notes:

--- indicates water sample was not collected because the spillway gates were closed at the time of the site visit.

J indicates analyte detected in field blank and/or associated sample.

"Compliance data" indicates bi-weekly sampling data used for consent decree calculation.

"N/A" indicates bi-weekly sampling data presented for informational purposes only and not used for consent decree calculation.

APPENDIX C

WEEKLY GRAB TOTAL PHOSPHORUS CONCENTRATION DATA FOR TAYLOR SLOUGH AND COASTAL BASINS

Date	S332DX	S18C
10/04/2010		5
10/05/2010	6	
10/12/2010	6	5
10/18/2010	7	5
10/25/2010	8	5
11/01/2010	6	5
11/08/2010	6	33
11/15/2010	5	3
11/22/2010	6	5
11/29/2010	6	4
12/06/2010	8	5
12/13/2010	10	4
12/20/2010	4	5
12/27/2010	7	7
01/03/2011		7
01/04/2011	6	
01/10/2011	6	7
01/18/2011	6	5
01/24/2011	5	7
01/31/2011	4	4
02/07/2011	5	4
02/14/2011	6	4
02/22/2011	9	4
03/07/2011	7	4
03/14/2011	8	6
03/21/2011	11	7
03/28/2011		7

Table C-1. Weekly Grab TP Concentration Data for Taylor Slough and Coastal Basins (ppb).

Date	S332DX	S18C
04/04/2011		6
04/05/2011	12	
04/11/2011	12	6
04/18/2011	12	7
04/25/2011	21	8
05/02/2011	12	9
05/09/2011	19	11
05/16/2011	17	12
05/23/2011	14	12
05/31/2011	15 (J)	11 (J)
06/06/2011	16	12
06/13/2011	16	13
06/20/2011	18	10
06/27/2011	14	9
07/05/2011	9	7
07/11/2011		7 (J)
07/12/2011	8	
07/18/2011	6	6
07/25/2011	5	5
08/01/2011	11	4
08/08/2011	7 (J)	4 (J)
08/15/2011	7	4
08/22/2011	8	5
08/29/2011	6	4
09/06/2011	7	4
09/12/2011	7	5
09/19/2011	7	4
09/26/2011	7	5

Note: -- indicates water sample was not collected.

APPENDIX D

CALCULATION METHODS

Long Term Marsh Concentration Levels for Loxahatchee National Wildlife Refuge

Long Term Marsh Concentration Levels:

C = 10.7172 - 0.54115 $+ 1.372\sqrt{7.5819 - 0.93105 + 0.02902216^2}$

Terms:

C = the natural log of the geometric mean total phosphorus concentration across 14 marsh stations.

S = average stage measured at gauges CA1-9, CA1-7, and CA1-8C on sampling date (feet).

This equation is applicable over a stage range of 15.42 to 17.14 feet. If the stage on any sampling date exceeds 17.14 feet, a stage of 17.14 feet should be used in calculating the long term concentration levels. The equation shall not apply to dates when the average stage is less than 15.42 feet.

(1991 Settlement Agreement entered as a Consent Decree in 1992 and modified in 1995, *Exhibit B, Appendix B, Attachment II, page B-7*)

Discharge Limits and OFW Standards for Shark River Slough

Interim Discharge Limit:

 $C = 11.16 - 0.00465Q + 1.397\sqrt{6.377 - 0.00591Q + 0.00000436Q^2}$

Long-Term Discharge Limit & OFW Standard:

 $C = 11.38 - 0.00538Q + 1.397\sqrt{2.493 - 0.00231Q + 0.00000170Q^2}$

Frequency Exceedance:

 $F = 48.411 - 0.02896Q + 1.397\sqrt{330.1 - 0.3071Q + 0.0002254Q^2}$

Terms:

Water Year = October through September

Q = total inflow to Shark River Slough for water year, S-12s + S-333 + any additional inflow from the WCAs established in the future, thousand acre-ft/yr (Kac-ft/yr).

C = limit on maximum flow-weighted-mean inflow concentration for any Water Year, composite of all inflows to Shark Slough (ppb).

F = exceedance for maximum frequency (percent) of inflow concentrations exceeding 10 ppb, computed from the time series of concentrations composited across all inflow structures on each sampling date with positive flow in a given Water Year.

The range of flow (Q) used in deriving the limits is 117 to 1061 Kac-ft/yr. If the total flow for any water year exceeds 1061 Kac-ft/yr, a flow of 1061 Kac-ft/yr should be used in calculating the discharge limits.

(1991 Settlement Agreement entered as a Consent Decree in 1992 and modified in 1995, *Exhibit B, Appendix A, Attachment I, page A-5*)

Note: Additional inflows, currently, comprises the discharges through S334, S355A and S335B, and S356. The latest TOC approved methodology to incorporate these additional inflows was documented in the Shark River Slough section of the Settlement Agreement July – September 2006 Report (dated November 9, 2011).

Discharge Limits and OFW Standards for Taylor Slough and Coastal Basins

Long-Term Flow-Weighted Discharge Limit & OFW Standard = 11.0 ppb

Frequency Exceedance:

Frequency of values > 10 ppb must be less than 53.1%.

Terms:

Limits are defined on a Water Year basis, October through September.

Basin flow is the total flow through structures S-332, S-175, S-18C, plus any new release points from this basin established in the future, thousand acre-ft/yr (Kac-ft/yr).

Limits apply to the flow-weighted-mean concentration for any Water Year, composite of all inflows to Taylor Slough (S-332) and Coastal Basin (S-18C).

Frequency exceedance is the exceedance for maximum frequency (percent) of inflow concentrations exceeding 10 ppb, computed from the time series of concentrations composited across all inflow structures on each sampling date with positive flow in a given Water Year.

(1991 Settlement Agreement entered as a Consent Decree in 1992 and modified in 1995, Exhibit B, Appendix A, Attachment II, page A-6)

APPENDIX E DOCUMENT REVISIONS

Page/Date	Original	Revision
A-2 12/19/11	Table A-1. Refuge monthly TP data (ppb).	The TP data for July, August, and September 2010 were misaligned. This problem has been fixed in this report. The data and compliance results in the main body are correct and have not been changed.
		The same errors are in Appendix A of reports for the third and fourth quarters of 2010 and first and second quarters of 2011. These previous reports were not changed. The data and compliance results within the main body of those reports are correct.

Table E-1. Revisions to this report since initial publication.