


## Weekly Climate Update October 21<sup>st</sup> 2008

- Tropical Atlantic /Caribbean temperature is still above normal which could support an increase in late season tropical activity in late October or early November.
- Developing negative subsurface temperature anomalies in the equatorial Pacific is a sign that an El Nino event is very unlikely. Currently the official climate outlook is calling for ENSO to remain in neutral conditions. This is different than climatology because wetter than normal conditions are less likely than the normal climatologic condition. The Position Analysis in slide 14 and 15 best illustrates this point.
- Summary of current global ocean-climate conditions can be found at the following link: [http://ioc3.unesco.org/oopc/state\\_of\\_the\\_ocean/all/](http://ioc3.unesco.org/oopc/state_of_the_ocean/all/)
- The new CPC official climate outlook was released last Thursday. The IRI and CFS models results suggest increased chances of normal to drier than normal conditions for this upcoming dry season.

# Table of Contents

	Slide
2007-2008 La Nina (Nino 3.4 index) compared with other ENSO events during the past several year	<u>2</u>
The evolution of sea surface temperature anomalies (SSTA) in the east-central equatorial Pacific ocean.....	<u>3</u>
The equatorial Pacific Ocean NINO Indices .....	<u>4</u>
Trend in the subsurface anomalies.....	<u>5</u>
Weekly Eastern and Central Equatorial Pacific subsurface Heat Anomaly.....	<u>6</u>
The Climate Forecasting System (CFS) sea temperature anomalies predicted.....	<u>7</u>
Verification of CFS Nino 3.4 Forecasts.....	<u>8</u>
Current Global Sea Surface Temperature Anomalies.....	<u>9</u>
NAO and Atlantic SST anomalies.....	<u>10</u>
Path of 2008 tropical storms to date.....	<u>11</u>
Evolution tropical Atlantic sea surface temperature indices.....	<u>12</u>
The <u>Official CPC Climate Outlook</u> .....	<u>13</u>
Lake Okeechobee October 1 <sup>st</sup> PA Percentiles.....	<u>14</u>
Lake Okeechobee October 1 <sup>st</sup> PA with Neutral ENSO.....	<u>15</u>
Lake Okeechobee October 1 <sup>st</sup> Neutral ENSO/AMO warm phase sub samplings.....	<u>16</u>
Backup Slides	

# El Nino-Southern Oscillation Index



2007-2008 La Nina compared with other ENSO events since 1981.

ENSO is currently in neutral conditions.

[http://ioc3.unesco.org/oopc/state\\_of\\_the\\_ocean/sur/pac/nino3.4.php](http://ioc3.unesco.org/oopc/state_of_the_ocean/sur/pac/nino3.4.php)

# Recent Evolution of Equatorial Pacific SST Departures (°C)

## Climate Prediction Center

### El Nino-Southern Oscillation Weekly Update

Nov.  
2007



Feb.  
2008

Time



July  
2008

Oct.  
2008



Since September 2008, positive sea surface temperature anomalies have weakened over the eastern equatorial Pacific, while negative SST anomalies have strengthened in the central Pacific.

Longitude

# Recent Evolution of Niño Region SST Departures (°C)

Climate Prediction Center

El Niño-Southern Oscillation Weekly Update



The latest weekly SST departures are:

Niño 4 -0.1°C


Niño 3.4 -0.2°C

Niño 3 0.0°C

Niño 1+2 -0.2°C




# Sub-Surface Temperature Departures ( $^{\circ}\text{C}$ ) in the Equatorial Pacific Ocean (October 20<sup>th</sup>, 2008) Bureau of Meteorology Research Centre



At this time neutral conditions are indicated by sea surface temperature (SST) anomalies. However, cool subsurface anomalies have appeared in the eastern and central equatorial Pacific.

## Central & Eastern Pacific Upper-Ocean (0-300 m) Weekly Heat Content Anomalies



The upper ocean heat content was below-average across the eastern half of the equatorial Pacific Ocean between October 2007 and April 2008, and above-average from early May 2008 through mid-August 2008 before returning to below-average within the most recent 60- days.


# Equatorial Pacific SST and Temperature Anomaly Forecast- NCEP Climate Forecast System Issued October 21<sup>st</sup> 2008



NWS/NCEP

Last update: Tue Oct 21 2008  
Initial conditions: 10Oct2008-19Oct2008


PDF correction: Forecast *Nino3.4* SST anomalies from CFS






## Verification of Nino 3.4

### CFS Niño3.4 SST Predictions from Different Initial Months



# Latest Weekly Sea Surface Temperature Anomaly

## National Climate Data Center



The Atlantic Main Hurricane Development Region is still above normal. With the location of the warm anomalies and together with being late in the hurricane season the Caribbean would be the most favorable region for development of storms that would affect south Florida.

# NAO and SST Anomaly in North Atlantic






Fig. NA2. Monthly standardized NAO index (top) derived from monthly standardized 500-mb height anomalies obtained from the NCEP CDAS in 20°N-90°N (<http://www.cpc.ncep.noaa.gov>). Time-Latitude section of SST anomalies averaged between 80°W and 20°W (bottom). SST are derived from the NCEP OI SST analysis, and anomalies are departures from the 1971-2000 base period means.

# Path and Strength of 2008 Tropical Storms and Hurricanes



Negative NAO anomalies tend to direct tropical storms and hurricanes further west with landfall often occurring before recurvature


# Evolution of Tropical Atlantic SST Indices




**Fig. A1a. Tropical Atlantic Variability region indices, calculated as the area-averaged monthly mean sea surface temperature anomalies (°C) for the TNA [60°W-30°W, 5°N-20°N], TSA [30°W-10°E, 20°S-0], and ATL3 [20°W-0, 2.5°S-2.5°N] regions, and Meridional Gradient Index, defined as differences between TNA and TSA. Data are derived from the NCEP OI SST analysis, and anomalies are departures from the 1971-2000 base period means.**

# CPC Seasonal Rainfall Outlook


November



December-February




March-May





# October 15<sup>th</sup> Position Analysis

## Lake Okeechobee SFWMM October 2008 Position Analysis




(See assumptions on the Position Analysis Results website)

Mon Oct 20 14:25:44 2008

# October 15<sup>th</sup> ENSO Neutral years Position Analysis

## Lake Okeechobee SFWMM October 2008 Position Analysis



(See assumptions on the Position Analysis Results website)

Mon Oct 20 14:59:59 2008


ENSO is officially in a neutral conditions according to CPC



# October 15<sup>th</sup> Position Analysis

## ENSO Neutral/AMO Warm sub sampling

### Lake Okeechobee SFWMM October 2008 Position Analysis




Mon Oct 20 14:26:14 2008


# Madden Julian Oscillation (MJO)

MJO is in the easterly phase over Florida this week

Historical (1947-1977)



Current state of MJO




The contrast between the westerly phase (green) and the easterly phase (brown) of the eastward propagating MJO is striking (see maps to left). During westerly MJO phase between 1949-1977, fifty tropical storms developed compared with 14 during easterly phase. The numbers were similarly disparate for hurricane formation, 24 to 6.

# Backup Slides with additional support material

Currently  
transitioning  
to cold phase of  
PDO


## PDO



## ENSO


La Nina predominates when  
PDO is in negative phase →

El Nino predominates when  
PDO is in positive phase →






# Correlation between RF and Atlantic Meridional Mode and Atlantic Multidecadal Oscillation

## AMM




## AMO



# Sea Surface Temperature Anomaly

Tropical Atlantic Main Development Region





## Madden-Julian Oscillation

