



Global sea surface anomaly
1 Jun 2019

EXTENDED HYDROLOGIC OUTLOOK

JUNE 11, 2019



Sea surface temperature anomaly / Anomalie de la température de la mer (C)



Snow depth / Épaisseur de la neige (cm)



Uncovered sea ice
Glace marine à découvert

Climatology 1995-2009 Climatologie



CMC Environnement Canada

CMC Environment Canada

Summary


- The Climate Prediction Center (CPC) is forecasting equal chances of above normal, normal and below normal rainfall for June through August.
- A weak El Niño is likely to continue through summer 2019 (70% chance) and possibly fall (55-60% chance). El Niño increases the chances of a wetter-than-normal dry season and decreases the potential for tropical storm activity from the Main Development Region in the Atlantic Ocean.
- Monitoring Atlantic Multidecadal Oscillation (AMO) index for switch to negative (cold) phase, this has the potential to contribute to drier-than-normal wet seasons.

U. S. Seasonal Outlooks


June-August 2019


The seasonal outlooks combine the effects of long-term trends, soil moisture, and, when appropriate, ENSO.

Precipitation



Temperature





Teleconnections to South Florida

Climate anomalies being related to each other at large distances:

El Niño Southern Oscillation (ENSO)

South Florida dry season (November through May) rainfall is positively correlated with El Niño which has a frequency that ranges between 3 to 7 years while rainfall is negatively correlated with La Niña November through March with a potential increase in tropical rainfall during La Niña

Atlantic Multidecadal Oscillation (AMO)

Average annual inflow to Lake Okeechobee is nearly 50% greater during the warm phase compared to the cold phase of the AMO, easterly flow toward south Florida affected by phase


Pacific Decadal Oscillation (PDO)

Increases variations of south Florida dry season rainfall


Current Global Sea Surface Temperature Anomalies

Global sea surface anomaly and snow cover
11 Jun 2019


Anomalie de la température de la mer et épaisseur de la neige
11 Juin 2019



Sea surface temperature anomaly / Anomalie de la température de la mer (C)



Snow depth / Épaisseur de la neige (cm)



Uncovered sea ice

Glace marine à découvert

Climatologie 1995-2009 Climatology





CMC Environnement Canada
CMC Environment Canada

Niño Region SST Departures ($^{\circ}$ C) Recent Evolution

The latest weekly SST
departures are:

Niño 4	1.1 $^{\circ}$ C
Niño 3.4	0.9 $^{\circ}$ C
Niño 3	0.6 $^{\circ}$ C
Niño 1+2	-0.4 $^{\circ}$ C




Weekly Heat Content Evolution in the Equatorial Pacific

In early August, October, November 2018 and in January-March 2019, positive subsurface temperature anomalies increased, partly due to downwelling Kelvin waves.


During May 2019, an upwelling Kelvin wave contributed to the reduction of positive subsurface anomalies and emergence of negative anomalies around 120°-90°W.

Since mid-May, a downwelling Kelvin wave resulted in the increase of positive subsurface anomalies in the central and east-central Pacific.


Equatorial oceanic Kelvin waves have alternating warm and cold phases. The warm phase is indicated by dashed lines. Downwelling and warming occur in the leading portion of a Kelvin wave, and up-welling and cooling occur in the trailing portion.



Warm Water Volume (5°N – 5°S , 120°E – 80°W)
and NINO 3.4 SST Anomaly




MEI.v2 Evolution of Current ENSO Event in Historical Context




MEI.v2 Evolution of Historical ENSO Events

(a) El Niño



(b) La Niña






NWS/NCEP/CPC

Last update: Tue Jun 11 2019
Initial conditions: 11May2019–20May2019

CFSv2 forecast Nino3.4 SST anomalies (K)



Latest 8 forecast members
Earliest 8 forecast members
Other forecast members

Forecast ensemble mean
NCDC daily analysis

(Model bias correct base period: 1999–2010; Climatology base period: 1982–2010)

IRI/CPC Pacific Niño

3.4 SST Model Outlook

The majority of models predict a weak El Niño to continue into the Northern Hemisphere winter 2019-20.




Figure provided by the International Research Institute (IRI) for Climate and Society (updated 20 May 2019).

Historical El Niño and La Niña Episodes Based on the ONI computed using ERSST.v5

Recent Pacific warm (red) and cold (blue) periods based on a threshold of +/- 0.5 °C for the Oceanic Niño Index (ONI) [3 month running mean of ERSST.v5 SST anomalies in the Nino 3.4 region (5N-5S, 120-170W)]. For historical purposes, periods of below and above normal SSTs are colored in blue and red when the threshold is met for a minimum of 5 consecutive over-lapping seasons.


The ONI is one measure of the El Niño-Southern Oscillation, and other indices can confirm whether features consistent with a coupled ocean-atmosphere phenomenon accompanied these periods. The complete table going back to DJF 1950 can be found [here](#).

Year	DJF	JFM	FMA	MAM	AMJ	MJJ	JJA	JAS	ASO	SON	OND	NDJ
2007	0.7	0.3	0.0	-0.2	-0.3	-0.4	-0.5	-0.8	-1.1	-1.4	-1.5	-1.6
2008	-1.6	-1.4	-1.2	-0.9	-0.8	-0.5	-0.4	-0.3	-0.3	-0.4	-0.6	-0.7
2009	-0.8	-0.7	-0.5	-0.2	0.1	0.4	0.5	0.5	0.7	1.0	1.3	1.6
2010	1.5	1.3	0.9	0.4	-0.1	-0.6	-1.0	-1.4	-1.6	-1.7	-1.7	-1.6
2011	-1.4	-1.1	-0.8	-0.6	-0.5	-0.4	-0.5	-0.7	-0.9	-1.1	-1.1	-1.0
2012	-0.8	-0.6	-0.5	-0.4	-0.2	0.1	0.3	0.3	0.3	0.2	0.0	-0.2
2013	-0.4	-0.3	-0.2	-0.2	-0.3	-0.3	-0.4	-0.4	-0.3	-0.2	-0.2	-0.3
2014	-0.4	-0.4	-0.2	0.1	0.3	0.2	0.1	0.0	0.2	0.4	0.6	0.7
2015	0.6	0.6	0.6	0.8	1.0	1.2	1.5	1.8	2.1	2.4	2.5	2.6
2016	2.5	2.2	1.7	1.0	0.5	0.0	-0.3	-0.6	-0.7	-0.7	-0.7	-0.6
2017	-0.3	-0.1	0.1	0.3	0.4	0.4	0.2	-0.1	-0.4	-0.7	-0.9	-1.0
2018	-0.9	-0.8	-0.6	-0.4	-0.1	0.1	0.1	0.2	0.4	0.7	0.9	0.8
2019	0.8	0.8	0.8	0.8								


CPC/IRI Probabilistic ENSO Outlook

Updated: 9 May 2019


El Niño conditions are favored to continue through winter 2019-20 with diminishing chances.




El Niño & La Niña Events (1950-2018), and Lake Okeechobee Watershed Rainfall & Net Inflow



Source: Cal Neidrauer (SFWMD)



Standardized Klotzbach/Gray Atlantic Multidecadal Oscillation Index (CSU)




2019 Tropical Outlook





2019 Atlantic Hurricane Season Outlook



Named storms
9-15

Hurricanes
4-8

Major hurricanes
2-4

Be prepared: Visit [hurricanes.gov](https://www.hurricanes.gov) and follow @NWS and @NHC_Atlantic on Twitter.

May 23, 2019

ATLANTIC BASIN SEASONAL HURRICANE FORECAST FOR 2019

Forecast Parameter and 1981-2010 Average (in parentheses)	Statistical Forecast	Final Forecast
Named Storms (NS) (12.1)	11.8	14
Named Storm Days (NSD) (59.4)	60.8	55
Hurricanes (H) (6.4)	6.9	6
Hurricane Days (HD) (24.2)	28.4	20
Major Hurricanes (MH) (2.7)	3.2	2
Major Hurricane Days (MHD) (6.2)	7.7	5
Accumulated Cyclone Energy (ACE) (106)	118	100
Net Tropical Cyclone Activity (NTC) (116%)	128	105

We have increased our forecast slightly and now believe that 2019 will have approximately average activity. There remains considerable uncertainty as to whether El Niño conditions will persist through the Atlantic hurricane season. The tropical Atlantic has warmed slightly faster than normal over the past few weeks and now has near-average sea surface temperatures. We anticipate a near-average probability for major hurricanes making landfall along the United States coastline and in the Caribbean. As is the case with all hurricane seasons, coastal residents are reminded that it only takes one hurricane making landfall to make it an active season for them. They should prepare the same for every season, regardless of how much activity is predicted.

Next Update: July 9, 2019


<https://tropical.colostate.edu/>

Source: Colorado State University/Tropical Meteorology Project

Dynamic Position Analysis

- Based on historical climatic conditions spanning the period 1965-2005
- Each year the model resets the initial stages for Lake Okeechobee (LOK) and the Water Conservation Areas (WCAs) to value on the 1st of the previous month and conditions the simulation using real time data during the previous month to achieve real time stage on the 1st of the current month for both Lake Okeechobee and the Water Conservation Areas
- Dynamic Position Analysis
 - Each 1-year simulation starts with current hydrologic conditions (e.g., 1-May-2019)
 - 41 1-year simulations of system response to historical rainfall conditions
 - Statistical summaries used to display projections


Lake Okeechobee SFWMM Jun 2019 Position Analysis




(See assumptions on the Position Analysis Results website)

Lake Okeechobee SFWMM Jun 2019 Position Analysis

All Simulated Years Plot PA_DPA2




Lake Okeechobee SFWMM Jun 2019 Position Analysis



(See assumptions on the Position Analysis Results website)

Lake Okeechobee SFWMM Jun 2019 Position Analysis



(See assumptions on the Position Analysis Results website)