



# EXTENDED HYDROLOGIC OUTLOOK JUNE 12, 2018



Sea surface temperature anomaly / Anomalie de la température de la mer (C)



Snow depth / Épaisseur de la neige (cm)



Uncovered sea ice  
Glace marine à découvert  
Climatologie 1995-2009 Climatologie



CMC Environnement Canada  
CMC Environment Canada


# Summary


- The Climate Prediction Center (CPC) is forecasting above normal rainfall (33-40% chance) for June through August.
- ENSO neutral conditions are present. ENSO-neutral is favored through September-November 2018, with the possibility of El Niño nearing 50% by winter 2018-19.
- Monitoring Atlantic Multidecadal Oscillation (AMO) index for switch to negative (cold) phase, this has the potential to contribute to drier-than-normal wet seasons.

# U. S. Seasonal Outlooks

June - August 2018

The seasonal outlooks combine the effects of long-term trends, soil moisture, and, when appropriate, ENSO.





# Teleconnections to South Florida

Climate anomalies being related to each other at large distances:

## El Niño Southern Oscillation (ENSO)

South Florida dry season (November through May) rainfall is positively correlated with El Niño which has a frequency that ranges between 3 to 7 years while rainfall is negatively correlated with La Niña November through March with a potential increase in tropical rainfall during La Niña

## Atlantic Multidecadal Oscillation (AMO)

Average annual inflow to Lake Okeechobee is nearly 50% greater during the warm phase compared to the cold phase of the AMO, easterly flow toward south Florida affected by phase


## Pacific Decadal Oscillation (PDO)

Increases variations of south Florida dry season rainfall


# Current Global Sea Surface Temperature Anomalies

Global sea surface anomaly and snow cover  
12 Jun 2018


Anomalie de la température de la mer et épaisseur de la neige  
12 Juin 2018



Sea surface temperature anomaly / Anomalie de la température de la mer (C)



Snow depth / Épaisseur de la neige (cm)



Uncovered sea ice

Glace marine à découvert

Climatologie 1995-2009 Climatology





CMC Environnement Canada  
CMC Environment Canada

# Niño Region SST Departures (°C) Recent Evolution

The latest weekly SST departures are:

Niño 4	0.2°C
Niño 3.4	0.0°C
Niño 3	0.2°C
Niño 1+2	-0.5°C




# Weekly Heat Content Evolution in the Equatorial Pacific

From August 2017- early January 2018, negative subsurface temperature anomalies persisted in the central and eastern Pacific Ocean.

From December 2017- May 2018, successive Kelvin waves contributed to the eastward shift of positive and negative subsurface temperature anomalies.


Over the last month, a downwelling Kelvin wave has contributed to the shift of positive temperature anomalies into the eastern Pacific.

Equatorial oceanic Kelvin waves have alternating warm and cold phases. The warm phase is indicated by dashed lines. Downwelling and warming occur in the leading portion of a Kelvin wave, and up-welling and cooling occur in the trailing portion.





### CFSv2 forecast Nino3.4 SST anomalies (K)



# IRI/CPC Pacific Niño

## 3.4 SST Model Outlook

The majority of models predict ENSO-neutral through summer 2018, with an elevated chance of El Niño by fall/winter 2018.




Figure provided by the International Research Institute (IRI) for Climate and Society (updated 18 May 2018).

# Historical El Niño and La Niña Episodes Based on the ONI computed using ERSST.v5

Recent Pacific warm (red) and cold (blue) periods based on a threshold of  $\pm 0.5$  °C for the Oceanic Nino Index (ONI) [3 month running mean of ERSST.v5 SST anomalies in the Nino 3.4 region (5N-5S, 120-170W)]. For historical purposes, periods of below and above normal SSTs are colored in blue and red when the threshold is met for a minimum of 5 consecutive over-lapping seasons.


The ONI is one measure of the El Niño-Southern Oscillation, and other indices can confirm whether features consistent with a coupled ocean-atmosphere phenomenon accompanied these periods. The complete table going back to DJF 1950 can be found [here](#).

Year	DJF	JFM	FMA	MAM	AMJ	MJJ	JJA	JAS	ASO	SON	OND	NDJ
2006	-0.8	-0.7	-0.5	-0.3	0.0	0.0	0.1	0.3	0.5	0.7	0.9	0.9
2007	0.7	0.3	0.0	-0.2	-0.3	-0.4	-0.5	-0.8	-1.1	-1.4	-1.5	-1.6
2008	-1.6	-1.4	-1.2	-0.9	-0.8	-0.5	-0.4	-0.3	-0.3	-0.4	-0.6	-0.7
2009	-0.8	-0.7	-0.5	-0.2	0.1	0.4	0.5	0.5	0.7	1.0	1.3	1.6
2010	1.5	1.3	0.9	0.4	-0.1	-0.6	-1.0	-1.4	-1.6	-1.7	-1.7	-1.6
2011	-1.4	-1.1	-0.8	-0.6	-0.5	-0.4	-0.5	-0.7	-0.9	-1.1	-1.1	-1.0
2012	-0.8	-0.6	-0.5	-0.4	-0.2	0.1	0.3	0.3	0.3	0.2	0.0	-0.2
2013	-0.4	-0.3	-0.2	-0.2	-0.3	-0.3	-0.4	-0.4	-0.3	-0.2	-0.2	-0.3
2014	-0.4	-0.4	-0.2	0.1	0.3	0.2	0.1	0.0	0.2	0.4	0.6	0.7
2015	0.6	0.6	0.6	0.8	1.0	1.2	1.5	1.8	2.1	2.4	2.5	2.6
2016	2.5	2.2	1.7	1.0	0.5	0.0	-0.3	-0.6	-0.7	-0.7	-0.7	-0.6
2017	-0.3	-0.1	0.1	0.3	0.4	0.4	0.2	-0.1	-0.4	-0.7	-0.9	-1.0
2018	-0.9	-0.8	-0.6	-0.4								

# CPC/IRI Probabilistic ENSO Outlook


Updated: 10 May 2018


ENSO-neutral is favored through September-November 2018, with the possibility of El Niño nearing 50% by Northern Hemisphere winter 2018-19.




Reference: Abtew, W. and P. Trimble. 2010. *El Niño Southern Oscillation Link to South Florida Hydrology and Water Management Applications*. Water Resources Management, 24(5):4255-4271, DOI:10.1007/s11269-010-9656-2.

## 2018 ENSO development likelihood





Standardized Klotzbach/Gray Atlantic Multidecadal Oscillation Index (CSU)




# 2018 Tropical Outlook





# 2018 Atlantic Hurricane Season Outlook



■ Above-normal ■ Near-normal ■ Below-normal season

Season probability

**Named storms**  
10-16

**Hurricanes**  
5-9

**Major hurricanes**  
1-4

NOAA

Be prepared: Visit [hurricanes.gov](https://hurricanes.gov) and follow @NWS and @NHC\_Atlantic on Twitter.

May 24, 2018


# 2018 FORECAST AS OF 31 MAY 2018

Forecast Parameter	Statistical Forecast	Analog Forecast	Final Forecast (Including Alberto)	1981-2010 Median
Named Storms (NS)	9.2	12.0	14	12.0
Named Storm Days (NSD)	41.3	57.3	55	60.1
Hurricanes (H)	4.9	7.3	6	6.5
Hurricane Days (HD)	16.9	20.6	20	21.3
Major Hurricanes (MH)	1.7	2.0	2	2.0
Major Hurricane Days (MHD)	3.3	2.1	4	3.9
Accumulated Cyclone Energy (ACE)	69	86	90	92
Net Tropical Cyclone Activity (NTC)	78	96	100	103

# Dynamic Position Analysis

- Based on historical climatic conditions spanning the period 1965-2005
- Each year the model resets the initial stages for Lake Okeechobee (LOK) and the Water Conservation Areas (WCAs) to value on the 1<sup>st</sup> of the previous month and conditions the simulation using real time data during the previous month to achieve real time stage on current month's 1<sup>st</sup> for both Lake Okeechobee and the Water Conservation Areas
- **Dynamic Position Analysis**
  - Each 1-year simulation starts with current hydrologic conditions (e.g., 1-Jun-2018)
  - 41 1-year simulations of system response to historical rainfall conditions
  - Statistical summaries used to display projections


# Lake Okeechobee SFWMM Jun 2018 Position Analysis



(See assumptions on the Position Analysis Results website)

# Lake Okeechobee SFWMM Jun 2018 Position Analysis


All Simulated Years Plot PA



(See assumptions on the Position Analysis Results website)

# Lake Okeechobee SFWMM Jun 2018 Position Analysis

AMO Warm / ENSO Neutral Analog Years Plot PA




(See assumptions on the Position Analysis Results website)

Analog Years are years  
with similar climatological conditions  
to the current year.


# Lake Okeechobee SFWMM Jun 2018 Position Analysis

All ENSO Neutral Years Plot PA



(See assumptions on the Position Analysis Results website)

# Lake Okeechobee SFWMM Jun 2018 Position Analysis



(See assumptions on the Position Analysis Results website)