

MEMORANDUM

TO: Jason Engle, Chief, Engineering Division (USACE)
FROM: John Mitnik, Chief District Engineer (SFWMD)
DATE: May 14, 2026
SUBJECT: System Operational Position Statement May 12, 2026 to May 18, 2026

This Position Statement is to provide operational input for the one-week period from May 12, 2026 to May 18, 2026 based on system conditions and data observed during the previous Monday to Sunday 7-day period.

Current climate conditions: District May rainfall to date is below normal (83% normal). The rainfall forecast (issued May 13) calls for near to below normal rainfall for the coming 7-day period and above normal for the following period.

Climate and weather forecasts: The most recent CPC precipitation outlook for May 2026 is equal chances of below, normal and above normal rainfall (EC) for the entire District. ENSO-neutral conditions favored through Apr - Jun 2026 (80% chance). In May - July 2026, El Niño is likely to emerge (61% chance) and persist through at least the end of 2026. The 3-month window May 2026 – Jul 2026 indicates increased chances (40-50%) of above normal rainfall for the entire District. The 3-month window Jun 2026 – Aug 2026 shows equal chances of below, normal and above normal rainfall (EC) for the state of Florida. The 3-month windows from Jul 2026 – Sep 2026 to Aug 2026 – Oct 2026 indicate slightly increased chances (33-40%) of below normal for the Kissimmee River and north, and increased chances (40-50%) of below Normal rainfall for the remainder of the District. The transition into the 2026 – 2027 Dry Season goes through the 3-month window of Sep 2024 – Nov 2024 showing equal chances of below, normal and above normal (EC) for south Florida. The precipitation outlook for the 2026 - 2027 Dry Season indicates slightly increased chances (33-40%) to increased chances (40-50%) of above normal rainfall for south Florida.

Hydrologic and tropical outlooks: Current climatological conditions are Normal. Current hydrological conditions are Normal. The lake stage is projected to remain in Zone D3 as per LOSOM modeling assumptions.



Water-supply conditions: The Lake Okeechobee seasonal net inflow outlook is Normal and at Low risk for water supply. The multi-seasonal net inflow outlook is Wet and at Low risk for water supply.

Estuary conditions: For the past 7-day periods, May 4 to May 10, 2026, total inflow to the Caloosahatchee River Estuary averaged approximately 350 cfs with 200 cfs coming from Lake Okeechobee through S-77. Salinities in the upper estuary were within the optimal range (0-10) for tape grass at S-79 and Val I-75, and in damaging range (>15) at Ft. Myers. Salinities were in the upper stressed range (>25) at Cape Coral, Shell Point and Sanibel for adult eastern oysters. Total inflow to the St. Lucie Estuary was about 200 cfs with no flow coming from Lake Okeechobee, C-23 Basin, C-24 Basin, and C-44 Basin, about 50 cfs coming from the Ten Mile Creek Basin, and about 150 cfs coming from Tidal Basin. The average salinity in the middle estuary was within the optimal range (10-25) for adult eastern oysters.

Lake Okeechobee stage and ecological conditions:

On May 10, the daily average Lake Okeechobee stage was 10.07 feet NAVD88 (11.38 feet NGVD29), which placed it within the lower portion of Zone D (Zone D3 of the PA25 simulation) of the Lake Okeechobee System Operating Manual (LOSOM). Lake stage decreased by 0.22 feet over the preceding 7-day period. ENSO-neutral is favored through April-June. The District will continue to monitor conditions in the estuaries, as well as the systemwide conditions. Normal Lake Operations continue pursuant to the considerations in LOSOM as informed by PA25. It is recommended that flow targets for the Caloosahatchee Estuary should rely on basin flows to ensure the delivery of the Minimum Flow and Level, but use Lake Okeechobee flows from S-77 to ensure S-79 flows remain above a targeted steady release of 350 cfs; flow targets for the St. Lucie Estuary and Lake Worth Lagoon should remain at 0 cfs consistent with Normal Operations within Zone D. The District will continue to monitor salinity conditions in the estuaries and water supply conditions within the Lake Okeechobee Service Area. The USACE typically implements the releases to the estuaries over a 7-day period starting on Saturday and ending on Friday.

Forecast Modeling Based on PA25 Simulation



The current and seven prior years' annual stage hydrographs for Lake Okeechobee in comparison to the ecological envelope (light grey).

Navigation and recreation conditions: Multiple navigational locks on Lake Okeechobee's north shore are temporarily closed for safety purposes. S-135/G-36/S-127/S-131 Boat Locks are completely closed to navigation. The S-193 Boat Lock will remain open only on Saturdays and Sundays. Boaters are encouraged to exercise caution on the Kissimmee River (C-38 Canal). Public access along the L-29, L-67A and L-67C Canals near Tamiami Trail in Miami-Dade County is closed for maintenance activities.

STOF water supply conditions: Current Lake Okeechobee stage is sufficiently high that water supply deliveries to the Seminole Tribe of Florida (STOF) Brighton Reservation, if needed, will not be impacted. When Lake Okeechobee stage recedes below 8.75 feet NAVD88 (10 feet NGVD29) and 6.75 feet NAVD88 (8 feet NGVD29), water supply delivery is not achievable via Pump Station G-207 on the Harney Pond Canal and Pump Station G-208 on the Indian Prairie Canal, respectively, as the respective canals become disconnected from Lake Okeechobee.

Algal Bloom conditions: The Fish and Wildlife Research Institute reported on May 8, 2026, that *Karenia brevis*, the Florida red tide dinoflagellate, was not observed at bloom concentrations in any samples collected within the District region. In the most recent non-obscured satellite image from May 9, 2026, NOAA's Harmful Algal Bloom Monitoring System suggests moderate to high cyanobacteria potential in Fisheating Bay and along much of the Indian Prairie shoreline.

STA conditions:


In STA-1E, Central Flow-way is offline for construction activities. An operational restriction is in place in the Western Flow-way for post-construction vegetation grow-in. Online treatment cells are at or near target stage. In STA-1W, Eastern Flow-way is offline for vegetation management activities. Treatment cells are at or near target stage. Vegetation in the Western and Eastern Flow-ways is highly stressed. The 365-day PLRs for the Northern and Western Flow-ways are below 1.0 g/m²/yr. In STA-2, treatment cells are at target stage. The 365-day PLRs for all Flow-ways are below 1.0 g/m²/yr. In STA-3/4, an operational restriction is in place in the Eastern Flow-way and Central Flow-way for vegetation management activities. Most treatment cells are at or slightly above target stage. Vegetation in the Central Flow-way is highly stressed. The 365-day PLR for the Eastern, Central, and Western Flow-ways are below 1.0 g/m²/yr. For the current operational period, USACE is not requesting flows south from Lake Okeechobee towards the WCAs. The District will continue to work with the USACE to manage Lake Okeechobee levels in an effort to curtail harmful discharges over this year. To help with this objective the District will move as much water south through the Stormwater Treatment Areas as possible under the current permits as regional conditions allow.

WCA conditions: On May 10, the daily average stage in WCA-1 was at 14.52 feet NAVD88 (16.11 feet NGVD29), in Zone A and 0.35 feet above regulation schedule. The daily average stage in WCA-2A was at 10.33 feet NAVD88 (11.92 feet NGVD29), in Zone A and 0.91 feet above regulation schedule. The daily average stage in WCA-3A was at 6.41 feet NAVD88 (7.93 feet NGVD29), in Zone C and 1.72 feet below regulation schedule. Over the 7-day period, May 4, 2026 to May 10, 2026, no regulatory releases were sent from Lake Okeechobee south to the STAs. No Lake regulatory releases reached Lake Worth Lagoon through the C-51 canal during this period.


ENP conditions: Releases from WCA-3A to the ENP continue through the S333 structures. In ENP, both Taylor Slough and Shark River Slough have less connection from north to south with Taylor Slough shows more connection to the coast. WCA-1 has water depths near or above average with some ponding beginning in the SE portion. WCA-2A has water depths an average in the central and south but drier to the north. WCA-3A remains very dry, having received the least amount of rainfall in the system over the last month. Stages decreased in Taylor Slough over the past week but remain above the recent average. Salinity decreased on average in Florida Bay compared to last week and remains above the estimated historical average and within the WY2001-2016 Interquartile Range (IQR) for all three regions. The Tamiami Trail Flow Formula (TTFF) recommends 113 cfs of daily target releases from WCA-3A to ENP. The District recommends continuing with the current operations for the releases from WCA-3A in accordance with the Combined Operating Plan.

May 2026 Conditional Position Analysis (CPA) results for Lake Okeechobee, WCA-1, WCA-2A and WCA-3A under LOSOM Recovery Operations.


LOK




WCA1 Site 8C




S11B_H



WCA3AAvg





Eastern (top panel), Central (middle panel) and Western (bottom panel) Florida Bay daily average salinities with WY2001-2016 interquartile (25-75 percentile) ranges (IQR) and estimated historical daily average salinities. The hyper salinity threshold indicates the level at which salinities start to become harmful to seagrass.