MEMORANDUM

TO: John Mitnik, Assistant Executive Director

THROUGH: Peter Kwiatkowski, Section Administrator, Resource Evaluation

FROM: SFWMD Staff Water Supply Advisory Team

DATE: January 3, 2023

SUBJECT: Water Supply Report

District-wide Conditions

All the United States Geological Survey (USGS) real-time wells in the Kissimmee Basin (KB) are in the median and upper percentile ranges for this time of year. The wells in the Upper KB are mostly completed in the Floridan aquifer and the wells in the Lower KB are surficial aquifer system wells. Surface and groundwater levels decreased in about 90 percent of the KB stations over the last week.

Upper East Coast (UEC) surface water levels increased, and groundwater stations decreased during the last week. Stages in UEC canals C-23, C-24, and C-25 are 22.88, 20.90, and 22.47 feet, all above the fourteen feet agricultural cut-off. All the UEC wells are at median and higher levels for this time of year.

In the Lower East Coast, about 70 percent of the surface and groundwater stations recorded decreasing trends over the past seven days. Surface water levels are on the low side in the C-111 basin (S-176 and S-177). About 90 percent of the LEC surficial aquifer system stations are in the median and upper percentile ranges for this time of year.

Groundwater levels decreased in all the Lower West Coast (LWC) stations over the last week. All the monitored wells in the Surficial aquifer system and the Sandstone aquifer are in the median and upper percentile ranges for this time of year. About 15 percent of the Lower Tamiami aquifer stations and about 30 percent of the Mid-Hawthorn aquifer wells are in the lower percentile ranges.

Figure 1 shows a statistical comparison between current groundwater levels and long-term historical monthly average groundwater levels at representative wells throughout the District.




Figure 1. Current Groundwater Level Conditions

Water Supply Technical Input to LORS2008

The Palmer Drought Index for Lake Okeechobee (LOK) Tributary Conditions was 0.74 which is classified as "normal to extremely wet" and is in the "low" risk category for water supply. The projected LOK stage for the next two months is Intermediate Sub-Band, and the risk to water supply is categorized as "low". The Climate Prediction Center's (CPC) Precipitation Outlook is projected as "normal" for one month and "below normal" for three months, leaving the 1-month projection in the "low" risk category and the 3-month projection in the "moderate" risk category. The LOK Seasonal Net Inflow Outlook is "dry" and is in the "moderate" risk for water supply. The LOK Multi-Seasonal Net Inflow Outlook is in the "normal" range with "moderate" risk to water supply. The stages in WCA 1, WCA-2, and WCA-3 are all above line 1 and are in the "low" risk category. The Year-Round Irrigation Rule is in effect for the three LEC Service Areas. All three LEC Service Areas are in the "low" risk category for water supply. **Figure 2** summarizes the water supply risk indicators.

LORS2008 Implementation on 01/02/2023 (ENSO Condition- La Niña Watch): Status for week ending 01/02/2023:

Water Supply Risk Evaluation

Area	Indicator	Value	Color Coded Scoring Scheme
LOK	Projected LOK Stage for the next two months	Intermediate Sub-band	L
	Palmer Drought Index for LOK Tributary Conditions	0.74 (Normal to Extremely Wet)	L
	CPC Precipitation Outlook	1 month: Normal	L
		3 months: Below Normal	М
	LOK Seasonal Net Inflow Outlook	0.27 ft	М
	ENSO Forecast	Dry	
	LOK Multi-Seasonal Net Inflow Outlook	2.65 ft	M
	ENSO Forecast	Normal	IVI
WCAs	WCA 1: 3 Station Average (Sites 1-8C)	Above Line 1 (17.36 ft)	L
	WCA 2A: Site 2-17	Above Line 1 (12.47 ft)	L
	WCA-3A: 3 Station Average (Sites 63, 64, and 65)	Above Line 1 (10.21 ft)	L
LEC	Service Area 1	Year-Round Irrigation Rule in effect	L
	Service Area 2	Year-Round Irrigation Rule in effect	L
	Service Area 3	Year-Round Irrigation Rule in effect	L

Note: The water supply risk classification based on the Palmer index, as well as the LOK seasonal and multi-seasonal net inflow outlooks use slightly different classification intervals than those used by the 2008-LORS.