MEMORANDUM

то:	John Mitnik, Assistant Executive Director		
THROUGH:	Peter Kwiatkowski, Section Administrator, Resource Evaluation		
FROM:	SFWMD Staff Water Supply Advisory Team		
DATE:	June 2 nd , 2020		
SUBJECT:	Water Supply Report		

District-wide Conditions

Surface and groundwater levels showed mixed trends throughout the District over the last week. The majority of the United States Geological Survey (USGS) real-time wells in the Kissimmee Basin (KB) within the District boundaries are at median levels for this time of year. The wells in the Upper KB are completed in the Floridan aquifer and the wells in the Lower KB are completed in the surficial aquifer system. Most of the surface and groundwater stations throughout the KB recorded decreases in water levels over the last week.

Stages in the Upper East Coast (UEC) canals C-23, C-24, and C-25 are 21.59, 19.71, and 19.31 feet, all above the fourteen feet agricultural cut-off. The surficial aquifer stations are at median levels and higher. Surface and groundwater levels increased in about three quarters of the Lower East Coast (LEC) stations over the past week. Approximately ten percent of the Biscayne aquifer wells are at median levels, with the remainder in the upper percentile ranges for this time of year.

Groundwater levels increased in most of the Lower West Coast (LWC) stations over the last seven days. Water Shortage restrictions remain in effect for Lee County at this time. Since last week all of the Surficial aquifer wells have recovered into their median percentile ranges. The majority of the Lower Tamiami aquifer wells are at median levels and higher for this time of year. Sandstone aquifer monitor wells have also climbed into their median and upper percentile ranges. Approximately fifty percent of the Mid-Hawthorn aquifer monitor wells are in the lower percentile ranges, with the remainder in the upper percentile ranges. **Figure 1** summarizes current conditions.




Figure 1. Real-Time Groundwater Level Map

Water Supply Technical Input to LORS2008

The Palmer Index for Lake Okeechobee (LOK) Tributary Conditions is –1.88 classified as "dry," and is in the "moderate" risk category for water supply. The LOK stage for the next two months is projected to be in the Beneficial Use Band, and the risk to water supply is categorized as "moderate." The Climate Prediction Center's (CPC) Precipitation Outlook is projected as "above normal" for one month and "above normal" for three months, leaving both the one-month outlook and three-month outlook in the "low" risk category. The LOK Seasonal Net Inflow Forecast is in the "normal to extremely wet" category and is in the "low" risk category. The Multi-Seasonal Net Inflow Forecast is in the "normal" range with "moderate" risk to water supply. The stages in WCA 1, WCA 2A and WCA 3A are all above line 1 and are in the "low" risk category. Year-Round Irrigation Rule is in effect for the LEC Service Areas. All Service Areas are in the "low" risk category.

LORS2008 Implementation on 06/1/2020 (ENSO Neutral Condition):

Status for week ending 6/1/2020:

Water Supply Risk Evaluation

Area	Indicator	Value	Color Coded Scoring Scheme	
LOK	Projected LOK Stage for the next two months	Beneficial Use sub band	М	
	Palmer Index for LOK Tributary Conditions	-1.88 (Dry)	М	
	CPC Precipitation Outlook	1 month: Above Normal	L	
	OF C Fredpitation Outlook	3 months: Above Normal	L	
	LOK Seasonal Net Inflow Outlook 2.81 ft			
	ENSO Forecast (positive)	Normal to Extremely Wet		
	LOK Multi-Seasonal Net Inflow Outlook	3.06 ft		
	ENSO Forecast (positive)	Normal	М	
WCAs	WCA 1: 3 Station Average (Site 1-7, 1-8T and 1-9)	Above Line 1 (16.11 ft)	L	
	WCA 2A: Site 2-17	Above Line 1 (12.15 ft)	L	
	WCA-3A: 3 Station Average (Site 63, 64 and 65)	Above Line 1 (9.31 ft)	L	
LEC	Service Area 1	Year-Round Irrigation Rule in effect	L	
	Service Area 2	Year-Round Irrigation Rule in effect	L	
	Service Area 3	Year-Round Irrigation Rule in effect	L	

Note: The water supply risk classification based on the Palmer index, as well as the LOK seasonal and multi-seasonal net inflow outlooks use slightly different classification intervals than those used by the 2008-LORS.

Figure 2. Water Supply Risk Indicators