

MEMORANDUM

TO: John Mitnik, Division Director, Operations, Engineering, and Construction

THROUGH: Peter Kwiatkowski, Section Administrator, Resource Evaluation

FROM: SFWMD Staff Water Supply Advisory Team


DATE: June 27, 2017

SUBJECT: Water Supply Report

District-wide Conditions

Surface and groundwater levels showed mixed trends throughout the District over the last week. The majority of United States Geological Survey (USGS) real-time wells in the Kissimmee Basin (KB) within the District boundaries are at median levels. Approximately two thirds of surface and groundwater stations across the KB recorded increases in water levels over the last week. Stages in the Upper East Coast (UEC) canals C-23, C-24, and C-25 are at 21.30, 19.11, and 19.26 feet. Two thirds of UEC surficial aquifer wells are in the 76th to 90th percentile range or higher for this time of year, with the remainder at median levels. Surface and groundwater levels increased in approximately half of the Lower East Coast (LEC) monitoring stations over the past week. Water levels are still a little low in parts of Everglades National Park, and C-111 basin. The majority of the USGS Biscayne aquifer monitor wells are within their median percentile range or higher with the recent rain.

In the Lower West Coast (LWC), groundwater levels decreased in most of the monitor wells over the last seven days. About two thirds of the wells in the Surficial aquifer are in the upper 76th to 90th percentile range, with most of the remainder at median levels. The majority of the Lower Tamiami aquifer wells are in the upper 76th to 90th percentile range or higher. Approximately seventy five percent of the Sandstone aquifer monitor wells are at median levels, with most of the remainder in the upper 76th to 90th percentile range for this time of year. About two thirds of the Mid-Hawthorn aquifer monitor wells are at median levels or higher, with most of the remainder in the lower 10th to 24th percentile range. **Figure 1** summarizes current water level conditions.



Explanation - Percentile classes (symbol color based on most recent measurement)							Wells	Springs
●	●	●	●	●	●	●	●	●
Low	<10	10-24	25-75	76-90	>90	High	Not Ranked	Real-Time
	Much Below Normal	Below Normal	Normal	Above Normal	Much Above Normal			Continuous
								Periodic Measurements

Figure 1. [Florida Real-Time Groundwater Level Network Map](#)

Water Supply Technical Input to LORS2008

The Palmer Index for Lake Okeechobee (LOK) Tributary Conditions is -1.87, classified as “dry,” and is in the “moderate” risk category. The LOK stage for the next two months is projected to be in the Beneficial Use Sub-Band, and the risk to water supply is categorized as “moderate.” The Climate Prediction Center’s (CPC) Precipitation Outlook is projected as “normal” for one month and “normal” for three months, leaving both the one month outlook and the three month outlook in the “low” risk category. The LOK Seasonal Net Inflow Forecast is in the “normal” range, with “low” risk to water supply. The Multi-Seasonal Net Inflow Forecast is projected as “wet” with “low” risk to water supply. The stages in all Water Conservation Areas are above line 1 and are in the “low” risk category. Year-Round Irrigation Rule is in effect for the LEC Service Areas.

Figure 2 summarizes the water supply risk indicators.

LORS2008 Implementation on 6/26/2017 (ENSO Neutral Condition):

Status for week ending 6/26/2017:

District wide, Raindar rainfall was 0.62 inches for the week. Lake stage on 6/26/2017 was 12.28 ft, up 0.24 ft from last week.

The updated June 2017 SFWMM Dynamic Position Analysis [percentile graph](#) for Lake Okeechobee show that the current lake stage is in the Beneficial Use Operational Sub-Band.

The LORS2008 tributary [indices](#) are classified as **Wet**. The PDSI indicates dry condition and the LONIN is Wet. The classification is based on the wetter of the two.

Water Supply Risk Evaluation

Area	Indicator	Value	Color Coded Scoring Scheme
LOK	Projected LOK Stage for the next two months	Beneficial Use Sub Band	M
	Palmer Index for LOK Tributary Conditions	-1.87 (Dry)	M
	CPC Precipitation Outlook	1 month: Normal	L
		3 months: Normal	L
	LOK Seasonal Net Inflow Outlook	3.68 ft (Normal)	L
	ENSO La Nina Years	4.23 ft (Wet)	L
	LOK Multi-Seasonal Net Inflow Outlook		
WCAs	WCA 1: Site 1-7, Site 1-8T, & Site 1-9 Average	Above Line 1 (16.55 ft)	L
	WCA 2A: Site 2-17 HW	Above Line 1 (14.20 ft)	L
	WCA-3A: 3 Station Average (Site 63, 64 and 65)	Above Line 1 (11.17 ft)	L
LEC	Service Area 1	Year-Round Irrigation Rule in effect	L
	Service Area 2	Year-Round Irrigation Rule in effect	L
	Service Area 3	Year-Round Irrigation Rule in effect	L

Note: The water supply risk classification based on the Palmer index, as well as the LOK seasonal and multi-seasonal net inflow outlooks use slightly different classification intervals than those used by the 2008-LORS.

Figure 2. Water Supply Risk Indicators