

MEMORANDUM

TO: John Mitnik, Assistant Executive Director

THROUGH: Peter Kwiatkowski, Section Administrator, Resource Evaluation

FROM: SFWMD Staff Water Supply Advisory Team

DATE: July 6th, 2021

SUBJECT: Water Supply Report

District-wide Conditions

Surface and groundwater levels generally increased throughout the District over the last week. Most of the United States Geological Survey (USGS) real-time wells in the Kissimmee Basin (KB) within the District boundaries are at median levels for this time of year. The wells in the northern portion of the KB are mostly completed in the Floridan aquifer and the wells in southern KB in the surficial aquifer system. The majority of surface and groundwater stations throughout the KB recorded increases in water levels over the last week.

Stages in the Upper East Coast (UEC) canals C-23, C-24, and C-25 are 20.86, 18.82, and 19.64 feet, all above the fourteen feet agricultural cut-off. About half of the surficial aquifer stations are at median levels for this time of year, with the remainder in the lower percentile ranges. Surface and groundwater levels increased in about half of the Lower East Coast (LEC) stations during the past week. Approximately eighty percent of the Biscayne aquifer wells are at median levels and higher for this time of year. Groundwater levels are on the low side in Everglades National Park, Water Conservation Area 3A (WCA 3A) and C-111 Basin.

Groundwater levels increased in most of the stations on the Lower West Coast (LWC) over the last seven days. Approximately three quarters of the surficial aquifer wells are at median levels for this time of year. About two thirds of the Lower Tamiami aquifer wells are in the lower percentile ranges, with the remainder at median levels and higher. Approximately eighty percent of Sandstone aquifer wells are in the lower percentile ranges for this time of year, with the remainder at median levels. About two thirds of the Mid-Hawthorn aquifer monitor wells are in the lower percentile ranges, with the remainder at median levels for this time of year. **Figure 1** summarizes current conditions.




Figure 1. Real-Time Groundwater Level Map

Water Supply Technical Input to LORS2008

The Palmer Index for Lake Okeechobee (LOK) Tributary Conditions was -2.21 and is classified as “extremely dry,” in the “high” risk category for water supply. The LOK stage for the next two months is projected to be in the Base Flow Sub-band, and the risk to water supply is categorized as “moderate.” The Climate Prediction Center’s (CPC) Precipitation Outlook is projected as “normal” for one month and “normal” for three months, leaving both the one-month and three-month outlook in the “low” risk category. The LOK Seasonal Net Inflow Forecast is in the “normal to extremely wet” category and is in the “low” risk category. The Multi-Seasonal Net Inflow Forecast is in the “normal” range with “moderate” risk to water supply. The stage in WCA 1 and WCA 2A are above line 1 and are in the “low” risk category. WCA 3A stages is between line 1 and line 2 and is in the “moderate” risk category. Year-Round Irrigation Rule is in effect for the LEC Service Areas. All Service Areas are in the “low” risk category for water supply. **Figure 2** summarizes the water supply risk indicators.

LORS2008 Implementation on 7/5/2021 (ENSO Condition- ENSO-neutral):

Status for week ending 7/5/2021:

Water Supply Risk Evaluation

Area	Indicator	Value	Color Coded Scoring Scheme
LOK	Projected LOK Stage for the next two months	Base Flow Sub-band	M
	Palmer Drought Index for LOK Tributary Conditions	-2.21 (Extremely Dry)	H
	CPC Precipitation Outlook	1 month: Normal	L
		3 months: Normal	L
	LOK Seasonal Net Inflow Outlook	2.53 ft	L
	ENSO Forecast	Normal to Extremely Wet	L
	LOK Multi-Seasonal Net Inflow Outlook	2.65 ft	M
ENSO Forecast		Normal	M
WCAs	WCA 1: Site 1-8C	Above Line 1 (16.01 ft)	L
	WCA 2A: Site 2-17	Above Line 1 (12.21 ft)	L
	WCA-3A: 3 Station Average (Site 63, 64 and 65)	Line 1- Line 2 (8.73 ft)	M
LEC	Service Area 1	Year-Round Irrigation Rule in effect	L
	Service Area 2	Year-Round Irrigation Rule in effect	L
	Service Area 3	Year-Round Irrigation Rule in effect	L

Note: The water supply risk classification based on the Palmer index, as well as the LOK seasonal and multi-seasonal net inflow outlooks use slightly different classification intervals than those used by the 2008-LORS.

Figure 2. Water Supply Risk Indicators