

Disclaimer: Information contained in the report addresses environmental conditions only and is not the official South Florida Water Management District operations recommendation or decision.

M E M O R A N D U M

TO: John Mitnik, Assistant Executive Director, Executive Office Staff

FROM: SFWMD Staff Environmental Advisory Team

DATE: March 23, 2022

SUBJECT: Weekly Environmental Conditions for Systems Operations

Summary

Weather Conditions and Forecast

A cold front will cross the Gulf of Mexico and move into north Florida late Wednesday. Ahead of the front, a good moisture supply will support isolated or widely scattered showers across SFWMD that should be greatest in coverage during the afternoon over the eastern part of the area. A squall line immediately preceding the cold front is forecast to bring heavy rains into the far northwestern part of the SFWMD before daybreak on Thursday. Heavy rains will push farther south into the northern and northwestern parts of the SFWMD early on Thursday. Additional rains could occur during the evening and overnight over parts of the SFWMD before the last of any rains exit the southeast by or just after sunrise on Friday. Following the frontal passage, a cooler and much drier air mass will overspread the SFWMD during the day on Friday. A secondary frontal passage without any rain is then expected late Saturday or Sunday, which will reinforce an unseasonably cool and dry weather pattern through early next week. Some moderation of temperatures is likely by late Tuesday. However, there is a good chance that a pattern of no rainfall would continue through at least mid-week.

Kissimmee

Flow at S-59 and S-61 is being increased to manage stage reversals on lakes East Toho and Toho. Flow at S-65 / S-65A is also being increased slowly to manage the stage reversals on KCH and the Kissimmee River; water depth on the Kissimmee River floodplain increased, with a mean depth of 0.29 feet as of March 20, 2022. The concentration of dissolved oxygen in the Kissimmee River remained well above the concentration of concern, with an average of 8.3 mg/L for the week ending on March 20, 2022.

Lake Okeechobee

Lake Okeechobee stage was 14.13 feet NGVD on March 20, 2022, with water levels 0.66 feet lower than a month ago (**Figure LO-1**). Lake stage fell back to within the ecological envelope on January 1, 2022, after being above the envelope since late September 2021,

and having spent a total of 279 days (79%) in 2021 above the envelope (**Figure LO-2**). Average daily inflows (excluding rainfall) increased from the previous week, going from 345 cfs to 705 cfs. Average daily outflows (excluding evapotranspiration) decreased from the previous week, going from 3,550 cfs to 2,408 cfs. Nineteen snail kite nests were observed on Lake Okeechobee, mostly within the Indian Prairie marsh. Recent satellite imagery (March 20, 2022) showed a few scattered areas of low to moderate bloom potential, in Fisheating Bay and along the northwestern shoreline (**Figure LO-6**).

Estuaries

Total inflow to the St. Lucie Estuary averaged 460 cfs over the past week with no flow coming from Lake Okeechobee. Mean salinities decreased at all sites in the estuary over the past week. Salinity at the US1 Bridge was in the good range (10-26) for adult eastern oysters.

Total inflow to the Caloosahatchee Estuary averaged 1,970 cfs over the past week with 1,420 cfs coming from the Lake. Mean surface salinities remained the same at S-79 and Val I-75, increased at Ft. Myers, and decreased at the remaining sites in the estuary over the past week. Salinities were in the good range (0-10) for tape grass at Val I-75 and Ft. Myers. Salinities were in the good range (10-30) for adult eastern oysters at Cape Coral and Shell Point, and in the fair range at Sanibel.

Stormwater Treatment Areas

For the week ending Sunday, March 20, 2022, approximately 3,300 acre-feet of Lake Okeechobee water was delivered to the FEBs / STAs. The total amount of Lake releases sent to the FEBs / STAs in WY2022 (since May 1, 2021) is approximately 97,400 acre-feet. The total amount of inflows to the STAs in WY2022 is approximately 1,001,000 acre-feet. Most STA cells are at or near target stage, except portions of STA-5/6 cells that are drying out. STA-1E Western Flow-way is offline for the Restoration Strategies project to fill and grade Cells 5 and 7, and STA-1E Eastern Flow-way is offline for vegetation management activities related to Tropical Storm Eta. Additionally, STA-3/4 Eastern Flow-way is offline for vegetation rehabilitation/drawdown, STA-2 Flow-way 2 is offline for construction activities, and STA-5/6 Flow-way 4 is offline for vegetation management activities. Operational restrictions are in place in STA-1W Western, Eastern, and Northern Flow-ways for construction activities. Operational restrictions are in effect in STA-1E Central Flow-way and STA-2 Flow-ways 3 and 4 for vegetation management activities, and in STA-2 Flow-way 1 following dry-out conditions. This week, if 2008 LORS recommends Lake releases to the WCAs and conditions allow, releases will be sent to STA-2.

Everglades

Stage changes within most of the WCAs were optimal to only slightly suboptimal. While depths in WCA-2A remain above schedule continued discharges from that basin to WCA-3A North via the northern perimeter of that basin are benefitting the ecology of that particularly sensitive region. Wading bird foraging and nesting expectations remain at best average this year, but optimism was boosted by some unexpected nesting across

the system. Most of the eastern CSSS subpopulation percent dry conditions have reached the target of 40% dry, but recessions in sub population Echo are slow. Stages fell in Taylor Slough, but the conditions remain fair within Florida Bay as salinities rise slowly.

Biscayne Bay

Total inflow to Biscayne Bay averaged 202 cfs and the previous 30-day mean inflow averaged 229 cfs. The seven-day mean salinity was 28.0 at BBCW8 and 25.8 at BBCW10, both below the preferred maximum salinity of 35 for these sites. Salinity data provided as a courtesy by Biscayne National Park.

Supporting Information

Kissimmee Basin

Upper Kissimmee

On March 20, 2022, lake stages were 56.8 feet NGVD (1.0 feet below schedule) in East Lake Toho, 53.6 feet NGVD (1.2 feet below schedule) in Lake Toho, and 50.1 feet NGVD (0.9 feet below schedule) in Lakes Kissimmee-Cypress-Hatchineha (KCH) (**Table KB-1, Figures KB-1-3**).

Lower Kissimmee

Discharge to the Kissimmee River on March 20, 2022 was 1,020 cfs at S-65 and 940 cfs at S-65A; discharge from the Kissimmee River were 930 cfs at S-65D and 790 cfs at S-65E (**Table KB-2**). Headwater stages were 46.5 feet NGVD at S-65A and 26.6 feet NGVD at S-65D on March 20, 2022. With lower water temperatures, the concentration of dissolved oxygen is well above the region of concern, with an average of 8.3 mg/L for the week ending on March 20, 2022 (**Table KB-2, Figure KB-4**). Flow at S-65/S-65A is being increased slowly to manage the stage reversals on KCH and the Kissimmee River, and water depth on the Kissimmee River floodplain increased slightly with a mean depth of 0.29 feet as of March 20, 2022 (**Figure KB-5**).

Water Management Recommendations

Managed stage recessions for snail kite nesting season were started on Lakes Toho and East Toho on January 15, 2022 to gradually reduce lake stages to their low pools by June 1. Due to the recent stage reversals, adjust flow at S-59 and S-61 to return lake stages to their projected recession lines by about March 23. In Kissimmee-Cypress-Hatchineha, continue flow at ~1,050 cfs to control its stage reversal, then begin adjustments to resume a slow stage decline while maintaining at least 300 cfs at S-65A and following the IS-14-50 discharge plan (**Figure KB-6**) for S-65 and S-65A.

Table KB-1. Average discharge for the preceding seven days and Sunday's average daily stage and departures from KCL flood regulation or temporary schedules. All data are provisional.

Water Body	Structure	Stage Monitoring Site	7-Day Average Discharge (cfs)	Lake Stage (feet NGVD) ^a	Schedule Type ^b	Schedule Stage (feet NGVD)	Departure from Regulation (feet)	
							3/20/22	3/13/22
Lakes Hart and Mary Jane	S-62	LKMJ	219	60.5	R	60.9	-0.4	-0.4
Lakes Myrtle, Preston and Joel	S-57	S-57	70	60.8	R	60.8	0.0	0.1
Alligator Chain	S-60	ALLI	215	63.9	R	63.9	0.0	0.0
Lake Gentry	S-63	LKGT	304	61.3	R	61.4	-0.1	0.0
East Lake Toho	S-59	TOHOE	811	56.8	R	57.8	-1.0	-1.0
Lake Toho	S-61	TOHOW S-61	1,630	53.6	R	54.8	-1.2	-1.3
Lakes Kissimmee, Cypress and Hatchineha	S-65	KUB011 LKIS5B	978	50.1	R	51.0	-0.9	-1.4

a. Names of in-lake monitoring sites and structures used to determine lake stage. If more than one site is listed, an average is reported.

b. A: projected recession line; R: USACE regulation schedule; S: temporary recession target line; T: temporary schedule; NA: not applicable or not available.




Figure KB-1. East Lake Toho regulation schedule, stage, discharge and rainfall.




Figure KB-2. Lake Toho regulation schedule, stage, discharge and rainfall.




Figure KB-3. Lakes Kissimmee, Cypress and Hatchineha regulation schedule, stage, discharge and rainfall.

Table KB-2. One- and seven-day average discharge and stage at Lower Kissimmee basin structures, river channel dissolved oxygen concentrations and water depths in the Phase I area floodplain. All data are provisional.

Metric	Location	Daily Average	Average for Previous Seven Day Periods			
		3/20/22	3/20/22	3/13/22	3/6/22	2/27/22
Discharge	S-65	1,020	980	440	400	710
Discharge	S-65A ^a	940	880	410	360	630
Headwater Stage (feet NGVD)	S-65A	46.5	46.4	46.4	46.4	46.3
Discharge	S-65D ^b	930	770	380	470	950
Headwater Stage (feet NGVD)	S-65D ^c	26.6	26.6	26.5	26.5	26.5
Discharge (cfs)	S-65E ^d	790	660	310	430	870
Discharge (cfs)	S-67	0	0	0	0	0
Dissolved Oxygen (mg/L) ^e	Phase I, II/III river channel	8.5	8.3	7.9	8.0	7.6
Mean depth (feet) ^f	Phase I floodplain	0.29	0.28	0.24	0.24	0.34

a. Combined discharge from main and auxiliary structures.

b. Combined discharge from S-65D, S-65DX1 and S-65DX2.

c. Average stage from S-65D and S-65DX1.


d. Combined discharge from S-65E and S-65EX1.

e. Dissolved oxygen is the average of values from sondes KRBN, PC62, PC33, PD62R and PD42R.

f. One-day spatial average obtained from the South Florida Water Depth Assessment Tool (SFWDAT).

Table KB-3. Discharge rate of change limits for S65/S-65A (revised 1/14/19).

Discharge (cfs)	Maximum Rate of Increase (cfs/day)	Maximum Rate of Decrease (cfs/day)
0-300	100	-50
301-650	150	-75
651-1,400	300	-150
1,401-3,000	600	-600
>3,000	1,000	-2,000



Report Date: 3/22/2022; data are through: 3/20/2022



Figure KB-4. Restored Kissimmee river channel mean daily dissolved oxygen concentration (mg/L), S-65A discharge (cfs), temperature (°C) and rainfall (inches). Dissolved oxygen (DO) and temperature are mean daily values averaged for PC62, KRBN, PC33, PD62R, and PD42R with an average of three stations reporting this week. Rainfall values are daily totals for Kissimmee River (Pool BCD) AHED watershed.




Figure KB-5. Phase I area Kissimmee River floodplain water depths (from left to right) one year ago, one month ago and current.




Figure KB-6. IS-14-50 Discharge Plan for S65/S65A with discharge rate of change limits (revised 1/14/19).

Lake Okeechobee

Lake Okeechobee stage was 14.13 feet NGVD on March 20, 2022, with water levels 0.66 feet lower than a month ago (**Figure LO-1**). Lake stage remains in the Low sub-band (**Figure LO-2**) and has been within the ecological envelope for almost 12 weeks (**Figure LO-3**). According to NEXRAD, 0.92 inches of rain fell directly on the Lake last week.

Average daily inflows (excluding rainfall) increased from the previous week, going from 345 cfs to 705 cfs. Average daily outflows (excluding evapotranspiration) decreased from the previous week, going from 3,550 cfs to 2,408 cfs. Highest inflows came from the Kissimmee River through the S-65E structure (663 cfs). The highest outflow (1,524 cfs) was to the west via the S-77 structure, while flows south via the S-350 structures decreased from 1,199 cfs to 274 cfs. Flows east were 440 cfs and 171 cfs via the S-308 structure and the L-8 canal through the S-271 structure (formerly Culvert 10A), respectively. **Figures LO-4 and LO-5** show the combined average daily inflows and outflows for the Lake over the past eight weeks, and average inflows and outflows last week, respectively. These data are provisional and are subject to change.

The most recent snail kite nesting survey recorded ten more nests on Lake Okeechobee, for a total of nineteen nests thus far, most of which were found in the Indian Prairie marsh.

The most recent satellite image (March 20, 2022) from the NOAA cyanobacteria monitoring product derived from EUMETSAT's Sentinel 3 OLCI sensor showed a few scattered areas of low to moderate bloom potential, in Fisheating Bay and along the northwestern shoreline (**Figure LO-6**).

1 Month Ago:
02/18/2022

Current:
03/20/2022





Figure LO-1. Lake Okeechobee water depth estimates based on South Florida Water Depth Assessment Tool (SFWDAT).

Lake Okeechobee Water Level History and Projected Stages



LORS-2008 - Adopted by USACE 28-April-2008

Figure LO-2. Recent Lake Okeechobee stages with projected stages based on a dynamic position analysis.

Lake Okeechobee Stage vs Ecological Envelope




Figure LO-3. The prior six years of annual stage hydrographs for Lake Okeechobee in comparison to the ecological envelope.




Figure LO-4. Major inflows (orange) to and outflows east and west (blue) from Lake Okeechobee. Outflows south are shown in green. Flows into Lake Okeechobee from the L-8 canal through S-271 (formerly Culvert 10A) or from the C-44 canal through the S-308 are included as inflows. Conversely, flows from Lake Okeechobee into the L-8 or C-44 canals are included with outflows. Inflows are shown as positive values; outflows are negative. Outflows through the S-77 (Caloosahatchee) and S-308 (C-44 Canal) structures are based on downstream gauges to include flows to lock openings for navigation.




Figure LO-5. Inflows into Lake Okeechobee from Indian Prairie basins, Taylor Creek/Nubbin Slough, Kissimmee River and Fisheating Creek, and outflows to the west via S-77, to the east via S-308, to the south via S-351, S-352, S-354, and to southeast via S-271 (formerly Culvert 10A) for the week of March 14, 2022 – March 20, 2022.




Figure LO-6. Cyanobacteria bloom potential on March 20, 2022 based on NOAA's harmful algal bloom monitoring system. Gray color indicates cloud cover.

Estuaries

St. Lucie Estuary

Over the past week, mean total inflow to the St. Lucie Estuary was 458 cfs (**Figures ES-1 and ES-2**) and the previous 30-day mean inflow was 312 cfs. For comparison, the historical provisional mean inflows from the contributing areas are shown in **Figure ES-2**.

Over the past week, salinities decreased at all sites within the estuary (**Table ES-1 and Figure ES-3**). The seven-day moving average of the surface and bottom salinities at the US1 Bridge was 20.4. Salinity conditions in the middle estuary were estimated to be within the good range for adult eastern oysters (**Figure ES-4**). Larval oyster recruitment rates reported by the Florida Fish and Wildlife Research Institute were low in February, as expected during the cooler months when most oysters are not spawning (**Figure ES-5**).

Caloosahatchee River Estuary

Over the past week, mean total inflow to the Caloosahatchee River Estuary was 1,973 cfs (**Figures ES-6 and ES-7**) and the previous 30-day mean inflow was 2,034 cfs. For comparison, the historical provisional mean inflows from the contributing areas are shown in **Figure ES-7**.

Over the past week, surface salinities remained the same at S-79 and Val I-75, increased at Ft. Myers, and decreased at the remaining sites in the estuary (**Table ES-2 and Figures ES-8 and ES-9**). The seven-day mean surface salinities (**Table ES-2**) were in the good range (0-10) for tape grass at Val I-75 and at Ft. Myers. The seven-day mean surface salinity values were within the good range for adult eastern oysters at Cape Coral and Shell Point, and in the fair range at Sanibel (**Figure ES-10**). Larval oyster recruitment rates reported by the Fish and Wildlife Research Institute were low at Iona Cove and Bird Island in February, as expected during the cooler months when most oysters are not spawning (**Figures ES-11 and ES-12**).

Surface salinity at Val I-75 was forecasted for the next two weeks, using an autoregression model (Qiu and Wan, 2013¹) coupled with a linear reservoir model for the tidal basin. Model scenarios included pulse releases at S-79 ranging from 0 to 1500 cfs and steady releases at 2,000 cfs with estimated tidal basin inflows of 53 cfs. Model results from all scenarios predict daily salinity to be 1.6 or lower and the 30-day moving average surface salinity to be 0.5 or lower at Val I-75 at the end of the two-week period (**Table ES-3 and Figure ES-13**). This keeps predicted salinities at Val I-75 within the LORS 2008 salinity range (0.0-5.0).

¹ Qiu, C., and Y. Wan. 2013. Time series modeling and prediction of salinity in the Caloosahatchee River Estuary. *Water Resources Research* 49:5804-5816.

Red Tide

The Florida Fish and Wildlife Research Institute reported on March 18, 2022, that *Karenia brevis*, the Florida red tide dinoflagellate, was not observed in any samples collected statewide. On the east coast, red tide was not observed in samples collected from Palm Beach County.

Water Management Recommendations

Lake stage is in the Low Sub-Band. Tributary conditions are dry. The LORS2008 release guidance suggests up to 450 cfs release at S-79 to the Caloosahatchee River Estuary and up to 200 cfs release at S-80 to the St. Lucie Estuary.




Figure ES-1. Basins, water control structures and salinity monitoring sites in the St. Lucie Estuary.




Figure ES-2. Total daily inflows from Lake Okeechobee and runoff from the C-44, C-23, C-24, Ten Mile Creek, and Tidal Basins into the St. Lucie Estuary.

Table ES-1. Seven-day mean salinity at oyster monitoring sites in the St. Lucie Estuary. Current means are in bold font; previous week's means are in parentheses. The envelope reflects the preferred salinity range for adult eastern oysters (*Crassostrea virginica*) in the middle estuary. Data are provisional.

Sampling Site	Surface	Bottom	Envelope
HR1 (North Fork)	13.1 (17.6)	16.9 (20.2)	NA ^a
US1 Bridge	20.0 (22.6)	20.9 (23.3)	10.0 – 26.0
A1A Bridge	27.7 (28.7)	29.6 (30.1)	NA ^a

a. The envelope is not applicable.




Figure ES-3. Mean daily salinity at the A1A, US1 and HR1 sites in the St. Lucie Estuary.




Figure ES-4. Seven-day moving average of the surface and bottom salinities at the US1 Bridge in the St. Lucie Estuary.




Figure ES-5. Mean oyster recruitment at the Rio oyster monitoring station and 14-day mean salinity at US1 Bridge.




Figure ES-6. Basins, water control structures and salinity monitoring sites in the Caloosahatchee River Estuary.




Figure ES-7. Total daily inflows from Lake Okeechobee, and runoff from the C-43 and Tidal basins into the Caloosahatchee River Estuary.

Table ES-2. Seven-day mean salinity at six monitoring sites in the Caloosahatchee River Estuary. Current means are in bold font; previous week's means are in parentheses. The envelope at I-75 is for the protection of tape grass in the upper estuary and the envelope in the lower estuary is the preferred salinity range for adult eastern oysters (*Crassostrea virginica*). Data are provisional.

Sampling Site	Surface	Bottom	Envelope
S-79 (Franklin Lock)	0.2 (0.2)	0.2 (0.2)	NA ^a
Val I-75	0.2 (0.2)	0.2 (0.2)	0.0 – 5.0 ^b
Fort Myers Yacht Basin	2.4 (2.1)	3.8 (2.5)	NA ^a
Cape Coral	11.1 (11.7)	13.3 (13.0)	10.0 – 30.0
Shell Point	26.2 (27.0)	27.0 (27.6)	10.0 – 30.0
Sanibel	31.2 (31.9)	31.9 (32.3)	10.0 – 30.0

a. The envelope is not applicable.

b. The envelope is based on the predicted 30-day mean for the next two weeks.




Figure ES-8. Mean daily salinity at upper Caloosahatchee River Estuary monitoring sites and mean daily flow at S-79.




Figure ES-9. Mean daily surface salinity at lower Caloosahatchee River Estuary monitoring sites and mean daily flow at S-79.




Figure ES-10. Seven-day moving average of surface and bottom salinities at Cape Coral, Shell Point and Sanibel monitoring sites in the Caloosahatchee River Estuary.




Figure ES-11. Mean oyster recruitment at the Iona Cove oyster monitoring station and 14-day mean salinity at Cape Coral.




Figure ES-12. Mean oyster recruitment at the Bird Island oyster monitoring station and 14-day mean salinity at Shell Point.

Table ES-3. Predicted salinity at Val I-75 in the Caloosahatchee River Estuary at the end of the forecast period for various S-79 flow release scenarios.

Scenario	Simulated S-79 Flow (cfs)	Tidal Basin Runoff (cfs)	Daily Salinity	30-Day Mean Salinity
A	0	53	1.6	0.5
B	450	53	0.8	0.4
C	800	53	0.4	0.4
D	1000	53	0.3	0.4
E	1500	53	0.3	0.4
F	2000	53	0.3	0.3




Figure ES-13. Forecasted Val I-75 site surface salinity assuming no pulse release at S-79.

Stormwater Treatment Areas

STA-1E: STA-1E Western Flow-way is offline for the Restoration Strategies project to fill and grade Cells 5 and 7, and the Eastern Flow-way is offline for vegetation management activities related to Tropical Storm Eta. Operational restrictions are in place in STA-1E Central Flow-way for vegetation management activities. Online treatment cells are at or near target stage and vegetation in the flow-ways is stressed and highly stressed. The 365-day phosphorus loading rates (PLRs) is high for the Central Flow-way (**Figure S-1**).

STA-1W: Operational restrictions are in place in STA-1W Western, Eastern, and Northern Flow-ways due to construction activities. Most treatment cells are at or near target stage. Vegetation in the flow-ways is stressed and highly stressed. The 365-day PLRs for the Northern and Western, and Eastern Flow-ways are $1.0 \text{ g/m}^2/\text{year}$ or lower (**Figure S-2**).


STA-2: STA-2 Flow-way 2 is offline for construction activities. Operational restrictions are in place in STA-2 Flow-ways 3 and 4 for vegetation management activities, and in Flow-way 1 following dryout conditions. Online treatment cells are near or below target stage. Vegetation in Flow-ways 1 and 3 is stressed, and in Flow-ways 2, 4 and 5 is highly stressed. The 365-day PLRs for Flow-ways 1, 4 and 5 are below $1.0 \text{ g/m}^2/\text{year}$. The 365-day PLR for Flow-way 3 is high (**Figure S-3**).

STA-3/4: STA-3/4 Eastern Flow-way is offline for vegetation rehabilitation/drawdown. Most online treatment cells are at or near target stage. Vegetation in the Eastern and Central Flow-ways is highly stressed and in the Western Flow-way is stressed. The 365-day PLRs for the Central and Western Flow-ways are below $1.0 \text{ g/m}^2/\text{year}$ (**Figure S-4**).

STA-5/6: STA-5/6 Flow-way 4 is offline for vegetation management activities. Most treatment cells are below target stage. All treatment cells have highly stressed vegetation conditions except Flow-ways 7 and 8 which are healthy. The 365-day PLRs for flow-ways 1, 6, 7, and 8 are below $1.0 \text{ g/m}^2/\text{year}$. The 365-day PLRs for flow-ways 4 and 5 are high. (**Figure S-5** and **S-6**).

For definitions on STA operational language see glossary following figures.

STA-1E Weekly Status Report – 3/14/2022 through 3/20/2022



















STA-1E Flow-Way Status				As of 3/20/2022		STA-1E Flow & Phosphorus Concentration			
Flow-Way	Vegetation Status <small>Healthy ----- Stressed</small> 	365-day P Loading Rate <small>(below 1.0 g P /m²/yr is optimal)</small>	Online / Offline / Restrictions	Stage Based: Relative to Target Stage (TS)			7-day	28-day	365-day
				 Deep Water Level (> 2.8' above TS)	 High Water Level (1.5' – 2.8' above TS)				
Eastern			Offline, Tropical Storm Eta repairs starting 01/24/2022	 0.2' – 1.5' above TS	 Target Stage (TS +/- 0.2')		266	1,804	178,117
				 Low Water Level (<0.2' below TS)					
Central			Vegetation Rehab	Depth / Area Based: Percent of Area Dry					
Western			Offline, construction activities starting 11/01/2019	 0-25% Dry	 50-75% Dry		78	75	122
				 25-50% Dry	 75-100% Dry				
							</		

Figure S-1. STA-1E Weekly Status Report

STA-1W Weekly Status Report – 3/14/2022 through 3/20/2022



STA-1W Flow-Way Status

Flow-Way	Vegetation Status Healthy --- Stressed	365-day P Loading Rate (below 1.0 g P /m ² /yr is optimal)	Online / Offline / Restrictions
Northern			Construction
Western			Construction
Eastern			Construction
Cell 7		N/A	Construction
Cell 8		N/A	Construction


As of 3/20/2022	
Stage Based: Relative to Target Stage (TS)	
	Deep Water Level (> 2.8' above TS)
	High Water Level (1.5' – 2.8' above TS)
	0.2' – 1.5' above TS
	Target Stage (TS +/- 0.2')
	Low Water Level (<0.2' below TS)
Depth / Area Based: Percent of Area Dry	
	0-25% Dry
	25-50% Dry
	50-75% Dry
	75-100% Dry









STA-1W Flow & Phosphorus Concentration










	7-day	28-day	365-day
Total Inflow, ac-ft	799	3,371	57,503
--Lake Inflow, ac-ft	0	N/A	5,400
Total Outflow, ac-ft	0	0	52,974
Inflow Conc., ppb	76	80	162
Outflow Conc., ppb	N/A	N/A	24
Includes Preliminary Data			

Figure S-2. STA-1W Weekly Status Report

STA-2 Weekly Status Report – 3/14/2022 through 3/20/2022




STA-2 Flow-Way Status			
Flow-Way	Vegetation Status Healthy ----- Stressed	365-day P Loading Rate (below 1.0 g P /m ² /yr is optimal)	Online / Offline / Restrictions
1	←  →		Post Dryout
2	Offline, construction activities as of 9/7/2021		
3	←  →		Vegetation Rehab
4	←  →		Vegetation Rehab
5	←  →		Online

As of 3/20/2022	
Stage Based: Relative to Target Stage (TS)	
	Deep Water Level (> 2.8' above TS)
	High Water Level (1.5' – 2.8' above TS)
	0.2' – 1.5' above TS
	Target Stage (TS +/- 0.2')
	Low Water Level (<0.2' below TS)
Depth / Area Based: Percent of Area Dry	
	0-25% Dry
	25-50% Dry
	50-75% Dry
	75-100% Dry

STA-2 Flow & Phosphorus Concentration			
	7-day	28-day	365-day
Total Inflow, ac-ft	3,176	8,828	314,857
--Lake Inflow, ac-ft	1,700	N/A	58,300
Total Outflow, ac-ft	1,165	4,898	328,378
Inflow Conc., ppb	72	66	89
Outflow Conc., ppb	13	14	15
Includes Preliminary Data			

Figure S-3. STA-2 Weekly Status Report


STA-3/4 Weekly Status Report – 3/14/2022 through 3/20/2022



STA-3/4 Flow-Way Status				STA-3/4 Flow & Phosphorus Concentration			
Flow-Way	Vegetation Status Healthy ----- Stressed	365-day P Loading Rate (below 1.0 g P /m ² /yr is optimal)	Online / Offline / Restrictions	As of 3/20/2022			
				Stage Based: Relative to Target Stage (TS)			
Eastern	Offline, vegetation management drawdown as of 3/1/2021			<div>Deep Water Level (> 2.8' above TS)</div> <div>High Water Level (1.5' – 2.8' above TS)</div> <div>0.2' – 1.5' above TS</div> <div>Target Stage (TS +/- 0.2')</div> <div>Low Water Level (<0.2' below TS)</div>			
				Depth / Area Based: Percent of Area Dry			
Central	← Healthy →	1.0	Online	0-25% Dry	50-75% Dry		
Western	← Healthy →	1.0	Online	25-50% Dry	75-100% Dry		
				Includes Preliminary Data			
				<div>Total Inflow, ac-ft</div> <div>--Lake Inflow, ac-ft</div> <div>Total Outflow, ac-ft</div> <div>Inflow Conc., ppb</div> <div>Outflow Conc., ppb</div>			
				7-day	28-day	365-day	
				28	29	358,565	
				0	N/A	37,900	
				79	84	316,363	
				31	30	65	
				9	9	15	

Figure S-4. STA-3/4 Weekly Status Report

STA-5/6 Weekly Status Report – 3/14/2022 through 3/20/2022




STA-5/6 Flow-Way Status			
Flow-Way	Vegetation Status Healthy ----- Stressed	365-day P Loading Rate (below 1.0 g P /m ² /yr is optimal)	Online / Offline / Restrictions
1			Online
2		N/A	Online
3		N/A	Online
4	Offline, vegetation management starting 01/24/2022		
5			Online

As of 3/20/2022	
Stage Based: Relative to Target Stage (TS)	
	Deep Water Level (> 2.8' above TS)
	High Water Level (1.5' – 2.8' above TS)
	0.2' – 1.5' above TS
	Target Stage (TS +/- 0.2')
	Low Water Level (<0.2' below TS)
Depth / Area Based: Percent of Area Dry	
	0-25% Dry
	25-50% Dry
	50-75% Dry
	75-100% Dry

STA-5/6 Flow & Phosphorus Concentration			
	7-day	28-day	365-day
Total Inflow, ac-ft	0	30	177,781
--Lake Inflow, ac-ft	0	N/A	8,800
Total Outflow, ac-ft	0	11	169,236
Inflow Conc., ppb	N/A	67	243
Outflow Conc., ppb	N/A	22	50
Includes Preliminary Data			

Figure S-5. STA-5/6 Weekly Status Report (Flow-ways 1 – 5)

STA-5/6 Weekly Status Report – 3/14/2022 through 3/20/2022



STA-5/6 Flow-Way Status				As of 3/20/2022	
Flow-Way	Vegetation Status Healthy ----- Stressed ←-----→	365-day P Loading Rate (below 1.0 g P /m ² /yr is optimal)	Online / Offline / Restrictions	Stage Based: Relative to Target Stage (TS)	
				<div>Deep Water Level (> 2.8' above TS)</div> <div>High Water Level (1.5' – 2.8' above TS)</div> <div>0.2' – 1.5' above TS</div> <div>Target Stage (TS +/- 0.2')</div> <div>Low Water Level (<0.2' below TS)</div>	
6	←-----→	1.0	Online		
7	←-----→	1.0	Online		
8	←-----→	1.0	Online		
				Depth / Area Based: Percent of Area Dry	
				0-25% Dry	50-75% Dry
				25-50% Dry	75-100% Dry

Figure S-6. STA-5/6 Weekly Status Report (Flow-ways 6 – 8)

Basic Concepts and Definitions for STA Weekly Status Report

- **Inflow:** Sum of flow volume at all inflow structures to an STA.
- **Lake Inflow:** Portion of the STA total inflow volume that originates from Lake Okeechobee.
- **Outflow:** Sum of flow volume at outflow structures from an STA.
- **Total Phosphorus (TP):** Total mass of phosphorus in all its forms; including particulate, dissolved, etc.
- **Inflow Concentration:** TP concentration is the mass of TP in micrograms per liter of water, $\mu\text{g/L}$ or ppb. Inflow concentration refers to the flow-weighted mean TP from all inflow structures over a period of time.
- **Outflow Concentration:** The flow-weighted mean TP from all outflow structures over a period of time. The outflow concentration represents the reduction of inflow TP achieved by STA treatment of the inflow water.
- **WQBEL:** The STA outflow concentration that is required upon completion of the Restoration Strategies projects by December 2025. The outflow concentration shall not exceed 13 ppb as an annual flow weighted mean in more than 3 out of 5 water years on a rolling basis and shall not exceed 19 ppb as an annual flow weighted in any water year.
- **Flow-Way (FW):** One or more treatment cells connected in series. Cells typically have emergent aquatic vegetation (EAV) in the front portion of the flow-way followed by a mix of EAV and submerged aquatic vegetation (SAV)
- **Vegetation Status:** Healthy means the vegetation condition is good and will allow the STA to perform as designed. Stressed means the vegetation is showing signs of poor health, such as browning or areas of vegetation die-off, or the cell contains undesirable vegetation such as floating exotic vegetation requiring treatment. The TP reduction capability of the STA is affected when the vegetation condition is poor.
- **Phosphorus Loading Rate (PLR):** Mass of inflow TP in grams, divided by total treatment area of STA in square meters, per year. In general, a 365-day value of less than 1.0 is needed for an STA to perform optimally. A PLR of 2.0 is considered very high and a PLR of 3.0 is considered extremely high. The TP reduction capability of the STA is affected when the PLR is high, very high and extremely high.
- **Online:** Online status means the FW can receive and treat inflow.
- **Online with Restriction:** The FW can receive and treat inflow, but the amount of flow or water level may be limited temporarily. For example, a vegetation rehabilitation effort may require reduced flows through an area while the new plants are establishing, or nesting by protected species may require a certain water level not to be exceeded.
- **Offline:** The FW is unable to receive and treat inflow due to repairs, construction, or other prohibitive reasons.
- **Depth:** Difference between the average surface water level in a cell and the average ground elevation in that cell. Target depths, or depths between flow events, are between 1.25 ft to 1.5 ft. As depth approaches or drops below zero, an increasing percentage of the cell is considered dry and STA conditions deteriorate. An increase in depth above target depth is expected with increasing flow. However, as depth increases much above the target depth and is sustained over a period of time, it can be detrimental to vegetation health and overall STA treatment performance.
- **Note:** The data provided in this summary report were developed using a combination of provisional and quality-assured flow and water quality data. In some cases, best professional judgment was used to estimate missing data and revise questionable data. Values provided are not considered final but are appropriate for use in STA operational decision-making.

Everglades

Water Conservation Area Regulation Schedules

WCA-1: The 1-8C recession slowed again. The average on Sunday was 0.24 feet above the falling Zone A1 regulation line. WCA-2A: Stage continued recession at S11B headwater last week. The average at that gauge on Sunday was 0.87 feet higher than the flat regulation line. WCA-3A: Last week the Three Gauge Average stages increased as a result of rain; average stage was 0.83 feet (0.94 feet the previous week) below the falling regulation line on Sunday. WCA-3A: Stage continued recession at gauge 62 (Northwest corner) last week, the average on Sunday was 0.98 feet below the flat Upper schedule line. (**Figures EV-1 through EV-4**).

Water Depths

The SFWDAT tool indicates that water depths have been showing a fairly stable recession in WCAs 1 and 2 over the last two months while conditions further south are drying out more rapidly. The effects of the water inputs to northeastern WCA-3A are now visible as slightly increased water levels. North to South hydrologic connectivity continues to diminish but remains within Everglades National Park's Shark River Slough. Stages are well below ground across southern BCNP as is typical but not favorable for this time of year (**Figure EV-5**). Comparing current WDAT water depths to the depth one month ago, stages are decreasing slowly across most of the EPA. Looking back one year, most of the EPA south of WCA-2A is lower in depth compared to one year ago. Northeastern WCA-3A central BCNP are slightly wetter than a year ago (**Figure EV-6**). Comparing current depths to the past 20 years, the eastern half WCA-3A is below the 20th percentile. This past week's rainfall shows an immediate impact over much of the area with at least half of the EPA now above the median depths (**Figure EV-7**).

Taylor Slough and Florida Bay

A spatial average of 0.73 inches of rain fell over Taylor Slough and Florida Bay during the week ending Sunday, 3/20. Stages in Taylor Slough decreased an average of 0.09 feet over this past week with the largest weekly change of -0.20 feet in the northern Taylor Slough area (**Figure EV-8**). The Slough, as a whole, is still 8 inches higher than average while the northern parts are 14 inches higher than the historical average for this time of year (pre-Florida Bay initiative which started in 2017). Taylor Slough is still on track to be mostly dry in approximately 2 weeks (**Figure EV-9**) which is a month later than what typically occurred in the past suggesting that water management this year has been able to expand the hydroperiod in the Slough despite the drier than average conditions. Slowing the recession in the Slough will allow water movements towards the Bay to be expedited once the wet season begins

Salinities in Florida Bay averaged an increase of 0.7 over the week ending 3/20, with individual station changes ranging from -0.8 to +2.9 (**Figure EV-8**). The nearshore stations increased while the Bay stations showed slight decreases over the week. Weekly flow from the 5 main creeks was about -2,000 acre-feet. The average salinities in all 3 regions of the Bay are increasing and in the middle of their interquartile ranges (**Figure**

EV-10). Current condition across the Bay is averaging 1.4 psu higher than the long-term mean for this time of year.

Water Management Recommendations

Conserving water in the northern basins, then allowing that water to move downstream as we transition into the dry season maximizes the ecological benefit of freshwater on the landscape. This recommendation is currently being epitomized with WCA-1 and STA2 discharges hydrating northern WCA-2A, being picked up by S7 which then supplies NE and NW WCA-3A North. This operation is more ecologically advantageous than outflow from the S-11 structures as long as a recession rate can be maintained in WCA-2A.

Recent inflows via the S-150 were anecdotally confirmed once again this week from the air as having a positive result on stages in northeastern WCA-3A. If conditions at all allow operational discharges into both the western and the eastern WCA-3A water control structures, this operation has greater benefit than discharges to the west alone. Continued freshwater to the Taylor Slough area, a discharge amount that maintains stage will help expedite deliveries to the south when the wet season begins. Individual regional recommendations can be found in **Table EV-2**.

Table EV-2. Previous week's rainfall and water depth changes in Everglades basins.

Everglades Region	Rainfall (inches)	Stage change (feet)
WCA-1	0.25	-0.06
WCA-2A	0.30	-0.10
WCA-2B	0.60	-0.11
WCA-3A	1.36	+0.05
WCA-3B	0.62	-0.04
ENP	0.92	-0.07




Figure EV-1. WCA-1 stage hydrographs and regulation schedule.




Figure EV-2. WCA-2A stage hydrographs and regulation schedule.




Figure EV-3. WCA-3A stage hydrographs (three-gauge average, S-333 headwater) and regulation schedule.




Figure EV-4. WCA-3A stage hydrograph (Deer gauge; Site 62) and CA62 regulation schedule.




Figure EV-5. Everglades water depths from two months ago (left), one month ago (center) and present (right), based on SFWDAT.




Figure EV-6. Present Everglades water depths (left) and water depth changes from one month (center) and one year (right) ago, based on SFWDAT.




Figure EV-7. Present water depths compared to the day of year median over the previous 20 years.




Figure EV-8. Taylor Slough water depths with changes since a week ago and Florida Bay salinities with changes since a week ago.

SFWMD Everglades Ecological Recommendations, February 22, 2022 (red is new)

Area	Weekly change	Recommendation	Reasons
WCA-1	Stage decreased by 0.06'	Allow water to move south from this basin until stages reach the regulation schedule. Maintain a recession rate of less than 0.10 feet per week.	Protect within basin and downstream habitat and wildlife. Stages are above the 90 th percentile.
WCA-2A	Stage decreased by 0.10'	Conserve water in this basin letting the water move south when conditions allow, with northern WCA-3A as the priority for receiving discharge. A recession rate less than 0.10 feet per week has an ecological benefit.	Protect within basin and downstream habitat and wildlife. Protect peat soil and future wading bird forage as the dry season progresses.
WCA-2B	Stage decreased by 0.11'	Conserve water in this basin, maintain a minimum input to maintain stage while moving water south when conditions allow.	Protect within basin and downstream habitat and wildlife.
WCA-3A NE	Stage had a net zero change	Conserve water in this basin, while letting the water move south when conditions allow. Returning to a recession rate less than 0.10 feet per week has an ecological benefit.	Protect within basin peat soils, and downstream habitat and wildlife. Inflows via the S-150 are anecdotally observed to be positively impacting stages in NE WCA-3A North.
WCA-3A NW	Stage decreased by 0.04'	Conserve water in this basin letting the water move south when conditions allow.	
Central WCA-3A S	Stage increased by 0.04'	Return to a recession rate that is less than 0.10 feet per week. Allow flows to move south as conditions allow.	Protect within basin and downstream habitat and wildlife.
Southern WCA-3A S	Stage increased by 0.20'		
WCA-3B	Stage decreased by 0.04'	Return to a recession rate of less than 0.10 feet per week in this basin, letting the water move south when conditions allow.	Protect within basin and downstream habitat and wildlife.
ENP-SRS	Stage decreased by 0.07'	Make discharges to ENP according to COP and TTFF protocol while adaptively considering upstream and downstream ecological conditions.	Protect within basin and upstream habitat and wildlife.
Taylor Slough	Stage changes ranged from -0.04' to -0.21'	Move water southward as possible.	When available, provide freshwater buffer for downstream conditions.
FB- Salinity	Salinity changes ranged -0.8 to +2.9	Move water southward as possible.	When available, provide freshwater to maintain low salinity buffer and promote water movement.

Biscayne Bay

As shown in **Figure BB-1**, mean total inflow to Biscayne Bay was 202 cfs and the previous 30-day mean inflow was 229 cfs. The seven-day mean salinity was 28.0 at BBCW8 and 25.8 at BBCW10, both below the preferred maximum salinity of 35 for these sites. Data provided by Biscayne National Park.




Figure BB-1. Seven-day mean salinity at BBCW8 and BBCW10 and total daily flow in Biscayne Bay. Total daily flow was calculated using flow from structures S20G, S20F, S21, S21A, S123, and S700P.