

Disclaimer: Information contained in the report addresses environmental conditions only and is not the official South Florida Water Management District operations recommendation or decision.

M E M O R A N D U M

TO: John Mitnik, Assistant Executive Director, Executive Office Staff

FROM: SFWMD Staff Environmental Advisory Team

DATE: April 6, 2022

SUBJECT: Weekly Environmental Conditions for Systems Operations

Summary

Weather Conditions and Forecast

A stationary front moved north of the Lake early in the week leading to afternoon thunderstorms over the interior of the District from Lake Okeechobee to the Kissimmee Valley. Precipitation will maximize where low-level flow converges with the east coast sea breeze boundary. However, due to accelerated storm motion, peak rainfall values are likely to be much lower later in the week. On Wednesday and Thursday, afternoon thunderstorms are likely over the northern part of the District and along the east coast in relation to the next frontal system. On Friday, a frontal passage is expected, and there could be light rain along the southeast coast. A dry period is expected to begin this weekend. Much below average District-wide rainfall is expected for the first 7-day period ending next Monday.

Kissimmee

Flow at S-59 and S-61 was increased to slow the rise in stage in East Toho and Toho above their respective recession lines. Flow at S-65/S-65A is being increased slowly to manage the stage reversals on Kissimmee-Cypress-Hatchineha and the Kissimmee River; water depth on the Kissimmee River floodplain remained steady, with a mean depth of 0.23 feet as of April 3, 2022. The concentration of dissolved oxygen in the Kissimmee River has remained well above the region of concern, with an average of 8.1 mg/L for the week ending on April 3, 2022.

Lake Okeechobee

Lake Okeechobee stage was 13.75 feet NGVD on April 3, 2022, with water levels 0.66 feet lower than a month ago. Lake stage has been within the ecological envelope for over 13 weeks, having spent a total of 279 days (79%) in 2021 above the envelope. Average daily inflows (excluding rainfall) increased slightly from the previous week; 843 cfs to 881 cfs. Average daily outflows (excluding evapotranspiration) decreased from 4,322 cfs to 3,859 cfs. Thirty-one snail kite nests have been recorded on the Lake thus far. Recent satellite imagery (March 28, 2022) showed a few scattered areas of low to moderate bloom potential in Fisheating Bay and along the northwestern shoreline.

Estuaries

Total inflow to the St. Lucie Estuary averaged 755 cfs over the past week with no flow coming from Lake Okeechobee. Mean salinities increased at all sites within the estuary over the past week. Salinity at the US1 Bridge was in the good range (10-26) for adult eastern oysters. Total inflow to the Caloosahatchee Estuary averaged 1,930 cfs over the past week with 1,510 cfs coming from the Lake. Mean surface salinities remained the same at S-79, Val I-75, and Shell Point, and decreased slightly at the remaining sites in the estuary over the past week. Salinities were in the good range (0-10) for tape grass at Val I-75 and Ft. Myers. Salinities were in the good range (10-30) for adult eastern oysters at Cape Coral and Shell Point and in the fair range at Sanibel. Total inflow to Biscayne Bay averaged 265 cfs, and the previous 30-day mean inflow averaged 267 cfs. The seven-day mean salinity was 27.1 at BBCW8 and 29.8 at BBCW10, both below the preferred maximum salinity of 35 for these sites. Salinity data are provided as a courtesy by Biscayne National Park.

Stormwater Treatment Areas

Approximately 100 ac-ft of Lake Okeechobee water was delivered to the FEBs/STAs over the past week. Lake releases sent to the FEBs/STAs in WY2022 (since May 1, 2021) is approximately 104,200 ac-feet. The total inflows to the STAs in WY2022 is approximately 1,013,000 ac-feet. Most STA cells are at or near target stage, except portions of STA-5/6 cells that are drying out. STA-1E Western Flow-way is offline for post-construction vegetation grow in. STA-1E Eastern Flow-way is offline for vegetation management activities related to Tropical Storm Eta. STA-3/4 Eastern Flow-way is offline for vegetation rehabilitation/drawdown, STA-2 Flow-way 2 is offline for construction activities, and STA-5/6 Flow-way 4 is offline for vegetation management activities. Operational restrictions are in place in STA-1W Western, Eastern, and Northern Flow-ways for construction activities, in STA-1E Central Flow-way and STA-2 Flow-ways 3 and 4 for vegetation management activities, and in STA-2 Flow-way 1 following dryout conditions. This week, if 2008 LORS recommends Lake releases to the WCAs and conditions allow, releases will be sent to STA-2.

Everglades

Stage changes within most of the WCAs are now fair. While depths in WCA-2A remain above schedule, continued discharges from that basin into WCA-3A North via the northern perimeter are benefitting the ecology of that particularly sensitive region. But, as stages fall at the S-11B headwaters, those discharges are expected to cease. FWC reports the WCA-3B wildfire is out, with about 10,000 acres burned with no negative impacts detected so far. Wading bird nesting expectations are average this year. Conditions are becoming appropriate for foraging in WCA-1 and WCA-2, but overall, the nesting effort is relatively low for this time of year. Wood Storks have initiated nesting very late. The eastern Cape Sable Seaside Sparrow subpopulation percent dry conditions have reached the target of 40% dry, but recessions in subpopulation Echo remain slow. Stages fell in Taylor Slough, especially in the north, and the salinities within Florida Bay increased with negative creek flow increasing salinities within the mangrove ecotone.

Supporting Information

Kissimmee Basin

Upper Kissimmee

On April 3, 2022, lake stages were 56.7 feet NGVD (0.6 feet below schedule) in East Lake Toho, 53.4 feet NGVD (0.9 feet below schedule) in Lake Toho, and 50.5 feet NGVD (0.4 feet below schedule) in Lakes Kissimmee-Cypress-Hatchineha (KCH) (**Table KB-1, Figures KB-1-3**).

Lower Kissimmee

Discharges to the Kissimmee River on April 3, 2022 were 1,190 cubic feet per second (cfs) at S-65 and 1,160 cfs at S-65A; discharges from the Kissimmee River were 1,020 cfs at S-65D and 880 cfs at S-65E (**Table KB-2**). Headwater stages were 46.4 feet NGVD at S-65A and 26.9 feet NGVD at S-65D on April 3, 2022. Although water temperature has begun to rise, the concentration of dissolved oxygen is well above the region of concern, with an average of 8.1 mg/L for the week ending on April 3, 2022 (**Table KB-2, Figure KB-4**). Flow at S-65/S-65A is being increased slowly to manage the stage reversals on KCH and the Kissimmee River, and water depth on the Kissimmee River floodplain has remained steady with a mean depth of 0.23 feet as of April 3, 2022 (**Figure KB-5**).

Water Management Recommendations

Following the most recent rainfall, flow is being increased at S-59, S-61 and S-65 to address the resulting stage reversals.

Table KB-1. Average discharge for the preceding seven days and Sunday's average daily stage and departures from Kissimmee Chain of Lakes flood regulation or temporary schedules. All data are provisional.

Water Body	Structure	Stage Monitoring Site	7-Day Average Discharge (cfs)	Lake Stage (feet NGVD) ^a	Schedule Type ^b	Schedule Stage (feet NGVD)	Departure from Regulation (feet)	
							4/3/22	3/27/22
Lakes Hart and Mary Jane	S-62	LKMJ	178	60.7	R	60.7	0.0	-0.4
Lakes Myrtle, Preston and Joel	S-57	S-57	48	60.9	R	60.6	0.3	0.0
Alligator Chain	S-60	ALLI	195	63.8	R	63.5	0.3	0.0
Lake Gentry	S-63	LKGT	364	61.5	R	61.0	0.5	-0.1
East Lake Toho	S-59	TOHOE	422	56.7	R	57.3	-0.6	-1.1
Lake Toho	S-61	TOHOW S-61	858	53.4	R	54.3	-0.9	-1.4
Lakes Kissimmee, Cypress and Hatchineha	S-65	KUB011 LKIS5B	1,145	50.5	R	50.9	-0.4	-0.7

a. Names of in-lake monitoring sites and structures used to determine lake stage. If more than one site is listed, an average is reported.

b. A: projected recession line; R: USACE regulation schedule; S: temporary recession target line; T: temporary schedule; NA: not applicable or not available.




Figure KB-1. East Lake Toho regulation schedule, stage, discharge and rainfall.




Figure KB-2. Lake Toho regulation schedule, stage, discharge and rainfall.




Figure KB-3. Lakes Kissimmee, Cypress and Hatchineha regulation schedule, stage, discharge and rainfall.

Table KB-2. One- and seven-day average discharge and stage at Lower Kissimmee basin structures, river channel dissolved oxygen concentrations and water depths in the Phase I area floodplain. All data are provisional.

Metric	Location	Daily Average 4/3/22	Average for Previous Seven Day Periods			
			4/3/22	3/27/22	3/20/22	3/13/22
Discharge	S-65	1,190	1,140	1,050	980	440
Discharge	S-65A ^a	1,160	1,080	980	880	410
Headwater Stage (feet NGVD)	S-65A	46.4	46.3	46.4	46.4	46.4
Discharge	S-65D ^b	1,020	980	950	770	380
Headwater Stage (feet NGVD)	S-65D ^c	26.9	26.8	26.6	26.6	26.5
Discharge (cfs)	S-65E ^d	880	860	830	660	310
Discharge (cfs)	S-67	0	0	0	0	0
Dissolved Oxygen (mg/L) ^e	Phase I, II/III river channel	7.4	8.1	8.3	8.3	7.9
Mean depth (feet) ^f	Phase I floodplain	0.23	0.24	0.28	0.28	0.24

a. Combined discharge from main and auxiliary structures.

b. Combined discharge from S-65D, S-65DX1 and S-65DX2.

c. Average stage from S-65D and S-65DX1.


d. Combined discharge from S-65E and S-65EX1.

e. Dissolved oxygen is the average of values from sondes KRBN, PC62, PC33, PD62R and PD42R.

f. One-day spatial average obtained from the South Florida Water Depth Assessment Tool (SFWDAT).

Table KB-3. Discharge rate of change limits for S65/S-65A (revised 1/14/19).

Discharge (cfs)	Maximum Rate of Increase (cfs/day)	Maximum Rate of Decrease (cfs/day)
0-300	100	-50
301-650	150	-75
651-1,400	300	-150
1,401-3,000	600	-600
>3,000	1,000	-2,000



Report Date: 4/5/2022; data are through: 4/3/2022



Figure KB-4. Restored Kissimmee river channel mean daily dissolved oxygen concentration (mg/L), S-65A discharge (cfs), temperature (°C) and rainfall (inches). Dissolved oxygen (DO) and temperature are mean daily values averaged for PC62, KRBN, PC33, PD62R, and PD42R with an average of three stations reporting this week. Rainfall values are daily totals for Kissimmee River (Pool BCD) AHED watershed.




Figure KB-5. Phase I area Kissimmee River floodplain water depths (from left to right) one year ago, one month ago and current.

SOUTH FLORIDA WATER MANAGEMENT DISTRICT

Stage and Discharge Guidance for 2021-2022.

Zone	KCH Stage (ft NGVD)	S-65/S-65A Discharge*
A	Above regulation schedule line.	Flood control releases as needed with no limits on the rate of discharge change.
B1	In flood control buffer zone (0.5 ft below the schedule line).	Adjust S-65 discharge so that S-65A discharge is between 1400 cfs at the buffer zone line and 3000 cfs at the schedule line.
B2	Between the Flood Control Buffer and the 50.0 ft line.	Adjust S-65 discharge to maintain at least 1400 cfs at S-65A. Use ± 0.2 ft buffer (gray band) above and below the 50.0 ft line to decide when to begin ramping up to 1400 cfs or down to 300 cfs; do not continue reducing discharge if stage rises back to or above the threshold stage line.
B3	Between the 50.0 ft line and 49 ft.	Adjust S-65 discharge to maintain at least 300 cfs at S-65A.
B4	Between 48.5 ft to 49 ft.	Adjust S-65 discharge to maintain S-65A discharge between 0 cfs at 48.5 ft and 300 cfs at 49 ft.
C	Below 48.5 ft.	0 cfs.


*Changes in discharge should not exceed limits in inset table below.

Table KB-3. Discharge Rate of Change Limits for S65/S65A (revised 1/14/19).

Q (cfs)	Maximum rate of INCREASE (cfs/day)	Maximum rate of DECREASE (cfs/day)
0-300	100	-50
301-650	150	-75
651-1400	300	-150
1401-3000	600	-600
>3000	1000	-2000

sfwmd.gov

2021-2022 Discharge Plan for S-65/S-65A



Other Considerations

- When possible, limit lake ascension rate in the Jun 1 - Aug 15 window to 0.25 ft per 7 days in Lakes Kissimmee, Cypress, Hatchineha (S-65), East Toho (S-59) and Toho (S-61).
- If outlook is for extreme dry conditions meet with KB staff to discuss modifications to this plan.

Slide Revised 1/3/2022

10

Figure KB-6. IS-14-50 Discharge Plan for S65/S65A with discharge rate of change limits (revised 1/14/19).

Lake Okeechobee

Lake Okeechobee stage was 13.75 feet NGVD on April 3, 2022, with water levels 0.66 feet lower than a month ago (**Figure LO-1**). Lake stage remains in the Low sub-band (**Figure LO-2**) and has been within the ecological envelope for over 13 weeks (**Figure LO-3**). According to NEXRAD, 0.21 inches of rain fell directly on the Lake last week.

Average daily inflows (excluding rainfall) increased from the previous week, 843 cfs to 881 cfs. Average daily outflows (excluding evapotranspiration) decreased from 4,322 cfs to 3,859 cfs. Highest inflows came from the Kissimmee River through the S-65E structure (861 cfs). The highest outflow (1,948 cfs) was to the west via the S-77 structure, while flows south via the S-350 structures decreased from 1,751 cfs to 1,144 cfs (S-351: 344 cfs; S-352: 345 cfs; and S-354: 455 cfs). Flows east were 556 cfs and 211 cfs via the S-308 structure and the L-8 canal through the S-271 structure (formerly Culvert 10A), respectively. **Figures LO-4 and LO-5** show the combined average daily inflows and outflows for the Lake over the past eight weeks, and average inflows and outflows last week, respectively. These data are provisional and are subject to change.

The most recent snail kite nesting survey recorded 12 more nests on Lake Okeechobee, for a total of 31 nests thus far, most of which were found in the Indian Prairie marsh. Eighteen nests are active with one nest about to fledge, and 13 nests have failed.

The most recent satellite image (March 28, 2022) from the NOAA cyanobacteria monitoring product derived from EUMETSAT's Sentinel 3 OLCI sensor showed scattered areas of low to moderate bloom potential in Fisheating Bay and along the northwestern shoreline (**Figure LO-6**).




Figure LO-1. Lake Okeechobee water depth estimates based on South Florida Water Depth Assessment Tool (SFWDAT).




Figure LO-2. Recent Lake Okeechobee stages with projected stages based on a dynamic position analysis.

Lake Okeechobee Stage vs Ecological Envelope




Figure LO-3. The prior six years of annual stage hydrographs for Lake Okeechobee in comparison to the ecological envelope.




Figure LO-4. Major inflows (orange) to and outflows east and west (blue) from Lake Okeechobee. Outflows south are shown in green. Flows into Lake Okeechobee from the L-8 canal through S-271 (formerly Culvert 10A) or from the C-44 canal through the S-308 are included as inflows. Conversely, flows from Lake Okeechobee into the L-8 or C-44 canals are included with outflows. Inflows are shown as positive values; outflows are negative. Outflows through the S-77 (Caloosahatchee) and S-308 (C-44 Canal) structures are based on downstream gauges to include flows to lock openings for navigation.




Figure LO-5. Inflows into Lake Okeechobee from Indian Prairie basins, Taylor Creek/Nubbin Slough, Kissimmee River and Fisheating Creek, and outflows to the west via S-77, to the east via S-308, to the south via S-351, S-352, S-354, and to southeast via S-271 (formerly Culvert 10A) for the week of March 28, 2022 – April 3, 2022.




Figure LO-6. Cyanobacteria bloom potential on March 28, 2022 based on NOAA's harmful algal bloom monitoring system. Gray color indicates cloud cover.

Estuaries

St. Lucie Estuary

Over the past week, mean total inflow to the St. Lucie Estuary was 755 cfs (**Figures ES-1 and ES-2**), and the previous 30-day mean inflow was 470 cfs. For comparison, the historical provisional mean inflows from the contributing areas are shown in **Figure ES-2**.

Over the past week, salinities increased at all sites within the estuary (**Table ES-1 and Figure ES-3**). The seven-day moving average of the surface and bottom salinities at the US1 Bridge was 21.7. Salinity conditions in the middle estuary were estimated to be within the good range for adult eastern oysters (**Figure ES-4**). Larval oyster recruitment rates reported by the Fish and Wildlife Research Institute were low in February and March, as expected during the cooler months when most oysters are not spawning (**Figure ES-5**).

Caloosahatchee River Estuary

Over the past week, mean total inflow to the Caloosahatchee River Estuary was 1,930 cfs (**Figures ES-6 and ES-7**), and the previous 30-day mean inflow was 1,979 cfs. For comparison, the historical provisional mean inflows from the contributing areas are shown in **Figure ES-7**.

Over the past week, salinities remained the same at S-79, Val I-75, and Shell Point, and decreased slightly at Ft. Myers, Cape Coral, and Sanibel (**Table ES-2 and Figures ES-8 and ES-9**). The seven-day mean surface salinities (**Table ES-2**) were in the good range (0-10) for tape grass at Val I-75 and at Ft. Myers. The seven-day mean surface salinity values were within the good range for adult eastern oysters at Cape Coral and Shell Point, and in the fair range at Sanibel (**Figure ES-10**). Larval oyster recruitment rates reported by the Fish and Wildlife Research Institute were low at Iona Cove and Bird Island in February and March, as expected during the cooler months when most oysters are not spawning (**Figures ES-11 and ES-12**).

Surface salinity at Val I-75 was forecasted for the next two weeks, using an autoregression model (Qiu and Wan, 2013¹) coupled with a linear reservoir model for the tidal basin. Model scenarios included pulse releases at S-79 ranging from 0 to 1500 cfs and steady releases at 2,000 cfs with estimated tidal basin inflows of 47 cfs. Model results from all scenarios predict daily salinity to be 2.1 or lower and the 30-day moving average surface salinity to be 0.5 or lower at Val I-75 at the end of the two-week period (**Table ES-3 and Figure ES-13**). This keeps predicted salinities at Val I-75 within the LORS 2008 salinity range (0.0-5.0).

¹ Qui, C., and Y. Wan. 2013. Time series modeling and prediction of salinity in the Caloosahatchee River Estuary. *Water Resources Research* 49:5804-5816.

Red Tide

The Florida Fish and Wildlife Research Institute reported on April 1, 2022, that *Karenia brevis*, the Florida red tide dinoflagellate, was not observed in any samples collected within the District region. On the east coast, red tide was not observed in samples from Palm Beach County.

Water Management Recommendations

Lake stage is in the Low Sub-Band. Tributary conditions are dry. The LORS2008 release guidance suggests up to 450 cfs release at S-79 to the Caloosahatchee River Estuary and up to 200 cfs release at S-80 to the St. Lucie Estuary.




Figure ES-1. Basins, water control structures and salinity monitoring sites in the St. Lucie Estuary.




Figure ES-2. Total daily inflows from Lake Okeechobee and runoff from the C-44, C-23, C-24, Ten Mile Creek, and Tidal Basins into the St. Lucie Estuary.

Table ES-1. Seven-day mean salinity at oyster monitoring sites in the St. Lucie Estuary. Current means are in bold font; previous week's means are in parentheses. The envelope reflects the preferred salinity range for adult eastern oysters (*Crassostrea virginica*) in the middle estuary. Data are provisional.

Sampling Site	Surface	Bottom	Envelope
HR1 (North Fork)	16.5 (14.7)	18.6 (17.0)	NA ^a
US1 Bridge	21.3 (19.9)	22.1 (21.0)	10.0 – 26.0
A1A Bridge	28.6 (28.3)	32.0 (30.7)	NA ^a

a. The envelope is not applicable.




Figure ES-3. Mean daily salinity at the A1A, US1 and HR1 sites in the St. Lucie Estuary.




Figure ES-4. Seven-day moving average of the surface and bottom salinities at the US1 Bridge in the St. Lucie Estuary.




Figure ES-5. Mean oyster recruitment at the Rio oyster monitoring station and 14-day mean salinity at US1 Bridge.




Figure ES-6. Basins, water control structures and salinity monitoring sites in the Caloosahatchee River Estuary.




Figure ES-7. Total daily inflows from Lake Okeechobee, and runoff from the C-43 and Tidal basins into the Caloosahatchee River Estuary.

Table ES-2. Seven-day mean salinity at six monitoring sites in the Caloosahatchee River Estuary. Current means are in bold font; previous week's means are in parentheses. The envelope at I-75 is for the protection of tape grass in the upper estuary and the envelope in the lower estuary is the preferred salinity range for adult eastern oysters (*Crassostrea virginica*). Data are provisional.

Sampling Site	Surface	Bottom	Envelope
S-79 (Franklin Lock)	0.2 (0.2)	0.2 (0.2)	NA ^a
Val I-75	0.2 (0.2)	0.2 (0.2)	0.0 – 5.0 ^b
Fort Myers Yacht Basin	2.8 (2.7)	3.5 (4.8)	NA ^a
Cape Coral	11.6 (11.9)	12.6 (13.6)	10.0 – 30.0
Shell Point	26.2 (26.1)	27.0 (27.1)	10.0 – 30.0
Sanibel	31.2 (31.4)	32.1 (32.3)	10.0 – 30.0

a. The envelope is not applicable.

b. The envelope is based on the predicted 30-day mean for the next two weeks.




Figure ES-8. Mean daily salinity at upper Caloosahatchee River Estuary monitoring sites and mean daily flow at S-79.




Figure ES-9. Mean daily surface salinity at lower Caloosahatchee River Estuary monitoring sites and mean daily flow at S-79.




Figure ES-10. Seven-day moving average of surface and bottom salinities at Cape Coral, Shell Point and Sanibel monitoring sites in the Caloosahatchee River Estuary.




Figure ES-11. Mean oyster recruitment at the Iona Cove oyster monitoring station and 14-day mean salinity at Cape Coral.




Figure ES-12. Mean oyster recruitment at the Bird Island oyster monitoring station and 14-day mean salinity at Shell Point.

Table ES-3. Predicted salinity at Val I-75 in the Caloosahatchee River Estuary at the end of the forecast period for various S-79 flow release scenarios.

Scenario	Simulated S-79 Flow (cfs)	Tidal Basin Runoff (cfs)	Daily Salinity	30-Day Mean Salinity
A	0	47	2.1	0.5
B	450	47	1.1	0.4
C	800	47	0.6	0.3
D	1000	47	0.4	0.3
E	1500	47	0.3	0.3
F	2000	47	0.3	0.3




Figure ES-13. Forecasted Val I-75 site surface salinity assuming no pulse release at S-79.

Stormwater Treatment Areas

STA-1E: STA-1E Western Flow-way is offline for post-construction vegetation grow in, and the Eastern Flow-way is offline for vegetation management activities related to Tropical Storm Eta. Operational restrictions are in place in STA-1E Central Flow-way for vegetation management activities. Online treatment cells are at or near target stage, and vegetation in the flow-ways is stressed and highly stressed. The 365-day phosphorus loading rates (PLRs) is high for the Central Flow-way (**Figure S-1**).

STA-1W: Operational restrictions are in place in STA-1W Western, Eastern, and Northern Flow-ways due to construction activities. Most treatment cells are at or near target stage. Vegetation in the flow-ways is stressed and highly stressed. The 365-day PLRs for the Northern and Western, and Eastern Flow-ways are $1.0 \text{ g/m}^2/\text{year}$ or lower (**Figure S-2**).


STA-2: STA-2 Flow-way 2 is offline for construction activities. Operational restrictions are in place in STA-2 Flow-ways 3 and 4 for vegetation management activities, and in Flow-way 1 following dryout conditions. Online treatment cells are near or below target stage. Vegetation in Flow-ways 1 and 3 is stressed, and in Flow-ways 2, 4 and 5 is highly stressed. The 365-day PLRs for Flow-ways 1, 4 and 5 are below $1.0 \text{ g/m}^2/\text{year}$. The 365-day PLR for Flow-way 3 is high (**Figure S-3**).



STA-3/4: STA-3/4 Eastern Flow-way is offline for vegetation rehabilitation/drawdown. Most online treatment cells are at or near target stage. Vegetation in the Eastern and Central Flow-ways is highly stressed and in the Western Flow-way is stressed. The 365-day PLRs for the Central and Western Flow-ways are below $1.0 \text{ g/m}^2/\text{year}$ (**Figure S-4**).










STA-5/6: STA-5/6 Flow-way 4 is offline for vegetation management activities. Most treatment cells are below target stage. All treatment cells have highly stressed vegetation conditions except Flow-ways 7 and 8 which are healthy. The 365-day PLRs for flow-ways 1, 6, 7, and 8 are below $1.0 \text{ g/m}^2/\text{year}$. The 365-day PLRs for flow-ways 4 and 5 are high. (**Figure S-5 and S-6**).

For definitions on STA operational language see glossary following figures.

STA-1E Weekly Status Report – 3/28/2022 through 4/3/2022




STA-1E Flow-Way Status			
Flow-Way	Vegetation Status Healthy ----- Stressed ←-----→	365-day P Loading Rate (below 1.0 g P /m ² /yr is optimal)	Online / Offline / Restrictions
Eastern	Offline, Tropical Storm Eta repairs starting 01/24/2022		
Central	←-----→ 		Vegetation Rehab
Western	Offline, post-construction grow in starting 3/28/2022		

As of 4/3/2022	
Stage Based: Relative to Target Stage (TS)	
	Deep Water Level (> 2.8' above TS)
	High Water Level (1.5' – 2.8' above TS)
	0.2' – 1.5' above TS
	Target Stage (TS +/- 0.2')
	Low Water Level (<0.2' below TS)
Depth / Area Based: Percent of Area Dry	
	0-25% Dry
	25-50% Dry
	50-75% Dry
	75-100% Dry

STA-1E Flow & Phosphorus Concentration			
	7-day	28-day	365-day
Total Inflow, ac-ft	0	1,892	177,518
--Lake Inflow, ac-ft	0	N/A	3,300
Total Outflow, ac-ft	12	691	149,413
Inflow Conc., ppb	N/A	75	121
Outflow Conc., ppb	23	32	23
Includes Preliminary Data			

STA-1W Weekly Status Report – 3/28/2022 through 4/3/2022




STA-1W Flow-Way Status			
Flow-Way	Vegetation Status Healthy --- Stressed	365-day P Loading Rate (below 1.0 g P /m ² /yr is optimal)	Online / Offline / Restrictions
Northern			Construction
Western			Construction
Eastern			Construction
Cell 7		N/A	Construction
Cell 8		N/A	Construction

As of 4/3/2022	
Stage Based: Relative to Target Stage (TS)	
	Deep Water Level (> 2.8' above TS)
	High Water Level (1.5' – 2.8' above TS)
	0.2' – 1.5' above TS
	Target Stage (TS +/- 0.2')
	Low Water Level (<0.2' below TS)
Depth / Area Based: Percent of Area Dry	
	0-25% Dry
	25-50% Dry
	50-75% Dry
	75-100% Dry

STA-1W Flow & Phosphorus Concentration			
	7-day	28-day	365-day
Total Inflow, ac-ft	0	3,165	59,304
--Lake Inflow, ac-ft	0	N/A	5,400
Total Outflow, ac-ft	0	0	52,974
Inflow Conc., ppb	N/A	73	160
Outflow Conc., ppb	N/A	N/A	24
Includes Preliminary Data			

Figure S-2. STA-1W Weekly Status Report

STA-2 Weekly Status Report – 3/28/2022 through 4/3/2022




STA-2 Flow-Way Status			
Flow-Way	Vegetation Status Healthy ----- Stressed	365-day P Loading Rate (below 1.0 g P /m ² /yr is optimal)	Online / Offline / Restrictions
1			Post Dryout
2	Offline, construction activities as of 9/7/2021		
3			Vegetation Rehab
4			Vegetation Rehab
5			Online
















As of 4/3/2022	
Stage Based: Relative to Target Stage (TS)	
	Deep Water Level (> 2.8' above TS)
	High Water Level (1.5' – 2.8' above TS)
	0.2' – 1.5' above TS
	Target Stage (TS +/- 0.2')
	Low Water Level (<0.2' below TS)
Depth / Area Based: Percent of Area Dry	
	0-25% Dry
	25-50% Dry
	50-75% Dry
	75-100% Dry

STA-2 Flow & Phosphorus Concentration			
	7-day	28-day	365-day
Total Inflow, ac-ft	3,089	14,981	311,153
--Lake Inflow, ac-ft	100	N/A	51,700
Total Outflow, ac-ft	1,411	4,853	319,706
Inflow Conc., ppb	54	62	89
Outflow Conc., ppb	29	20	15
Includes Preliminary Data			


Figure S-3. STA-2 Weekly Status Report

STA-3/4 Weekly Status Report – 3/28/2022 through 4/3/2022



STA-3/4 Flow-Way Status				STA-3/4 Flow & Phosphorus Concentration			
Flow-Way	Vegetation Status <small>Healthy ----- Stressed</small> 	365-day P Loading Rate <small>(below 1.0 g P / m²/yr is optimal)</small>	Online / Offline / Restrictions	As of 4/3/2022			
				Stage Based: Relative to Target Stage (TS)			
Eastern				 Deep Water Level (> 2.8' above TS)			
				 High Water Level (1.5' – 2.8' above TS)			
Central			Online	 0.2' – 1.5' above TS			
				 Target Stage (TS +/- 0.2')			
Western			Online	 Low Water Level (<0.2' below TS)			
				Depth / Area Based: Percent of Area Dry			
				 0-25% Dry	 50-75% Dry		
				 25-50% Dry	 75-100% Dry		
Total Inflow, ac-ft				83	83	358,620	
--Lake Inflow, ac-ft				0	N/A	38,100	
Total Outflow, ac-ft				0	0	316,281	
Inflow Conc., ppb				31	31	65	
Outflow Conc., ppb				N/A	N/A	15	
Includes Preliminary Data							

STA-5/6 Weekly Status Report – 3/28/2022 through 4/3/2022




STA-5/6 Flow-Way Status			
Flow-Way	Vegetation Status Healthy ----- Stressed	365-day P Loading Rate (below 1.0 g P /m ² /yr is optimal)	Online / Offline / Restrictions
1			Online
2		N/A	Online
3		N/A	Online
4	Offline, vegetation management starting 01/24/2022		
5			Online

As of 4/3/2022	
Stage Based: Relative to Target Stage (TS)	
	Deep Water Level (> 2.8' above TS)
	High Water Level (1.5' – 2.8' above TS)
	0.2' – 1.5' above TS
	Target Stage (TS +/- 0.2')
	Low Water Level (<0.2' below TS)
Depth / Area Based: Percent of Area Dry	
	0-25% Dry
	25-50% Dry
	50-75% Dry
	75-100% Dry

STA-5/6 Flow & Phosphorus Concentration			
	7-day	28-day	365-day
Total Inflow, ac-ft	0	0	177,793
--Lake Inflow, ac-ft	0	N/A	8,800
Total Outflow, ac-ft	0	0	168,884
Inflow Conc., ppb	N/A	N/A	243
Outflow Conc., ppb	N/A	N/A	50
Includes Preliminary Data			

Figure S-5. STA-5/6 Weekly Status Report (Flow-ways 1 – 5)

STA-5/6 Weekly Status Report – 3/28/2022 through 4/3/2022



STA-5/6 Flow-Way Status				As of 4/3/2022	
Flow-Way	Vegetation Status Healthy ----- Stressed ←-----→	365-day P Loading Rate (below 1.0 g P /m ² /yr is optimal)	Online / Offline / Restrictions	Stage Based: Relative to Target Stage (TS)	
				<div>Deep Water Level (> 2.8' above TS)</div> <div>High Water Level (1.5' – 2.8' above TS)</div> <div>0.2' – 1.5' above TS</div> <div>Target Stage (TS +/- 0.2')</div> <div>Low Water Level (<0.2' below TS)</div>	
6	←-----→	1.0	Online		
7	←-----→	1.0	Online		
8	←-----→	1.0	Online		
				Depth / Area Based: Percent of Area Dry	
				<div>0-25% Dry</div>	<div>50-75% Dry</div>
				<div>25-50% Dry</div>	<div>75-100% Dry</div>

Figure S-6. STA-5/6 Weekly Status Report (Flow-ways 6 – 8)

Basic Concepts and Definitions for STA Weekly Status Report

- **Inflow:** Sum of flow volume at all inflow structures to an STA.
- **Lake Inflow:** Portion of the STA total inflow volume that originates from Lake Okeechobee.
- **Outflow:** Sum of flow volume at outflow structures from an STA.
- **Total Phosphorus (TP):** Total mass of phosphorus in all its forms; including particulate, dissolved, etc.
- **Inflow Concentration:** TP concentration is the mass of TP in micrograms per liter of water, $\mu\text{g/L}$ or ppb. Inflow concentration refers to the flow-weighted mean TP from all inflow structures over a period of time.
- **Outflow Concentration:** The flow-weighted mean TP from all outflow structures over a period of time. The outflow concentration represents the reduction of inflow TP achieved by STA treatment of the inflow water.
- **WQBEL:** The STA outflow concentration that is required upon completion of the Restoration Strategies projects by December 2025. The outflow concentration shall not exceed 13 ppb as an annual flow weighted mean in more than 3 out of 5 water years on a rolling basis and shall not exceed 19 ppb as an annual flow weighted in any water year.
- **Flow-Way (FW):** One or more treatment cells connected in series. Cells typically have emergent aquatic vegetation (EAV) in the front portion of the flow-way followed by a mix of EAV and submerged aquatic vegetation (SAV)
- **Vegetation Status:** Healthy means the vegetation condition is good and will allow the STA to perform as designed. Stressed means the vegetation is showing signs of poor health, such as browning or areas of vegetation die-off, or the cell contains undesirable vegetation such as floating exotic vegetation requiring treatment. The TP reduction capability of the STA is affected when the vegetation condition is poor.
- **Phosphorus Loading Rate (PLR):** Mass of inflow TP in grams, divided by total treatment area of STA in square meters, per year. In general, a 365-day value of less than 1.0 is needed for an STA to perform optimally. A PLR of 2.0 is considered very high and a PLR of 3.0 is considered extremely high. The TP reduction capability of the STA is affected when the PLR is high, very high and extremely high.
- **Online:** Online status means the FW can receive and treat inflow.
- **Online with Restriction:** The FW can receive and treat inflow, but the amount of flow or water level may be limited temporarily. For example, a vegetation rehabilitation effort may require reduced flows through an area while the new plants are establishing, or nesting by protected species may require a certain water level not to be exceeded.
- **Offline:** The FW is unable to receive and treat inflow due to repairs, construction, or other prohibitive reasons.
- **Depth:** Difference between the average surface water level in a cell and the average ground elevation in that cell. Target depths, or depths between flow events, are between 1.25 ft to 1.5 ft. As depth approaches or drops below zero, an increasing percentage of the cell is considered dry and STA conditions deteriorate. An increase in depth above target depth is expected with increasing flow. However, as depth increases much above the target depth and is sustained over a period of time, it can be detrimental to vegetation health and overall STA treatment performance.
- **Note:** The data provided in this summary report were developed using a combination of provisional and quality-assured flow and water quality data. In some cases, best professional judgment was used to estimate missing data and revise questionable data. Values provided are not considered final but are appropriate for use in STA operational decision-making.

Everglades

Water Conservation Area Regulation Schedules

WCA-1: The 1-8C gauge measurements fell faster than the slope of the regulation line last week, then flattened at the end of the week. The average on Sunday was 0.14 feet above the falling Zone A1 regulation line. WCA-2A: Stage continues to drop quickly at the S11B headwater last week. The average at that gauge on Sunday was 0.44 feet higher than the flat regulation line. WCA-3A: Last week, the Three Gauge Average stages declined faster than the slope of the Zone A line; average stage was 0.97 feet below the falling regulation line on Sunday. WCA-3A: Stage continued to recede at gauge 62 (Northwest corner) reversing late in the week. The average on Sunday was 1.12 feet below the flat Upper schedule line. (**Figures EV-1** through **EV-4**).

Water Depths

The SFWDAT tool indicates that water depths are below the soil surface across most of Big Cypress National Preserve (BCNP) and WCA-3A North as is typical but not favorable for this time of year. Central WCA-2A and northern WCA-1 are drying down providing wading bird foraging habitat. North to South hydrologic connectivity continues to diminish but remains within Everglades National Park's (ENP) Shark River Slough. (**Figure EV-5**). Comparing current WDAT water depths to the depth one month ago, stages decreased across the Everglades Protection Area, significantly so in western BCNP. Looking back one year, WCA-3A is significantly lower in depth compared to one year ago. WCA-2A, WCA-1, central BCNP and eastern ENP are slightly wetter than a year ago. (**Figure EV-6**). Comparing current depths to the past 20 years, portions of the eastern half WCA-3A fell into the 10th percentile. BCNP remains above the median but below the soil surface. Though flows into NE WCA-3A have diminished downstream hydration remains evident. (**Figure EV-7**).

Taylor Slough and Florida Bay

A spatial average of 0.46 inches of rain fell over Taylor Slough and Florida Bay on Sunday, 4/3/22, but it was only over the eastern areas. Stages in Taylor Slough decreased an average of 0.12 feet over this past week, with the largest weekly change of -0.29 feet in the northern Taylor Slough area (**Figure EV-8**). The Slough is still 9 inches higher than average, driven in large part by the northern areas, which are 15 inches higher than the historical average for this time of year (pre-Florida Bay initiative, which started in 2017). If these recession rates continue, the Slough would be mostly dry after 2 weeks (**Figure EV-9**). Historically, the Slough would be dry by early March, so the pulses of rain and continued water deliveries have prolonged the hydroperiod in this area despite the drier dry season.

Salinities in Florida Bay averaged an increase of 1.8 over the week ending 4/4/22, with individual station changes ranging from -0.2 to +4.9 (**Figure EV-8**). Most of the Bay stations increased with Garfield Bight (GB) going over 40 for the first time in 9 months. Creek flow from the 5 main creeks was negative for most of the week with saline water pushing upstream and raising salinity in the mangrove ecotone (TR on the map, **Figure EV-8**). The interquartile ranges of historical data increased again in April and keeps the current averages below the 75th percentile (**Figure EV-10**), but the western Florida Bay zone is already in the upper half of April's range.

Water Management Recommendations

Conserving water in the northern basins, then allowing that water to move downstream as we transition further into the dry season maximizes the ecological benefit of freshwater on the landscape. This recommendation is currently being epitomized with STA 2 discharges hydrating northern WCA-2A, being picked up by S7, which then supplies NE and NW WCA-3A North. Continued flows via this route are recommended as long as feasible.

Continued inflows via the S-150 have a positive result on stages in northeastern WCA-3A as seen from the air and in recent modeling results. If conditions allow, continued flow through S-150 is recommended due to it potentially having great ecological benefit by slowing recessions near the Alley North colony and the 7,000 White Ibis nesting there. If conditions allow operational discharges into both the western and the eastern WCA-3A water control structures, this operation has greater benefit than discharges to the west alone. Continued rain and freshwater flows to the Taylor Slough area have prolonged the hydroperiod there. If stage can be maintained, it will help expedite deliveries to the south when the wet season begins. Individual regional recommendations can be found in **Table EV-2**.

Table EV-2. Previous week's rainfall and water depth changes in Everglades basins.

Everglades Region	Rainfall (inches)	Stage change (feet)
WCA-1	0.62	-0.08
WCA-2A	0.50	-0.08
WCA-2B	0.48	-0.13
WCA-3A	0.23	-0.10
WCA-3B	0.18	-0.11
ENP	0.15	-0.11




Figure EV-1. WCA-1 stage hydrographs and regulation schedule.




Figure EV-2. WCA-2A stage hydrographs and regulation schedule.




Figure EV-3. WCA-3A stage hydrographs (three-gauge average, S-333 headwater) and regulation schedule.




Figure EV-4. WCA-3A stage hydrograph (Deer gauge; Site 62) and CA62 regulation schedule.




Figure EV-5. Everglades water depths from two months ago (left), one month ago (center) and present (right), based on SFWDAT.




Figure EV-6. Present Everglades water depths (left) and water depth changes from one month (center) and one year (right) ago, based on SFWDAT.




Figure EV-7. Present water depths (4/3/2022) compared to the day of year median over the previous 20 years.




Figure EV-8. Taylor Slough water depths with changes since a week ago and Florida Bay salinities with changes since a week ago.




Figure EV-9. Taylor Slough water depth time series.




Figure EV-10. Eastern (top panel), Central (middle panel) and Western (bottom panel) Florida Bay daily average salinities with interquartile (25-75 percentile) ranges.

Table EV-2. Weekly water depth changes and water management recommendations

SFWMDC Everglades Ecological Recommendations, February 22, 2022 (red is new)			
Area	Weekly change	Recommendation	Reasons
WCA-1	Stage decreased by 0.08'	Allow water to move south from this basin until stages reach the regulation schedule. Maintain the recession rate to less than 0.10 feet per week.	Protect within basin and downstream habitat and wildlife. Stages are above the 90 th percentile.
WCA-2A	Stage decreased by 0.08'	Conserve water in this basin letting the water move south when conditions allow, with northern WCA-3A as the priority for receiving discharge. Maintain the recession rate to less than 0.10 feet per week.	Protect within basin and downstream habitat and wildlife. Protect peat soil and future wading bird forage as the dry season progresses.
WCA-2B	Stage decreased by 0.13'	Conserve water in this basin, maintain a minimum input to maintain stage while moving water south when conditions allow.	Protect within basin and downstream habitat and wildlife.
WCA-3A NE	Stage decreased by 0.13'	Conserve water in this basin, while letting the water move south when conditions allow. Returning to a recession rate less than 0.10 feet per week has an ecological benefit.	Protect within basin peat soils, and downstream habitat and wildlife. Inflows via the S-150 are anecdotally observed to be positively impacting stages in NE WCA-3A North. Lower fire risk.
WCA-3A NW	Stage decreased by 0.07'	Conserve water in this basin letting the water move south when conditions allow. Maintaining a recession rate less than 0.10 feet per week has an ecological benefit.	
Central WCA-3A S	Stage decreased by 0.07'	Returning to a recession rate less than 0.10 feet per week has an ecological benefit.	Protect within basin and downstream habitat and wildlife. Lower fire risk.
Southern WCA-3A S	Stage decreased by 0.12'		
WCA-3B	Stage decreased by 0.11'	Returning to a recession rate of around 0.10 feet per week in this basin, letting the water move south when conditions allow.	Protect within basin and downstream habitat and wildlife. Lower fire risk.
ENP-SRS	Stage decreased by 0.11'	Make discharges to ENP according to COP and TTFF protocol while adaptively considering upstream and downstream ecological conditions.	Protect within basin and upstream habitat and wildlife.
Taylor Slough	Stage changes ranged from -0.29' to +0.03'	Move water southward as possible.	When available, provide freshwater buffer for downstream conditions.
FB- Salinity	Salinity changes ranged -0.2 to +4.9	Move water southward as possible.	When available, provide freshwater to maintain low salinity buffer and promote water movement.

Biscayne Bay

As shown in **Figure BB-1**, mean total inflow to Biscayne Bay was 265 cfs and the previous 30-day mean inflow was 267 cfs. The seven-day mean salinity was 27.1 at BBCW8 and 29.8 at BBCW10, both below the preferred maximum salinity of 35 for these sites. Data are provided by Biscayne National Park.




Figure BB-1. Seven-day mean salinity at BBCW8 and BBCW10 and total daily flow in Biscayne Bay. Total daily flow was calculated using flow from structures S20G, S20F, S21, S21A, S123, and S700P.