

Disclaimer: Information contained in the report addresses environmental conditions only and is not the official South Florida Water Management District operations recommendation or decision.

## **M E M O R A N D U M**

**TO:** John Mitnik, Assistant Executive Director, Executive Office Staff

**FROM:** SFWMD Staff Environmental Advisory Team

**DATE:** September 13, 2023

**SUBJECT:** Weekly Environmental Conditions for Systems Operations

### **Summary**

#### **Weather Conditions and Forecast**

The combination of slightly below-normal moisture levels and a lack of upper-level features will preclude typical wet season rains through Thursday. Under these conditions, widely scattered showers and thunderstorms are expected over the interior, resulting in daily rainfall totals nearly half of what is climatologically expected. Most of the rainfall will be limited to the coastal areas. On Thursday, an upper-level trough will push a frontal boundary southward into north Florida. Moisture pooling south of the frontal boundary will fuel scattered showers and thunderstorms, primarily over the northern interior in the afternoon. The frontal boundary will continue southward into central Florida on Friday and stall through the weekend, pooling more moisture. There is high confidence for widespread heavy rainfall on Friday, possibly in and around Lake Okeechobee. On Saturday, the frontal boundary will press southward, restricting the afternoon thunderstorm activity to areas south of the Lake. An upper-level disturbance passing through the northeastern US could cause the frontal boundary to lift northward on Monday, returning heavy rains to areas north of Lake Okeechobee. For the 7-day period ending next Tuesday morning, near to above average total SFWMD rainfall is likely.

#### **Kissimmee**

Releases were made from East Lake Toho and Lake Toho as needed to keep lake stage from exceeding their respective regulation schedules. Weekly average discharge on September 10, 2023, was 160 cfs and 310 cfs at S-65 and S-65A, respectively. Mean weekly water depth on the Kissimmee River floodplain decreased by 0.02 ft to 0.14 feet over the week ending September 10, 2023. The weekly average concentration of dissolved oxygen in the Kissimmee River increased from 5.6 mg/L last week to 6.5 mg/L for the week ending September 10, 2023, which is above both the potentially lethal level and the level that is physiologically stressful for largemouth bass and other sensitive species.

## **Lake Okeechobee**

Lake Okeechobee stage was 15.41 feet NGVD on September 10, 2023, which was 0.02 feet higher than the previous week and 0.07 feet higher than a month ago. Average daily inflows (excluding rainfall) decreased from 2,330 cfs the previous week to 2,010 cfs. Average daily outflows (excluding evapotranspiration) increased from 40 cfs the previous week to 680 cfs. The cyanobacteria index level was low to moderate along the northwestern regions of the Lake according to the September 10, 2023, satellite image from NOAA's Harmful Algal Bloom Monitoring System. Routine phytoplankton monitoring on September 5 - 6 detected microcystins toxins at 3 sites with a maximum concentration of just 0.5 µg/L recorded at PALMOUT1. Phytoplankton communities were dominated by *Microcystis aeruginosa* at 14 sites and *Planktolyngbya limnetica* at 3 sites, and the remaining 13 sites had mixed communities.

## **Estuaries**

Total inflow to the St. Lucie Estuary averaged 420 cfs over the past week with no flow coming from Lake Okeechobee. Mean salinities increased at the US1 and A1A bridge sites over the past week. Salinity in the middle estuary was in the optimal range (5-10) for adult eastern oysters.

Total inflow to the Caloosahatchee Estuary averaged 2,740 cfs over the past week with 640 cfs coming from Lake Okeechobee. Mean surface salinities increased at Ft. Myers and decreased at the remaining sites in the estuary over the past week. Salinities were in the optimal range (0-10) for tape grass in the upper estuary. Salinities were in the optimal range for adult eastern oysters at Cape Coral (10-25), and in the upper stressed range at Shell Point and Sanibel (>25).

## **Stormwater Treatment Areas**

For the week ending Sunday, September 10, 2023, 100 ac-ft of Lake Okeechobee water was delivered to the FEBs/STAs. The total amount of Lake releases sent to the FEBs/STAs in WY2024 (since May 1, 2023) is approximately 2,200 ac-feet. The total amount of inflows to the STAs in WY2024 is approximately 687,000 ac-feet. Most STA cells are above target stage. STA-1E Western Flow-way is offline for post-construction vegetation grow in. Operational restrictions are in effect in STA-1W Northern Flow-way, STA-2 Flow-ways 2 and 4, STA-3/4 Eastern Flow-way, and STA-5/6 Flow-way 4 for vegetation management activities. Additional restrictions are in effect in STA-1E Central Flow-way for a Restoration Strategies Science Plan study and in STA-2 Flow-way 3 for canal plug refurbishments. This week, if 2008 LORS recommends Lake releases to the WCAs and conditions allow, releases will be sent to STA-2.

## **Everglades**

Most regions of the WCAs transitioned from a good rate of stage change to a fair rate last week as recessions in the wet season are not generally ecologically desirable. Depths remain below average in WCA-2A after being above for several months and are likely not below average in northwestern WCA-3A. Conditions remain indicative of the potential for better wading bird nesting this year compared to the last two years. Stages decreased on average last week in Taylor Slough but remain above the historical average. Post-

hurricane changes dominated the salinity conditions in Florida Bay, salinity dropped on average across the bay but remains above the historical average.

### **Biscayne Bay**

Total inflow to Biscayne Bay averaged 600 cfs, and the previous 30-day mean inflow averaged 990 cfs. The seven-day mean salinity was 23.3 at BBCW8 and 20.5 at BBCW10, both within the ideal salinity range for estuarine organisms in this region (salinity less than 35). Data provided by Biscayne National Park.

## **Supporting Information**

### **Kissimmee Basin**

#### ***Upper Kissimmee***

On September 10, 2023, mean daily lake stages were 56.5 feet NGVD (0.2 feet below schedule) in East Lake Toho, 53.5 feet NGVD (0.2 feet below schedule) in Lake Toho, and 51.0 feet NGVD (0.2 feet below schedule) in Lakes Kissimmee-Cypress-Hatchineha (KCH) (**Table KB-1, Figures KB-1-3**).

#### ***Lower Kissimmee***

For the week ending September 10, 2023, mean weekly discharge was 160 cfs and 310 cfs at S-65 and S-65A, respectively. Mean weekly discharge from the Kissimmee River was 410 cfs at S-65D and 530 cfs at S-65E (**Table KB-2**). Mean weekly headwater stages were 46.4 feet NGVD at S-65A and 27.9 feet NGVD at S-65D on September 10, 2023. Mean weekly river channel stage increased by 0.1 feet to 33.1 ft NGVD over the week ending on September 10, 2023 (**Figure KB-4**). Mean weekly water depth on the Kissimmee River floodplain decreased by 0.02 ft to 0.14 feet over the week ending September 10, 2023 (**Table KB-2, Figure KB-5**). The weekly average concentration of dissolved oxygen in the Kissimmee River increased from 5.6 mg/L the previous week to 6.5 mg/L for the week ending September 10, 2023 (**Table KB-2, Figure KB-6**).

#### ***Water Management Recommendations***


Follow the IS-14-50 discharge plan for S-65/S-65A; maintain at least minimum flow (250-300 cfs) at S-65A. Maintain current headwater stage at S-65D for the time being.

**Table KB-1.** Average discharge for the preceding seven days, Sunday’s average daily stage and Sunday’s average daily departure from KCOL flood regulation lines or temporary schedules. All data are provisional.


Water Body	Structure	Stage Monitoring Site	Weekly (7-Day) Average Discharge (cfs)	Sunday Lake Stage (feet NGVD) <sup>a</sup>	Schedule Type <sup>b</sup>	Sunday Schedule Stage (feet NGVD)	Sunday Departure from Regulation (feet)	
							9/10/23	9/3/23
Lakes Hart and Mary Jane	S-62	LKMJ	150	59.9	R	60.0	-0.1	-0.1
Lakes Myrtle, Preston and Joel	S-57	S-57	20	60.9	R	61.0	-0.1	-0.1
Alligator Chain	S-60	ALLI	160	63.1	R	63.2	-0.1	0.0
Lake Gentry	S-63	LKGT	210	61.0	R	61.0	0.0	-0.1
East Lake Toho	S-59	TOHOE	400	56.5	R	56.7	-0.2	-0.1
Lake Toho	S-61	TOHOW S-61	760	53.5	R	53.7	-0.2	-0.2
Lakes Kissimmee, Cypress and Hatchineha	S-65	KUB011 LKIS5B	160	51.0	R	51.2	-0.2	-0.4

a. Names of in-lake monitoring sites and structures used to determine lake stage. If more than one site is listed, an average is reported.


b. A: projected recession line; R: USACE regulation schedule; S: temporary recession target line; T: temporary schedule; NA: not applicable or not available.



**Figure KB-1.** East Lake Toho regulation schedule, stage, discharge, and rainfall.



**Figure KB-2.** Lake Toho regulation schedule, stage, discharge, and rainfall.



**Figure KB-3.** Lakes Kissimmee, Cypress and Hatchineha regulation schedule, stage, discharge, and rainfall.

**Table KB-2.** One- and seven-day average discharge and stage at Lower Kissimmee basin structures, river channel dissolved oxygen concentrations and water depths in the Phase I area floodplain. All data are provisional.

Metric	Location	Sunday Daily Average	Weekly Average for Previous Seven Day Periods			
		9/10/23	9/10/23	9/3/23	8/27/23	8/20/23
Discharge	S-65	0	160	160	280	91
Discharge	S-65A <sup>a</sup>	300	310	310	320	310
Headwater Stage (feet NGVD)	S-65A	46.6	46.4	45.9	46.3	46.0
Discharge	S-65D <sup>b</sup>	500	410	410	590	800
Headwater Stage (feet NGVD)	S-65D <sup>c</sup>	27.8	27.9	27.7	27.8	27.8
Discharge (cfs)	S-65E <sup>d</sup>	600	530	570	690	1,100
Discharge (cfs)	S-67	0	0	0	0	0
Dissolved Oxygen (mg/L) <sup>e</sup>	Phase I, II/III river channel	5.4	6.5	5.6	5.3	4.6
River channel mean stage <sup>f</sup>	Phase I river channel	33.1	33.1	33.0	33.2	33.4
Mean depth (feet) <sup>g</sup>	Phase I floodplain	0.15	0.14	0.16	0.18	0.23

a. Combined discharge from main and auxiliary structures.

b. Combined discharge from S-65D, S-65DX1 and S-65DX2.

c. Average stage from S-65D and S-65DX1.

d. Combined discharge from S-65E and S-65EX1.


e. Dissolved oxygen is the average of values from sondes KRBN, PC62, PC33, PD62R and PD42R.

f. Mean of five river channel stations (PC62, KRDR02, KRBN, PC33, PC11) in the Phase I area.

g. One-day spatial average obtained from the South Florida Water Depth Assessment Tool (SFWDAT).




Stage and Discharge Hydrograph (report date: 9/12/2023; data through: 9/10/2023)




\*River Channel Stage is the average for PC62, KRDR02, KRBN, PC33, and PC11.



Figure KB-4. Kissimmee River stage, discharge, and rainfall.



**Figure KB-5.** Phase I area Kissimmee River floodplain water depths (from left to right) one year ago, one month ago, and current.



Report Date: 9/12/2023; data are through: 9/10/2023



**Figure KB-6.** Restored Kissimmee River channel mean daily dissolved oxygen concentration (mg/L), S-65A discharge (cfs), temperature (°C) and rainfall (inches). Dissolved oxygen (DO) and temperature are mean daily values averaged for PC62, KRDR02, KRBN, PC33, PC11, PD62R, and PD42R with an average of five stations reporting this week. Rainfall values are daily totals for Kissimmee River (Pool BCD) AHED watershed.


**Stage and Discharge Guidance for 2021-2023.**

Zone	KCH Stage (ft NGVD)	S-65/S-65A Discharge*
A	Above regulation schedule line.	Flood control releases as needed with no limits on the rate of discharge change.
B1	In flood control buffer zone (0.5 ft below the schedule line).	Adjust S-65 discharge so that S-65A discharge is between 1400 cfs at the buffer zone line and 3000 cfs at the schedule line.
B2	Between the Flood Control Buffer and the 50.0 ft line.	Adjust S-65 discharge to maintain at least 1400 cfs at S-65A. Use ± 0.2 ft buffer (gray band) above and below the 50.0 ft line to decide when to begin ramping up to 1400 cfs or down to 300 cfs; do not continue reducing discharge if stage rises back to or above the threshold stage line.
B3	Between the 50.0 ft line and 49 ft.	Adjust S-65 discharge to maintain at least 300 cfs at S-65A.
B4	Between 48.5 ft to 49 ft.	Adjust S-65 discharge to maintain S-65A discharge between 0 cfs at 48.5 ft and 300 cfs at 49 ft.
C	Below 48.5 ft.	0 cfs.

\*Changes in discharge should not exceed limits in inset table below.

Q (cfs)	Maximum rate of INCREASE (cfs/day)	Maximum rate of DECREASE (cfs/day)
0-300	100	-50
301-650	150	-75
651-1400	300	-150
1401-3000	600	-600
>3000	1000	-2000

## 2021-2023 Discharge Plan for S-65/S-65A



**Other Considerations**

- When possible, limit lake ascension rate in the Jun 1 - Aug 15 window to 0.25 ft per 7 days in Lakes Kissimmee, Cypress, Hatchineha (S-65), East Toho (S-59) and Toho (S-61).
- If outlook is for extreme dry conditions meet with KB staff to discuss modifications to this plan.

Slide Revised 1/3/2022

**Figure KB-7.** IS-14-50 Discharge Plan for S65/S65A with discharge rate of change limits (revised 1/14/19).

## Lake Okeechobee

Lake Okeechobee stage was 15.41 feet NGVD on September 10, 2023, which was 0.02 feet higher than the previous week and 0.07 feet higher than a month ago (**Figure LO-1**). Lake stage remained in the Low sub-band (**Figure LO-2**) and was 0.75 feet above the upper limit of the ecological envelope (**Figure LO-3**). According to NEXRAD, 0.87 inches of rain fell directly on the Lake last week.


Average daily inflows (excluding rainfall) decreased from 2,330 cfs the previous week to 2,010 cfs. The highest structure inflow came from the C-41A Canal via the S-84/84X (610 cfs). Average daily outflows (excluding evapotranspiration) increased from 40 cfs the previous week to 680 cfs. The highest average single structure outflow was recorded at the S-77 structure into the C-43 Canal (660 cfs). **Figures LO-4 and LO-5** show the combined average daily inflows and outflows for the Lake over the past eight weeks and average inflows and outflows last week, respectively. These data are provisional and are subject to change.

The cyanobacteria index level was low to moderate along the northwestern region of the Lake according to the September 10, 2023, satellite image from NOAA's Harmful Algal Bloom Monitoring System (**Figure LO-6**). Routine phytoplankton monitoring on September 5 - 6 detected microcystins toxins at 3 sites with a maximum concentration of just 0.5 µg/L recorded at PALMOUT1. Phytoplankton communities were dominated by *Microcystis aeruginosa* at 14 sites and *Planktolyngbya limnetica* at 3 sites, and the remaining 13 sites had mixed communities (**Figure LO-7**). All data presented in this report are provisional and are subject to change.

# Changes in Water Depth


1 Month Ago:  
08/11/2023

Current:  
09/10/2023



**Figure LO-1.** Lake Okeechobee water depth estimates based on South Florida Water Depth Assessment Tool (SFWDAT).

## Lake Okeechobee Water Level History and Projected Stages



**Figure LO-2.** Recent Lake Okeechobee stages with projected stages based on a dynamic position analysis.

# Lake Okeechobee Stage vs Ecological Envelope






Figure LO-3. The prior seven years of annual stage hydrographs for Lake Okeechobee in comparison to the ecological envelope.







**Figure LO-4.** Major inflows (orange) to and outflows east and west (blue) from Lake Okeechobee. Outflows south are shown in green. Flows into Lake Okeechobee from the L-8 canal through S-271 (formerly Culvert 10A) or from the C-44 canal through the S-308 are included as inflows. Conversely, flows from Lake Okeechobee into the L-8 or C-44 canals are included with outflows. Inflows are shown as positive values; outflows are negative. Outflows through the S-77 (Caloosahatchee) and S-308 (C-44 Canal) structures are based on downstream gauges to include flows to lock openings for navigation.



**Figure LO-5.** Inflows into Lake Okeechobee from Indian Prairie basins, Taylor Creek/Nubbin Slough, Kissimmee River and Fisheating Creek, and outflows to the west via S-77, to the east via S-308, to the south via S-351, S-352, S-354, and to southeast via S-271 (formerly Culvert 10A) for the week of September 4 - 10, 2023.



**Figure LO-6.** Cyanobacteria bloom index level on September 10, 2023, based on NOAA’s harmful algal bloom monitoring system. Gray color indicates cloud cover.



**Figure LO-7.** Total microcystins ( $\mu\text{g/L}$ ) data from September 5 - 6, 2023. Sampling locations are overlaid on the September 6, 2023, image from NOAA's harmful algal bloom monitoring system. Gray color indicates cloud cover.

## Estuaries

### ***St. Lucie Estuary***

Over the past week, mean total inflow to the St. Lucie Estuary was 420 cfs (**Figures ES-1 and ES-2**), and the previous 30-day mean inflow was 970 cfs. For comparison, the historical provisional mean inflows from the contributing areas are shown in **Figure ES-2**.

Over the past week, salinities increased at the US1 and A1A Bridge sites (**Table ES-1 and Figure ES-3**). The seven-day moving average of the surface and bottom salinities at the US1 Bridge was 16.5. Salinity conditions in the middle estuary were estimated to be within the optimal range for adult eastern oysters (**Figure ES-4**). The mean larval oyster recruitment rate reported by the Fish and Wildlife Research Institute (FWRI) was 3.2 spat/shell for August, which is the highest rate recorded yet this year (**Figure ES-5**).

### ***Caloosahatchee River Estuary***

Over the past week, mean total inflow to the Caloosahatchee River Estuary was 2,740 cfs (**Figures ES-6 and ES-7**), and the previous 30-day mean inflow was 2,130 cfs. For comparison, the historical provisional mean inflows from the contributing areas are shown in **Figure ES-7**.

Over the past week, surface salinity increased at Ft. Myers and decreased at the remaining sites in the estuary (**Table ES-2 and Figures ES-8 and ES-9**). The seven-day mean salinities (**Table ES-2**) were in the optimal range (0-10) for tape grass in the upper estuary. The seven-day mean salinity values were within the optimal range for adult eastern oysters at Cape Coral and in the stressed range at Shell Point and Sanibel (**Figure ES-10**). The mean larval oyster recruitment rate reported by the FWRI was 4.8 spat/shell at Iona Cove in August, which was a slight decrease from the previous month. At Bird Island, the reported rate was 22.5 spat/shell for August, which was a substantial increase from July (**Figures ES-11 and ES-12**).

Surface salinity at Val I-75 was forecasted for the next two weeks using an autoregression model (Qiu and Wan, 2013<sup>1</sup>) coupled with a linear reservoir model for the tidal basin. Model scenarios included pulse releases at S-79 ranging from 0 to 1,500 cfs, and a steady release at 2,000 cfs with estimated tidal basin inflows of 418 cfs. Model results from all scenarios predict daily salinity to be 2.2 or lower and the 30-day moving average surface salinity to be 0.5 or lower at Val I-75 at the end of the two-week period (**Table ES-3 and Figure ES-13**). This keeps predicted salinities in the upper estuary within the optimal salinity range (0-10) for tape grass.

---


<sup>1</sup> Qiu, C., and Y. Wan. 2013. Time series modeling and prediction of salinity in the Caloosahatchee River Estuary. *Water Resources Research* 49:5804-5816.

### Red Tide


The FWRI reported on September 8, 2023, that *Karenia brevis*, the Florida red tide dinoflagellate, was not observed in any samples collected within the District region.

### Water Management Recommendations

Lake stage is in the Low Sub-Band. Tributary conditions are near normal. The LORS2008 release guidance suggests up to 3,000 cfs release at S-79 to the Caloosahatchee River Estuary and up to 1,170 cfs release at S-80 to the St. Lucie Estuary.




**Figure ES-1.** Basins, water control structures and salinity monitoring sites in the St. Lucie Estuary.




**Figure ES-2.** Total daily inflows from Lake Okeechobee and runoff from the C-44, C-23, C-24, Ten Mile Creek, and Tidal Basins into the St. Lucie Estuary.

**Table ES-1.** Seven-day mean salinity at oyster monitoring sites in the St. Lucie Estuary. Current means are in bold font; previous week's means are in parentheses. The envelope reflects the optimum salinity range for adult eastern oysters (*Crassostrea virginica*) in the estuary. Data are provisional.

Sampling Site	Surface	Bottom	Optimum Envelope
HR1 (North Fork)	<b>8.4</b> (NR)	<b>14.3</b> (11.1)	10.0 – 25.0
US1 Bridge	<b>14.9</b> (11.9)	<b>18.1</b> (13.7)	10.0 – 25.0
A1A Bridge	<b>23.6</b> (22.3)	<b>28.6</b> (26.6)	10.0 – 25.0




**Figure ES-3.** Mean daily salinity at the A1A, US1 and HR1 sites in the St. Lucie Estuary.




**Figure ES-4.** Seven-day moving average of the surface and bottom salinities at the US1 Bridge in the St. Lucie Estuary.






**Figure ES-5.** Mean oyster recruitment at the Rio oyster monitoring station and 14-day mean salinity at US1 Bridge.




**Figure ES-6.** Basins, water control structures and salinity monitoring sites in the Caloosahatchee River Estuary.




**Figure ES-7.** Total daily inflows from Lake Okeechobee, and runoff from the C-43 and Tidal basins into the Caloosahatchee River Estuary.

**Table ES-2.** Seven-day mean salinity at six monitoring sites in the Caloosahatchee River Estuary. Current means are in bold font; previous week's means are in parentheses. The envelope in the upper estuary sites is for the protection of tape grass and the envelope in the lower estuary is the optimum salinity range for adult eastern oysters (*Crassostrea virginica*). Data are provisional.


Sampling Site	Surface	Bottom	Optimum Envelope
S-79 (Franklin Lock)	<b>0.4</b> (1.1)	<b>0.4</b> (1.2)	0.0 – 10.0
Val I-75	<b>0.6</b> (1.9)	<b>1.1</b> (2.4)	0.0 – 10.0
Fort Myers Yacht Basin	<b>7.2</b> (6.4)	<b>10.2</b> (6.7)	0.0 – 10.0
Cape Coral	<b>11.6</b> (14.3)	<b>15.0</b> (15.0)	10.0 – 25.0
Shell Point	<b>26.2</b> (27.9)	<b>27.7</b> (28.6)	10.0 – 25.0
Sanibel	<b>34.1</b> (34.5)	<b>33.1</b> (33.3)	10.0 – 25.0



**Figure ES-8.** Mean daily salinity at upper Caloosahatchee River Estuary monitoring sites and mean daily flow at S-79.



**Figure ES-9.** Mean daily surface salinity at lower Caloosahatchee River Estuary monitoring sites and mean daily flow at S-79.



**Figure ES-10.** Seven-day moving average of surface and bottom salinities at Cape Coral, Shell Point and Sanibel monitoring sites in the Caloosahatchee River Estuary.




Figure ES-11. Mean oyster recruitment at the Iona Cove oyster monitoring station and 14-day mean salinity at Cape Coral.





Figure ES-12. Mean oyster recruitment at the Bird Island oyster monitoring station and 14-day mean salinity at Shell Point.

**Table ES-3.** Predicted salinity at Val I-75 in the Caloosahatchee River Estuary at the end of the forecast period for various S-79 flow release scenarios.

Scenario	Simulated S-79 Flow (cfs)	Tidal Basin Runoff (cfs)	Daily Salinity	30-Day Mean Salinity
A	0	418	2.2	0.5
B	450	418	1.2	0.4
C	750	418	0.7	0.3
D	1,000	418	0.5	0.3
E	1,500	418	0.3	0.3
F	2,000	418	0.3	0.3



**Figure ES-13.** Forecasted Val I-75 site surface salinity assuming no pulse release at S-79.

## Stormwater Treatment Areas

**STA-1E:** STA-1E Western Flow-way is offline for post-construction vegetation grow in. An operational restriction is in place in STA-1E Central Flow-way for a Restoration Strategies Science Plan Study. Online treatment cells are at or above target stage. Vegetation in the flow-ways is stressed and highly stressed. The 365-day phosphorus loading rate (PLRs) for the Central Flow-way and Eastern Flow-ways are high (**Figure S-1**).

**STA-1W:** An operational restriction is in place in STA-1W Northern Flow-way for vegetation management activities. Treatment cells are above target stage. Vegetation in the flow-ways is stressed and highly stressed. The 365-day PLRs for the Eastern, Western, and Northern Flow-ways are high (**Figure S-1**).

**STA-2:** Operational restrictions are in place in Flow-ways 2 and 4 for vegetation management activities, and in STA-2 Flow-way 3 for canal plug refurbishments. Most online treatment cells are above target stage. Vegetation in Flow-ways 1 and 3 is stressed, and in Flow-ways 4 and 5 is highly stressed. The 365-day PLRs for Flow-ways 4 and 5 are below 1.0 g/m<sup>2</sup>/year. The 365-day PLRs for Flow-way 1 and 3 are high (**Figure S-2**).


**STA-3/4:** An operational restriction is in place in the Eastern Flow-way for post-drawdown vegetation grow-in. Online treatment cells are above target stage. Vegetation in the Central Flow-way is highly stressed and in the Western Flow-way is stressed. The 365-day PLRs for the Central and Western Flow-ways are below 1.0 g/m<sup>2</sup>/year (**Figure S-2**).

**STA-5/6:** An operational restriction is in place in Flow-way 4 for vegetation management (prescribed burn). Most treatment cells are at or above target stage. All treatment cells have highly stressed or stressed vegetation conditions except Flow-ways 7 which is healthy. The 365-day PLRs for Flow-ways 1, 6, 7, and 8 are below 1.0 g/m<sup>2</sup>/year, and the 365-day PLRs for Flow-ways 2, 4, and 5 are high, and the 365-day PLR for Flow-way 3 is very high. (**Figure S-3**).

For definitions on STA operational language see glossary following figures.



## Eastern Flow Path Weekly Status Report – 9/4/2023 through 9/10/2023




STA-1W	Flow-way Status
Western	<ul style="list-style-type: none"> <li>• High 365-day PLR</li> <li>• Highly stressed vegetation conditions</li> </ul>
Eastern	<ul style="list-style-type: none"> <li>• High 365-day PLR</li> <li>• Highly stressed vegetation conditions</li> </ul>
Northern	<ul style="list-style-type: none"> <li>• High 365-day PLR</li> <li>• Highly stressed vegetation conditions</li> <li>• Planting emergent vegetation</li> </ul>
Cell 7	<ul style="list-style-type: none"> <li>• Stressed vegetation conditions</li> </ul>
Cell 8	<ul style="list-style-type: none"> <li>• Construction activities</li> </ul>

STA-1E	Flow-way Status
Western	<ul style="list-style-type: none"> <li>• Offline for post-construction vegetation grow-in</li> </ul>
Central	<ul style="list-style-type: none"> <li>• High 365-day PLR</li> <li>• Highly stressed vegetation conditions</li> <li>• RSSP Study - flow restriction</li> </ul>
Eastern	<ul style="list-style-type: none"> <li>• High 365-day PLR</li> <li>• Stressed vegetation conditions</li> </ul>

**Figure S-1.** Eastern Flow Path Weekly Status Report

## Central Flow Path Weekly Status Report – 9/4/2023 through 9/10/2023




STA-3/4	Flow-way Status
<b>Western</b>	<ul style="list-style-type: none"> <li>• Stressed vegetation conditions</li> <li>• Nuisance vegetation control within inflow canal</li> </ul>
<b>Central</b>	<ul style="list-style-type: none"> <li>• Highly stressed vegetation conditions</li> <li>• Removal of floating tussocks</li> <li>• Nuisance vegetation control within inflow canal</li> </ul>
<b>Eastern</b>	<ul style="list-style-type: none"> <li>• Offline for post-drawdown vegetation establishment</li> <li>• Nuisance vegetation control within inflow canal</li> </ul>

STA-2	Flow-way Status
<b>Flow-way 1</b>	<ul style="list-style-type: none"> <li>• High 365-day PLR</li> <li>• Stressed vegetation conditions</li> </ul>
<b>Flow-way 2</b>	<ul style="list-style-type: none"> <li>• Post-construction vegetation grow-in</li> <li>• Planting emergent vegetation</li> </ul>
<b>Flow-way 3</b>	<ul style="list-style-type: none"> <li>• Refurbishments project - plug construction</li> <li>• Stressed vegetation conditions</li> </ul>
<b>Flow-way 4</b>	<ul style="list-style-type: none"> <li>• Planting emergent vegetation</li> <li>• Nuisance vegetation control</li> <li>• Highly stressed vegetation conditions</li> </ul>
<b>Flow-way 5</b>	<ul style="list-style-type: none"> <li>• Highly stressed vegetation conditions</li> </ul>

**Figure S-2.** Central Flow Path Weekly Status Report

## Western Flow Path Weekly Status Report – 9/4/2023 through 9/10/2023



**Figure S-3.** Western Flow Path Weekly Status Report

## Basic Concepts and Definitions for STA Weekly Status Report

- **Inflow:** Sum of flow volume at all inflow structures to an STA.
- **Lake Inflow:** Portion of the STA total inflow volume that originates from Lake Okeechobee.
- **Outflow:** Sum of flow volume at outflow structures from an STA.
- **Total Phosphorus (TP):** Total mass of phosphorus in all its forms; including particulate, dissolved, etc.
- **Inflow Concentration:** TP concentration is the mass of TP in micrograms per liter of water,  $\mu\text{g/L}$  or ppb. Inflow concentration refers to the flow-weighted mean TP from all inflow structures over a period of time.
- **Outflow Concentration:** The flow-weighted mean TP from all outflow structures over a period of time. The outflow concentration represents the reduction of inflow TP achieved by STA treatment of the inflow water.
- **WQBEL:** The STA outflow concentration that is required upon completion of the Restoration Strategies projects by December 2025. The outflow concentration shall not exceed 13 ppb as an annual flow weighted mean in more than 3 out of 5 water years on a rolling basis and shall not exceed 19 ppb as an annual flow weighted in any water year.
- **Flow-Way (FW):** One or more treatment cells connected in series. Cells typically have emergent aquatic vegetation (EAV) in the front portion of the flow-way followed by a mix of EAV and submerged aquatic vegetation (SAV)
- **Vegetation Status:** Healthy means the vegetation condition is good and will allow the STA to perform as designed. Stressed means the vegetation is showing signs of poor health, such as browning or areas of vegetation die-off, or the cell contains undesirable vegetation such as floating exotic vegetation requiring treatment. The TP reduction capability of the STA is affected when the vegetation condition is poor.
- **Phosphorus Loading Rate (PLR):** Mass of inflow TP in grams, divided by total treatment area of STA in square meters, per year. In general, a 365-day value of less than 1.0 is needed for an STA to perform optimally. A PLR of 2.0 is considered very high and a PLR of 3.0 is considered extremely high. The TP reduction capability of the STA is affected when the PLR is high, very high and extremely high.
- **Online:** Online status means the FW can receive and treat inflow.
- **Online with Restriction:** The FW can receive and treat inflow, but the amount of flow or water level may be limited temporarily. For example, a vegetation rehabilitation effort may require reduced flows through an area while the new plants are establishing, or nesting by protected species may require a certain water level not to be exceeded.
- **Offline:** The FW is unable to receive and treat inflow due to repairs, construction, or other prohibitive reasons.
- **Depth:** Difference between the average surface water level in a cell and the average ground elevation in that cell. Target depths, or depths between flow events, are between 1.25 ft to 1.5 ft. As depth approaches or drops below zero, an increasing percentage of the cell is considered dry and STA conditions deteriorate. An increase in depth above target depth is expected with increasing flow. However, as depth increases much above the target depth and is sustained over a period of time, it can be detrimental to vegetation health and overall STA treatment performance.
- **Note:** The data provided in this summary report were developed using a combination of provisional and quality-assured flow and water quality data. In some cases, best professional judgment was used to estimate missing data and revise questionable data. Values provided are not considered final but are appropriate for use in STA operational decision-making.

## **Everglades**

### ***Water Conservation Area Regulation Schedules***

Last week the Everglades Protection Area (EPA) only received a significant amount of rainfall for one day last week. WCA-1: Last week stage within the Refuge was relatively flat. The 3 Gauge average on Sunday was 0.22 feet below the Zone A1 regulation line. WCA-2A: Stage at the 2-17 continues to trend towards the regulation schedule line. The average on Sunday was 0.05 feet above the regulation line. WCA-3A: The 3-Gauge average receded towards the regulation line last week. The average stage on Sunday was 0.71 feet above that line and 0.79 feet below the EHWL. WCA-3A North: Gauge 62 (NW corner) continued to recede below the schedule line last week, averaging 0.14 feet below the Lower Schedule on Sunday. See figures **EV-1** through **EV-4**.

### ***Water Depths***

The SFWDAT tool illustrates water depths continuing to rise in WCA-1 and WCA-3A, with only southern Big Cypress National Preserve (BCNP) and eastern Everglades National Park (ENP) (rocky glades) with the potential for stages below ground. Drier conditions expand in spatial extent in southwestern BCNP. Ponded conditions have receded in eastern WCA-2A but continue to build in the upper reaches of the L-67s and across southern WCA-3A and within WCA-3B. Hydrologic connectivity is strong within all the major sloughs of ENP.

Comparing current WDAT water depths to one month ago conditions in southern WCA-1 and southern WCA-3A are slightly deeper, with WCA-2A and northern WCA-3A slightly lower in depth. Looking back a year ago, conditions are significantly wetter in eastern WCA-3A and along the northern border of WCA-2A, and generally deeper across the EPA with western BCNP the exception (**Figure EV-5** and **Figure EV-6**).

Comparing current conditions to the 20-year average on September 10th: Well above average depths conditions continue across most of the EPA, especially in WCA-3B and northeastern ENP where conditions have remained in the 90th percentile for most of 2023. Below average depths are welcomed in WCA-2A, but drier than average conditions in western BCNP are not (**Figure EV-7**).

### ***Taylor Slough and Florida Bay***

Total weekly rainfall averaged 0.2 inches in Taylor Slough and Florida Bay over the past week (Monday-Sunday) based on the 16 gauges used for this report. Total weekly rainfall ranged from 0.0 inches at Taylor Slough Bridge (TSB) in the northern slough to 0.5 inches at Long Sound (LS) in the eastern nearshore region. Most stages decreased across Taylor Slough, with an average decrease of 0.13 feet. Stage changes ranged from -0.28 feet at TSB in the northern slough to +0.02 feet at P37 in the southern slough (**Figure EV-8** and **Figure EV-9**). Taylor Slough water levels are above the historical average for this time of year by 3.9 inches compared to before the Florida Bay initiative (starting in 2017).

Average Florida Bay salinity was 28.1, 2.6 lower than the previous week. As increased salinity from hurricane-driven high-water levels during the previous week subsided, salinity decreased at most sites. Salinity changes ranged from a decrease of 13.2 at Joe Bay in the eastern nearshore region to an increase of 2.1 at Duck Key in the eastern region (**Figure EV-8**). Eastern and Central salinities are within the interquartile range (IQR) and Western salinities are at the 75<sup>th</sup> percentile (**Figure EV-10**). Average Florida Bay salinity remains above its historical average for this time of year by 1.4, a decrease of 0.9 relative to last week.

**Water Management Recommendations**

Depths in WCA-3A are well positioned to provide good wading bird foraging and be protective of peat soils this upcoming dry season. Keeping water within the system and flowing south has long term ecological benefits; however, operations that elevate the rate of a region’s change in stage in excess of 0.18 feet per week have detrimental ecological impacts on sensitive regions of the EPA. A distribution of flows into the northwestern WCA-3A that slowly builds water depths in that region is ecologically desirable. As conditions remain above the 90th percentile in NE Shark River Slough, continuing strong positive Taylor Slough creek flows to avoid salinity swings in the nearshore areas is ecologically beneficial. Individual regional recommendations can be found in **Table EV-2**.

**Table EV-2.** Previous week’s rainfall and water depth changes in Everglades basins.

Everglades Region	Rainfall (inches)	Stage change (feet)
WCA-1	0.43	-0.03
WCA-2A	0.28	-0.06
WCA-2B	0.95	-0.14
WCA-3A	0.51	-0.13
WCA-3B	0.24	-0.06
ENP	0.68	+0.01




Figure EV-1. WCA-1 stage hydrographs and regulation schedule.






Figure EV-2. WCA-2A stage hydrographs and regulation schedule.




**Figure EV-3.** WCA-3A stage hydrographs (three-gauge average, 3-69W) and regulation schedule.




**Figure EV-4.** WCA-3A stage hydrograph (Deer gauge; Site 62) and CA62 regulation schedule.







**Figure EV-5.** Everglades water depths from two months ago (left), one month ago (center) and present (right), based on SFWDAT.




**Figure EV-6.** Present Everglades water depths (left) and water depth changes from one month (center) and one year (right) ago, based on SFWDAT.




**Figure EV-7.** Present water depths (9/10/2023) compared to the day of year average over the previous 20 years.



**Figure EV-8.** Taylor Slough water depths with changes since a week ago and Florida Bay salinities with changes since a week ago.



**Figure EV-9.** Taylor Slough water depth time series.




**Figure EV-10.** Eastern (top panel), Central (middle panel) and Western (bottom panel) Florida Bay daily average salinities with interquartile (25-75 percentile) ranges.

**Table EV-2.** Weekly water depth changes and water management recommendations

<b>SFWMD Everglades Ecological Recommendations, September 12, 2023 (red is new)</b>			
	Weekly change	Recommendation	Reasons
<b>WCA-1</b>	Stage decreased by 0.03'	Ascension rate of less than +0.18' per week.	Protect within basin and downstream habitat and wildlife (apple snail reproduction).
<b>WCA-2A</b>	Stage decreased by 0.06'	Ascension rate of less than +0.18' per week.	Protect within basin and downstream habitat and wildlife (apple snail reproduction).
<b>WCA-2B</b>	Stage decreased by 0.14'	Ascension rate of less than +0.18' per week.	Protect within basin and downstream habitat and wildlife (apple snail reproduction).
<b>WCA-3A NE</b>	Stage decreased by 0.30'	Ascension rate of less than +0.18' per week.	Protect within basin and downstream habitat (peat soils) and wildlife (fish/crayfish reproduction).
<b>WCA-3A NW</b>	Stage decreased by 0.07'	Ascension rate of less than +0.18' per week.	
<b>Central WCA-3A S</b>	Stage decreased by 0.05'	Ascension rate of less than +0.18' per week.	Protect within basin and downstream habitat and wildlife (apple snail reproduction).
<b>Southern WCA-3A S</b>	Stage decreased by 0.10'		
<b>WCA-3B</b>	Stage decreased by 0.06'	Minimal ascension rate.	Protect within basin and downstream habitat and wildlife (apple snail reproduction). Allow for flow through.
<b>ENP-SRS</b>	Stage increased by 0.01'	Make discharges to ENP according to COP and TFFF protocol while adaptively considering upstream and downstream ecological conditions.	Protect within basin and upstream habitat and wildlife (apple snail reproduction).
<b>Taylor Slough</b>	Stage changes ranged from -0.28' to +0.02'	Move water southward as possible.	When available, provide freshwater to promote water movement.
<b>FB- Salinity</b>	Salinity changes ranged from -13.2 to +2.1	Move water southward as possible.	When available, provide freshwater to promote water movement.

## Biscayne Bay

As shown in **Figure BB-1**, mean total inflow to Biscayne Bay was 600 cfs, and the previous 30-day mean inflow was 990 cfs. The seven-day mean salinity was 23.3 at BBCW8 and 20.5 at BBCW10, both within the ideal salinity range for estuarine organisms in this region (salinity less than 35). Data were provided by Biscayne National Park.



**Figure BB-1.** Seven-day mean salinity at BBCW8 and BBCW10 and total daily flow in Biscayne Bay. Total daily flow was calculated using flow from structures S20G, S20F, S21, S21A, S123, and S700P.