

Disclaimer: Information contained in the report addresses environmental conditions only and is not the official South Florida Water Management District operations recommendation or decision.

M E M O R A N D U M

TO: John Mitnik, Assistant Executive Director, Executive Office Staff

FROM: SFWMD Staff Environmental Advisory Team

DATE: April 22, 2026

SUBJECT: Weekly Environmental Conditions for Systems Operations

Summary

Weather Conditions and Forecast

Monday, a frontal boundary moved through most of south Florida, bringing in a cooler and drier air mass that will remain in place through Wednesday. By the middle part of this week, a series of upper-air disturbances moving along the subtropical jet stream are forecast to reach the northern Gulf of Mexico. This will allow the air mass behind the front to slowly moderate and begin to moisten, supporting the development of a few light showers along portions of the east coast after midnight Wednesday. By Thursday afternoon, a continued increase in low-level moisture should support isolated showers and thunderstorms across the interior. Meanwhile, the trailing upper-air disturbance is forecast to dig over the eastern Gulf of Mexico, while the leading disturbance becomes stretched out over the western Atlantic. This will cause the frontal boundary south of the state to lift slightly northward, drawing moisture back into parts of south Florida. With this setup, scattered showers and thunderstorms are expected to develop near and along the frontal zone. Some of this thunderstorm activity could reach the southern Florida Keys Thursday night, and the southern Everglades and Miami-Dade County late Thursday into Friday. The coverage and placement of this storm activity will depend heavily on the exact position of the frontal boundary, which remains uncertain. However, with the boundary generally expected to remain south of the interior, most of the heavy showers and thunderstorms will likely remain over the Florida Keys. Southeasterly steering flow may allow scattered thunderstorms to develop along parts of the southwest coast Thursday afternoon. By Friday, the upper-air disturbance is forecast to move into the western Atlantic, pulling the deep moisture away from the region and allowing drier air to move back in. Despite this, limited moisture near the surface combined with warmer afternoon temperatures may still support brief, isolated showers or thunderstorms each afternoon this weekend, mainly across the southern interior. Early next week, a shift back to southwesterly steering flow, driven by another disturbance moving across north Florida, could focus some of the afternoon shower and thunderstorm activity along portions of the upper east coast. For the week ending next Tuesday morning, much below average total SFWMD rainfall is likely.

Kissimmee

In the past week, releases were made as needed from East Lake Toho and Lake Toho to continue snail kite nesting season stage recessions to reach low pool by June 1, 2026. Releases from Lakes Kissimmee-Cypress-Hatchineha followed the Headwaters Revitalization Schedule Increment 1 Temporary Deviation Discharge Plan. Weekly average discharge on April 19, 2026, was 640 cfs at S-65 and 550 cfs at S-65A. Mean weekly water depth on the Kissimmee River floodplain increased from the previous week's value of 0.33 feet to 0.34 feet. The weekly average concentration of dissolved oxygen in the Kissimmee River increased from the previous week's value of 7.4 mg/L to 7.8 mg/L, which is above both the potentially lethal level of 1.0 mg/L and the stressful level of 2.0 mg/L for Florida bass and other species.

Lake Okeechobee

Lake Okeechobee stage was 10.66 feet NAVD88 (11.97 ft NGVD29) on April 19, 2026, which was 0.11 feet lower than the previous week and 0.09 feet lower than a month ago. Average daily inflows (excluding rainfall) decreased from 710 cfs the previous week to 580 cfs. Average daily outflows (excluding evapotranspiration) increased considerably from the previous week, rising from 20 cfs to 1190 cfs. The most recent non-obscured satellite image from the April 18, 2026, NOAA's Harmful Algal Bloom Monitoring System suggests moderate to high cyanobacteria activity in Fisheating Bay and along much of the western side of the lake.

Estuaries

Total inflow to the St. Lucie Estuary averaged 996 cfs over the past week with no flow coming from Lake Okeechobee. Mean salinities decreased at all sites over the past week. Salinity in the middle estuary was in the lower stressed range (5-9) for adult eastern oysters.

Total inflow to the Caloosahatchee River Estuary averaged 526 cfs over the past week with 241 cfs coming from Lake Okeechobee. Over the past week, surface salinities increased at all sites in the estuary. Salinities were in the optimal range (0-10) for tape grass at S-79 and Val I-75, and the damaging range (>15) at Ft. Myers. Salinities were in the optimal range (10-25) for adult eastern oysters at Cape Coral and in the upper stressed range (>25) at Shell Point and Sanibel.

Stormwater Treatment Areas

For the week ending April 19, 2026, 700 ac ft of Lake Okeechobee water was delivered to the FEBs/STAs. The total amount of Lake releases sent to the FEBs/STAs in WY2026 is approximately 98,100 ac-feet. The total amount of inflows to the STAs in WY2026 is approximately 687,000 ac-feet. Most online STA treatment cells are at or above target stage. STA-1E Central Flow-way is offline for construction activities. STA-1W Eastern Flow-way is offline for vegetation management activities. Operational restrictions are in effect in STA-1E Western Flow-way and STA-3/4 Eastern Flow-way for vegetation management activities. This week, if LOSOM recommends Lake releases to the WCAs and conditions allow, releases will be sent to STA-2.

Everglades

Below-average rainfall occurred across most of the Everglades Protection Area (EPA), which corresponded with increasing trends in recession rates averaging -0.12 ft/week. Water depths are beginning to return to near normal or below average relative to the past 20 years in areas that have seen recent upticks following recent large rain events. WCA-3A and WCA-3B have continued to exhibit unwavering below-average depths. These conditions can have ecological consequences, including reduced already-limited prey populations, increased risk of damaging wildfires, enhanced peat oxidation, and potential ridge and slough degradation. Wading bird activity remains very limited with nesting in numbers only occurring in WCA-1, some foraging is occurring along the drying front in WCA-2A and there is very little foraging or nesting occurring in WCA-3A. Wood storks have abandoned nests system wide. Unless there is an unexpected large increase in nesting in the next few weeks, this nesting season could be one of the lowest nesting efforts observed in the Everglades in the last 30 years. Taylor Slough stages decreased last week but remain above the recent averages for this time of year by 5.7 inches. Average Florida Bay salinity increased slightly last week; however, all three regions are within their respective interquartile range.

Supporting Information

Kissimmee Basin

Upper Kissimmee

On April 19, 2026, mean daily lake stages were 54.8 feet NAVD88 (0.9 feet below schedule) in East Lake Toho, 51.9 feet NAVD88 (0.8 feet below schedule) in Lake Toho, and 48.6 feet NAVD88 (2.1 feet below the Increment 1 Temporary Deviation schedule) in Lakes Kissimmee-Cypress-Hatchineha (KCH) (**Table KB-1, Figures KB-1-3**).

Lower Kissimmee

For the week ending April 19, 2026, mean weekly discharge was 640 cfs at S-65 and 550 cfs at S-65A. Mean weekly discharge from the Kissimmee River was 700 cfs at S-65D and 560 cfs at S-65E (**Table KB-2**). Mean weekly headwater stages were 45.2 feet NAVD88 at S-65A and 30.0 feet NAVD88 at S-65D. Mean weekly river channel stage increased from the previous week's value of 31.7 feet to 32.6 feet NAVD88 (**Figure KB-4**). Mean weekly water depth on the Kissimmee River floodplain increased from the previous week's value of 0.33 feet to 0.34 feet (**Table KB-2, Figure KB-5**). The weekly average concentration of dissolved oxygen in the Kissimmee River increased from 7.4 mg/L the previous week to 7.8 mg/L (**Table KB-2, Figure KB-6**).

Water Management Recommendations

Continue the stage recessions in East Lake Toho and Lake Toho to reach their low pools on June 1, 2026. In KCH, follow the Headwaters Revitalization Schedule (HRS) Increment 1 Temporary Deviation Discharge Plan for S-65/S-65A (**Figure KB-7**). With KCH stage in Zone B3, target flows between 300 and 1,400 cfs at S-65A, using the Increment 1 Interpolation Tool to determine discharge relative to stage in KCH; if stage decreases into Zone B4, target flows of 300 cfs.

Table KB-1. Average discharge for the preceding seven days, Sunday’s average daily stage and Sunday’s average daily departure from Kissimmee Chain of Lakes (KCOL) flood regulation lines or temporary schedules. All data are provisional.

Water Body	Structure	Stage Monitoring Site	Weekly (7-Day) Average Discharge (cfs)	Sunday Lake Stage (feet NAVD88) ^a	Schedule Type ^b	Sunday Schedule Stage (feet NAVD88)	Sunday Departure from Regulation (feet)	
							4/19/26	4/12/26
Lakes Hart and Mary Jane	S-62	LKMJ	0	59.0	R	59.2	-0.2	-0.2
Lakes Myrtle, Preston and Joel	S-57	S-57	1	59.3	R	59.4	-0.1	0.0
Alligator Chain	S-60	ALLI	48	62.0	R	62.0	0.0	0.0
Lake Gentry	S-63	LKGT	63	59.6	R	59.5	0.1	-0.1
East Lake Toho	S-59	TOHOE	46	54.8	R	55.7	-0.9	-1.0
Lake Toho	S-61	TOHOW S-61	170	51.9	R	52.7	-0.8	-1.0
Lakes Kissimmee, Cypress and Hatchineha	S-65	KUB011 LKIS5B	640	48.6	T	50.7	-2.1	-2.0

a. Names of in-lake monitoring sites and structures used to determine lake stage. If more than one site is listed, an average is reported.

b. A: projected recession line; R: USACE regulation schedule; S: temporary recession target line; T: temporary schedule; NA: not applicable or not available.




Figure KB-1. East Lake Toho regulation schedule, stage, discharge, and rainfall.




Figure KB-2. Lake Toho regulation schedule, stage, discharge, and rainfall.




Figure KB-3. Lakes Kissimmee, Cypress and Hatchineha regulation schedule, stage, discharge, and rainfall.

Table KB-2. One- and seven-day average discharge and stage at Lower Kissimmee basin structures, river channel dissolved oxygen concentrations and water depths in the Phase I area floodplain. All data are provisional.

Metric	Location	Sunday Daily Average	Weekly Average for Previous Seven Day Periods			
		4/19/26	4/19/26	4/12/26	4/5/26	3/29/26
Discharge	S-65	440	640	590	380	420
Discharge	S-65A ^a	350	550	520	340	360
Headwater Stage (feet NAVD88)	S-65A	45.2	45.2	45.2	45.3	45.4
Discharge	S-65D ^b	610	700	490	420	410
Headwater Stage (feet NAVD88)	S-65D ^c	24.6	30.0	29.1	28.8	28.8
Discharge (cfs)	S-65E ^d	470	560	430	340	320
Discharge (cfs)	S-67	0	0	0	0	0
Dissolved Oxygen (mg/L) ^e	Phase I, II/III river channel	7.6	7.8	7.4	8.1	8.6
River channel mean stage (feet NAVD88) ^f	Phase I river channel	31.9	32.6	31.7	31.4	31.3
Mean depth (feet) ^g	Phase I & II/III floodplain	0.33	0.34	0.33	0.33	0.33

a. Combined discharge from main and auxiliary structures.

b. Combined discharge from S-65D, S-65DX1, and S-65DX2.

c. Average stage from S-65D and S-65DX1.


d. Combined discharge from S-65E and S-65EX1.

e. Dissolved oxygen is the average of values from sondes KRBN, PC62, PC33, PD62R, and PD42R.

f. Mean of five river channel stations (PC62, KRDR02, KRBN, PC33, PC11) in the Phase I area.

g. One-day spatial average obtained from the South Florida Water Depth Assessment Tool (SFWDAT).

Stage and Discharge Hydrograph (report date: 4/21/2026; data through: 4/19/2026)



*River Channel Stage is the average for PC62, KRDR02, KRBN, PC33, and PC11.



Figure KB-4. Kissimmee River stage, discharge, and rainfall.





Figure KB-5. Phase I-II-III area Kissimmee River floodplain water depths (from left to right) one year ago, one month ago, and current.



Report Date: 4/21/2026; data are through: 4/19/2026



Figure KB-6. Kissimmee River channel mean daily dissolved oxygen concentration (mg/L), S-65A discharge (cfs), temperature (°C) and rainfall (inches). Dissolved oxygen (DO) and temperature are mean daily values averaged for PC62, KRDR02, KRBN, PC33, PC11, PD62R, and PD42R with an average of five stations reporting this week. Rainfall values are daily totals for Kissimmee River (Pool BCD) AHED watershed.

HRS Increment 1 Temporary Deviation Discharge Plan for S-65/S-65A

Discharge Guidance for Increment 1 Temporary Deviation Discharge Plan		
ZONE	S-65 RELEASES	S-65A TARGET FLOWS
ZONE A	Releases for Flood Risk Management up to maximum structure capacity as determined by downstream constraints with a firm capacity of 3,000 cfs.	
ZONE B1	1,400 cfs minimum ramp to 3,000 cfs at Zone A boundary	S-65A releases between 1,400 cfs and 3,000 cfs at Zone A boundary based on Table 1
ZONE B2	Releases as needed to target flows at S-65A	Target S-65A flows of 1,400 cfs to meet ecological needs
ZONE B3	Releases as needed to target flows at S-65A	S-65A flows between 300 cfs and 1,400 cfs
ZONE B4	Releases as needed to target flows at S-65A	Target S-65A flows of 300 cfs
ZONE B5	Releases as needed to target flows at S-65A	Target S-65A flows of 150 cfs
ZONE C	0 cfs	Flow as needed to maintain optimum S-65A headwater




Table KB-3. Maximum Rate of Change Limits for S-65A

MAXIMUM Release Rate of Change Limits for S-65A. In general recommended rates of change will be slower than shown in this table.

Q (cfs)	Maximum rate of INCREASE (cfs/day)	Maximum rate of DECREASE (cfs/day)
0-300	50	-50
301-650	75	-75
651-1400	150	-150
1401-3000	300	-600
>3000	1000	-2000

CENTRAL AND SOUTHERN FLORIDA PROJECT
 2024 LAKE KISSIMMEE, HATCHINEHA & CYPRESS
 Temporary Deviation (Increment 1)
 DATED: May 2024
 DEPARTMENT OF THE ARMY, JACKSONVILLE DISTRICT
 CORPS OF ENGINEERS, JACKSONVILLE, FLORIDA

Other Considerations

- When possible, limit lake ascension rate in the Jun 1 - Aug 15 window to 0.25 ft per 7 days in Lakes Kissimmee, Cypress, Hatchineha (S-65), East Toho (S-59) and Toho (S-61).
- If outlook is for extreme dry conditions meet with KB staff to discuss modifications to this plan.

Figure KB-7. Headwaters Revitalization Schedule (HRS) Increment 1 Temporary Deviation Discharge Plan for S-65/S-65A.

Lake Okeechobee

Lake Okeechobee stage was 10.66 feet NAVD88 (11.97 ft NGVD29) on April 19, 2026, which was 0.11 feet lower than the previous week and 0.09 feet lower than a month ago (**Figure LO-1**). Lake stage is in Zone D of the regulation schedule, is 0.75 feet above the water shortage management band (**Figure LO-2**) and 0.40 feet below the ecological envelope (**Figure LO-3**). According to NEXRAD, 0.01 inches of rain fell directly over the lake during the previous week, and 1.26 inches were lost to evapotranspiration.

Average daily inflows (excluding rainfall) decreased from the previous week, dropping from 710 cfs to 580 cfs. The highest average inflow came from the Kissimmee River (560 cfs via S-65E(X1)). Average daily outflows (excluding evapotranspiration) increased considerably from the previous week, rising from 20 cfs to 1190 cfs. The largest release was to the west through the S-77 structure (480 cfs). **Figures LO-4 and LO-5** show the combined average daily inflows and outflows for the lake over the past eight weeks, and the average inflows and outflows last week, respectively.

In the most recent non-obscured satellite image from April 18, 2026, NOAA's Harmful Algal Bloom Monitoring System suggests moderate to high cyanobacteria activity in Fisheating Bay and along much of the western side of the lake. (**Figure LO-6**).

Note: All data presented in this report are provisional and are subject to change.

1 Month Ago:
03/19/2026

Current:
04/19/2026

10.75 ft
NAVD88

10.66 ft
NAVD88




Figure LO-1. Lake Okeechobee water depth estimates based on South Florida Water Depth Assessment Tool (SFWDAT).




Figure LO-2. Recent Lake Okeechobee stages with projected stages based on a conditional position analysis. Note: stages are in NGVD29, approximate NAVD88 values are shown for reference.

Lake Okeechobee Stage vs Ecological Envelope




Figure LO-3. The current and select prior year's annual lake stage hydrographs in comparison to the Lake Okeechobee ecological envelope (light grey).




Figure LO-4. Major inflows (orange) to and outflows east and west (blue) from Lake Okeechobee. Outflows south are shown in green. Flows into Lake Okeechobee from the L-8 canal through S-271 (formerly Culvert 10A) or from the C-44 canal through the S-308 are included as inflows. Conversely, flows from Lake Okeechobee into the L-8 or C-44 canals are included with outflows. Inflows are shown as positive values; outflows are negative. Outflows through the S-77 (Caloosahatchee) and S-308 (C-44 Canal) structures are based on downstream gauges to include flows to lock openings for navigation.




Figure LO-5. Inflows into Lake Okeechobee from Indian Prairie basins, Taylor Creek/Nubbin Slough, Kissimmee River and Fisheating Creek (currently no flow data available for FECR), and outflows to the west via S-77, to the east via S-308, to the south via S-351, S-352, S-354, and to southeast via S-271 (formerly Culvert 10A) for the week of Apr 13 - 19, 2026.




Figure LO-6. Cyanobacteria bloom index level on Lake Okeechobee, based on NOAA's harmful algal bloom monitoring system. Gray color indicates cloud cover. *Provisional NOAA image, subject to change*.

Estuaries

St. Lucie Estuary

Over the past week, mean total inflow to the St. Lucie Estuary was 996 cfs (**Figures ES-1 and ES-2**), and the previous 30-day mean inflow was 980 cfs. For comparison, the historical provisional mean inflows from contributing areas are shown in **Figure ES-2**.

Over the past week, salinities decreased at all sites (**Table ES-1 and Figure ES-3**). The seven-day moving average of the surface and bottom salinities at the US1 Bridge was 9.7. Salinity conditions in the middle estuary were estimated to be within lower stressed range (5-9) for adult eastern oysters (**Figure ES-4**). The mean larval oyster recruitment rate reported by the Fish and Wildlife Research Institute (FWRI) for March was 0 spat/shell at Rio, which is equal to the previous month (**Figure ES-5**).

Caloosahatchee River Estuary

Over the past week, mean total inflow to the Caloosahatchee River Estuary was 526 cfs (**Figures ES-6 and ES-7**), and the previous 30-day mean inflow was 876 cfs. For comparison, the historical provisional mean inflows from the contributing areas are shown in **Figure ES-7**.

Over the past week, surface salinities increased at all sites in the estuary (**Table ES-2 and Figures ES-8 and ES-9**). The seven-day mean water column salinities (**Table ES-2**) were in the optimal range (0-10) for tape grass at S-79 and Val I-75, and the damaging range (>15) at Ft. Myers. The seven-day mean salinity values were within the optimal range (10-25) for adult eastern oysters at Cape Coral and in the upper stressed range (>25) at Shell Point and Sanibel (**Figure ES-10**). The mean larval oyster recruitment rates reported by the FWRI in March were 0.03 spat/shell at Iona Cove and 2.9 spat/shell at Bird Island, which is an increase at both sites from the previous month (**Figures ES-11 and ES-12**).

Surface salinity at Val I-75 was forecast for the next two weeks using an autoregression model (Qiu and Wan, 2013¹) coupled with a linear reservoir model for the tidal basin. Model scenarios included pulse releases at S-79 ranging from 450 to 2,000 cfs, with estimated tidal basin inflows of 62 cfs. Model results from all scenarios predict daily salinity to be 5.9 or lower and the 30-day moving average surface salinity to be 6.1 or lower at Val I-75 at the end of the two-week period (**Table ES-3 and Figure ES-13**). This keeps predicted salinities in the upper estuary within the optimal salinity range (0-10) for tape grass.

¹ Qiu, C., and Y. Wan. 2013. Time series modeling and prediction of salinity in the Caloosahatchee River Estuary. *Water Resources Research* 49:5804-5816.

Red Tide

The FWRI reported on April 17, 2026, that *Karenia brevis*, the Florida red tide dinoflagellate, was not observed at bloom concentrations in any samples collected within the District region.

Water Management Recommendations

Lake stage is in Zone D. Current climatological and hydrological conditions are normal. The LOSOM release guidance suggests up to 2,000 cfs release at S-79 to the Caloosahatchee River Estuary and no releases at S-80 to the St. Lucie Estuary.




Figure ES-1. Basins, water control structures and salinity monitoring sites in the St. Lucie Estuary.




Figure ES-2. Total daily inflows from Lake Okeechobee and runoff from the C-44, C-23, C-24, Ten Mile Creek, and Tidal Basins into the St. Lucie Estuary.

Table ES-1. Seven-day mean salinity at oyster monitoring sites in the St. Lucie Estuary. Current means are in bold font; previous week's means are in parentheses. The envelope reflects the optimum salinity range for adult eastern oysters (*Crassostrea virginica*) in the estuary. Data are provisional.

Sampling Site	Surface	Bottom	Optimum Envelope
HR1 (North Fork)	2.6 (11.0)	10.5 (20.6)	10.0 – 25.0
US1 Bridge	8.7 (17.1)	10.7 (19.1)	10.0 – 25.0
A1A Bridge	18.7 (25.9)	24.0 (28.3)	10.0 – 25.0




Figure ES-3. Mean daily salinity at the A1A, US1, and HR1 sites in the St. Lucie Estuary.




Figure ES-4. Seven-day moving average of the surface and bottom salinities at the US1 Bridge in the St. Lucie Estuary.




Figure ES-5. Mean oyster recruitment at the Rio oyster monitoring station and 14-day mean salinity at US1 Bridge.




Figure ES-6. Basins, water control structures, and salinity monitoring sites in the Caloosahatchee River Estuary.




Figure ES-7. Total daily inflows from Lake Okeechobee, and runoff from the C-43 and Tidal basins into the Caloosahatchee River Estuary.

Table ES-2. Seven-day mean salinity at six monitoring sites in the Caloosahatchee River Estuary. Current means are in bold font; previous week's means are in parentheses. The envelope in the upper estuary sites is for the protection of tape grass and the envelope in the lower estuary is the optimum salinity range for adult eastern oysters (*Crassostrea virginica*). Data are provisional.

Sampling Site	Surface	Bottom	Optimum Envelope
S-79 (Franklin Lock)	6.7 (3.6)	6.7 (3.7)	0.0 – 10.0
Val I-75	6.3 (4.8)	9.1 (6.6)	0.0 – 10.0
Fort Myers Yacht Basin	14.3 (12.8)	16.6 (17.9)	0.0 – 10.0
Cape Coral	20.6 (20.4)	22.5 (21.8)	10.0 – 25.0
Shell Point	32.2 (30.3)	32.5 (31.2)	10.0 – 25.0
Sanibel	35.4 (33.9)	35.6 (34.9)	10.0 – 25.0




Figure ES-8. Mean daily salinity at upper Caloosahatchee River Estuary monitoring sites and mean daily flow at S-79.




Figure ES-9. Mean daily surface salinity at lower Caloosahatchee River Estuary monitoring sites and mean daily flow at S-79.




Figure ES-10. Seven-day moving average of surface and bottom salinities at Cape Coral, Shell Point, and Sanibel monitoring sites in the Caloosahatchee River Estuary.




Figure ES-11. Mean oyster recruitment at the Iona Cove oyster monitoring station and 14-day mean salinity at Cape Coral.




Figure ES-12. Mean oyster recruitment at the Bird Island oyster monitoring station and 14-day mean salinity at Shell Point.

Table ES-3. Predicted salinity at Val I-75 in the Caloosahatchee River Estuary at the end of the forecast period for various S-79 flow release scenarios.

Scenario	Simulated S-79 Flow (cfs)	Tidal Basin Runoff (cfs)	Daily Salinity	30-Day Mean Salinity
A	450	62	5.9	6.1
B	750	62	4.6	5.6
C	1,000	62	3.3	5.2
D	1,500	62	1.7	4.7
E	2,000	62	0.9	4.3




Figure ES-13. Surface salinity forecast at the Val I-75 site assuming a 450 cfs pulse release at S-79.

Stormwater Treatment Areas

STA-1E: STA-1E Central Flow-way is offline for construction activities. An operational restriction is in place in the Western Flow-way for post-construction vegetation grow-in. Online treatment cells are at or slightly above target stage. The 365-day PLR for the Western and Eastern Flow-way is below 1.0 g/m²/year (**Figure S-2**).

STA-1W: STA-1W Eastern Flow-way is offline for vegetation management activities. Treatment cells are at or slightly above target stage. Vegetation in the Western and Eastern Flow-ways is highly stressed. The 365-day PLRs for the Northern and Western Flow-ways are below 1.0 g/m²/year (**Figure S-2**).

STA-2: Treatment cells are at or slightly above target stage. The 365-day PLRs for all Flow-ways are below 1.0 g/m²/year (**Figure S-3**).

STA-3/4: An operational restriction is in place in the Eastern Flow-way for vegetation management activities. Most treatment cells are at or slightly above target stage. Vegetation in the Central Flow-way is highly stressed. The 365-day PLR for the Eastern, Central, and Western Flow-ways are below 1.0 g/m²/year (**Figure S-3**).

STA-5/6: Treatment cells are at or below target stage. All treatment cells have highly stressed vegetation conditions. The 365-day PLRs for all Flow-ways are below 1.0 g/m²/year. (**Figure S-4**).

For definitions on STA operational language see glossary following figures

Everglades Stormwater Treatment Areas - STAs

Estimated Inflow and Outflow Volumes

Apr. 13th, 2026 - Apr. 19th, 2026 *Includes preliminary data*

	Total Inflow (acre-feet)	Total Outflow (acre-feet)
STA-1E	400	1,800
STA-1W	2,300	4,700
STA-2	0	2,000
STA-3/4	8,500	9,700
STA-5/6	50	0

- Total WY2026 inflows to STAs (5/1/2025 to 4/19/2026): ~687,000 ac-ft
- Lake Okeechobee releases to FEBs/STAs
 - 4/13/2026 to 4/19/2026: 700 ac-ft
 - WY 2026: ~ 98,100 ac-ft
- Extensive vegetation management activities underway to address stressed and highly stressed vegetation in EAV cells
- All treatment cells are at or near target water depth




Figure S-1. STA depths and flow volumes

0 CFS Lake release capacity in Eastern Flow Path:
 4/20/2026 to 4/26/2026
 Subject to change weekly as dry season progresses




Figure S-2. Eastern Flow Path Weekly Status Report

400 CFS Lake release capacity in Central Flow Path:
4/20/2026 to 4/26/2026
Subject to change weekly as dry season progresses




Figure S-3. Central Flow Path Weekly Status Report

0 CFS Lake release capacity in Western Flow Path:
4/20/2026 to 4/26/2026




Figure S-4. Western Flow Path Weekly Status Report

Basic Concepts and Definitions for STA Weekly Status Report

- **Inflow:** Sum of flow volume at all inflow structures to an STA.
- **Lake Inflow:** Portion of the STA total inflow volume that originates from Lake Okeechobee.
- **Outflow:** Sum of flow volume at outflow structures from an STA.
- **Total Phosphorus (TP):** Total mass of phosphorus in all its forms; including particulate, dissolved, etc.
- **Inflow Concentration:** TP concentration is the mass of TP in micrograms per liter of water, $\mu\text{g/L}$ or ppb. Inflow concentration refers to the flow-weighted mean TP from all inflow structures over a period of time.
- **Outflow Concentration:** The flow-weighted mean TP from all outflow structures over a period of time. The outflow concentration represents the reduction of inflow TP achieved by STA treatment of the inflow water.
- **WQBEL:** The STA outflow concentration that is required upon completion of the Restoration Strategies projects by December 2025. The outflow concentration shall not exceed 13 ppb as an annual flow weighted mean in more than 3 out of 5 water years on a rolling basis and shall not exceed 19 ppb as an annual flow weighted in any water year.
- **Flow-Way (FW):** One or more treatment cells connected in series. Cells typically have emergent aquatic vegetation (EAV) in the front portion of the flow-way followed by a mix of EAV and submerged aquatic vegetation (SAV)
- **Vegetation Status:** Healthy means the vegetation condition is good and will allow the STA to perform as designed. Stressed means the vegetation is showing signs of poor health, such as browning or areas of vegetation die-off, or the cell contains undesirable vegetation such as floating exotic vegetation requiring treatment. The TP reduction capability of the STA is affected when the vegetation condition is poor.
- **Phosphorus Loading Rate (PLR):** Mass of inflow TP in grams, divided by total treatment area of STA in square meters, per year. In general, a 365-day value of less than 1.0 is needed for an STA to perform optimally. A PLR of 2.0 is considered very high and a PLR of 3.0 is considered extremely high. The TP reduction capability of the STA is affected when the PLR is high, very high and extremely high.
- **Online:** Online status means the FW can receive and treat inflow.
- **Online with Restriction:** The FW can receive and treat inflow, but the amount of flow or water level may be limited temporarily. For example, a vegetation rehabilitation effort may require reduced flows through an area while the new plants are establishing, or nesting by protected species may require a certain water level not to be exceeded.
- **Offline:** The FW is unable to receive and treat inflow due to repairs, construction, or other prohibitive reasons.
- **Depth:** Difference between the average surface water level in a cell and the average ground elevation in that cell. Target depths, or depths between flow events, are between 1.25 ft to 1.5 ft. As depth approaches or drops below zero, an increasing percentage of the cell is considered dry and STA conditions deteriorate. An increase in depth above target depth is expected with increasing flow. However, as depth increases much above the target depth and is sustained over a period of time, it can be detrimental to vegetation health and overall STA treatment performance.
- **Note:** The data provided in this summary report were developed using a combination of provisional and quality-assured flow and water quality data. In some cases, best professional judgment was used to estimate missing data and revise questionable data. Values provided are not considered final but are appropriate for use in STA operational decision-making.

Everglades

Water Conservation Area

Regulation Schedules

WCA-1: Stage at the 1-8C gauge remained above the regulation line last week with very little slope and is 0.32 feet above the A1 zone regulation line as of Sunday, April 19, 2026.

WCA-2A: Last week's depth change at the 2-17 gauge was minimal. Stage was 1.02 feet above the regulation line on Sunday.

WCA-3A: The 3-gauge average remains well within zone B, continuing on a steady downward trend last week. On Sunday, stages were 1.74 ft NGVD29 below the zone A regulation line. Stage at Gauge 62 (NW corner) continued a steep downward trend away from the regulation line and was below the Upper Schedule line by 2.1 feet on Sunday (**Figures EV-1 through EV-4**).

Water Depths

The SFWDAT model output for April 19, 2026, illustrates deeper water depths in both WCA-1 and WCA-2A compared to a month ago due to a reversal two weeks ago. Depths are now back on a trend of receding from north to south. WCA-1 has only a small area in the north approaching ground level. WCA-2A also only has a small portion of the north-central region approaching ground level and still remains slightly more ponded in the southeast. WCA-3A remains very dry, having received the least amount of rainfall in the system over the last month. In the Everglades National Park (ENP), Taylor Slough (TS) and Shark River Slough (SRS) are showing less connection from north to south with TS still showing potential connection to the coast vs SRS with now much more limited connection, if any. Comparing current conditions to depths over the last twenty years, WCA-1 depths are primarily around the 50th percentile and in the 80th percentile in the NE/SE portions of that basin. WCA 2A depths are around the 60th percentile in the central region and 80th percentile in the SE. BCNP depths continue to be above average in the center of the preserve, but even above average depths for this time of year are below ground along Tamiami Trail. Within WCA-3A and 3B, depths remain below the 10th percentile, although a portion of NW WCA3A (below Rotenberger WMA) is now showing depths closer to the 70th percentile. Conditions in northern and central ENP are now below average depths, while the southern coastline of ENP remains at above average depths (**Figures EV-5 through EV-6**).

Taylor Slough and Florida Bay

All stages decreased across Taylor Slough over the past week, with an average decrease of 0.23 feet for the week. Changes ranged from -0.59 feet at E112 in the northern slough to -0.06 feet at EPSW in the C-111 area (**Figure EV-7 and Figure EV-8**). The stage at Taylor Slough Bridge (TSB) again dropped below ground, indicating a lack of water at the head of the slough. Taylor Slough water levels are above the recent average (WY1993-2016) for this time of year by 5.7 inches compared to before the Florida Bay Initiative (starting in 2017), a decrease of 2.3 inches relative to last week. The stages at Craighead

Pond (CP) and TSB are below the estimated historical average by 0.18 and 1.06 feet, respectively.

Average salinity in Florida Bay was 34.3, an increase of 2.8 from last week. Salinity changes ranged from -3.4 at Garfield Bight (GB) in the western nearshore region to +14.3 at Joe Bay (JB) in the eastern nearshore region (**Figure EV-7**). Salinity remains above the estimated historical average and within the WY2001-2016 Interquartile Range (IQR) for all three regions (**Figure EV-9**). Salinity in the western region remains near the hypersalinity threshold. Bay-wide salinity remains just above its recent average (WY1993-2016) for this time of year by 0.5, an increase of 0.3 from last week's comparison.

Salinity at the Taylor River (TR) station in the mangrove zone (tracked for the Florida Bay MFL) was 20.0, an increase of 18.5 from last week (**Figure EV-10**). The 365-day moving sum of flow from the five major creeks (McCormick Creek, Taylor River, Mud Creek, Trout Creek, West Highway Creek) was unable to be assessed due to missing data.

Average rainfall across Taylor Slough and Florida Bay was 0.05 inches over the past week, based on the 18 gauges used for this report. Rainfall ranged from 0.00 inches at 11 stations to 0.42 inches at Garfield Bight (GB) (**Figure EV-11**). Wind directions and speeds in Florida Bay ranged from 1.7 mph E on April 19th to 25.8 mph NE on April 13th (**Figure EV-11**).

The Taylor River, Mud Creek, and West Highway Creek flow stations are currently offline until further notice, so data from all five major creeks are unable to be assessed. Some data was also missing from Trout Creek this week, but overall flows were likely net negative for the week (**Figure EV-13**). Average daily flow from Alligator Creek was -93 acre-feet, with net negative flows for the week (**Figure EV-13**).

Implications/considerations for water management.

- Average stage recessions continue to decline in WCA-3A. Due to current drought conditions, slower recessions in WCA-3A can help to protect the wetland ecology from damaging dry downs as the dry season continues.
 - WCA-3A and WCA-3B continue to experience unseasonably dry conditions.
 - Poor wading bird nesting is expected to continue for the fourth year into the 2026 nesting season.
 - With continued dry season drought conditions, conserving water within the WCAs, especially WCA-3A, will continue to be ecologically beneficial.
- Depths remain above average (~1.2 feet) in southern WCA-2A where shallower conditions on average are needed to recover ridge and slough habitat. However, holding water high in the north of the system, as further dry conditions are predicted, may prove ecologically beneficial as the Everglades dry down continues to accelerate due to increasing evapotranspiration. Input of water into areas prone to wildfire and peat oxidation may help to buffer the worst of the ecological stress.

- Taylor Slough depths decreased last week but remained above the recent average for this time of year. There is now more potential for slough connectivity extending down to Florida Bay and higher creek flows from previous rain are likely to help maintain consistent Florida Bay salinities. Salinities remain slightly above their recent average in Florida Bay.
 - Freshwater input into Taylor Slough and the C-111 basin could help moderate salinities and support recovery of estuarine conditions in Florida Bay.
- Conserving water in the WCAs while providing freshwater input to the sloughs of ENP will require careful consideration of a balance between the upstream and downstream ecological needs of the system. Individual regional recommendations can be found in **Table EV-2**.

Table EV-2. Previous week’s rainfall and water depth changes in Everglades basins.

Everglades Region	Rainfall (inches)	Stage change (feet)
WCA-1	0.33	-0.01
WCA-2A	0.17	-0.04
WCA-2B	0.10	-0.20
WCA-3A	0.26	-0.10
WCA-3B	0.16	-0.24
ENP	0.25	-0.15




Figure EV-1. WCA-1 stage hydrographs and regulation schedule.




Figure EV-2. WCA-2A stage hydrographs and regulation schedule. $11.94 - 1.49 = 10.45, 1.05$




Figure EV-3. WCA-3A stage hydrographs (three-gauge average, 3-69W) and regulation schedule. $8.24 - 1.5 = 6.74, 1.68$




Figure EV-4. WCA-3A stage hydrograph (Deer gauge; Site 62) and regulation schedule.




Figure EV-5. Everglades water depths from two months ago (left), one month ago (center) and present (right), based on SFWDAT.




Figure EV-6. Present water depths (April 19, 2026) compared to the day of year relative to average (percentile) over the previous 20 years.




Figure EV-7. Taylor Slough water depths and Florida Bay salinities with changes since a week ago.




Figure EV-8. Taylor Slough water depth time series for Taylor Slough Bridge (TSB; northern slough) and Craighead Pond (CP; southern slough).




Figure EV-9. Eastern (top panel), Central (middle panel) and Western (bottom panel) Florida Bay daily average salinities with WY2001-2016 interquartile (25-75 percentile) ranges (IQR) and estimated historical daily average salinities. The hypersalinity threshold indicates the level at which salinities start to become harmful to seagrass.




Figure EV-10. Daily average salinity at Taylor River (TR) tracked for the Florida Bay MFL criteria. The 365-day total creek flow MFL metric is not currently available due to missing creek flow data.




Figure EV-11. Daily average rain across Taylor Slough and Florida Bay, along with hourly average wind speed and direction (measured at Long Key) in Florida Bay over the past four weeks.




Figure EV-12. Top: daily average creek flow summed between the five major creeks with estimated historical daily flow over the past four weeks (**note:** data from Taylor River, Mud Creek and West Highway Creek are currently unavailable since November 21st, 2025). Bottom: Daily average Alligator Creek flow data. N/A indicates missing data.

Table EV-2. Weekly water depth changes and water management recommendations

SFWMD Everglades Ecological Recommendations, April 19, 2026 (red is new)			
	Weekly change	Recommendation	Reasons
WCA-1	Stage decreased by 0.01 feet.	A recession of no faster than 0.05 feet per week.	Conserve water, maintain within basin and downstream habitat and wildlife. Maintain maintenance access for vegetation management.
WCA-2A	Stage decreased by 0.04 feet	A recession of no faster than 0.05 feet per week.	Maintain within basin (north versus south) and downstream habitat and wildlife.
WCA-2B	Stage decreased by 0.20 feet	A recession of no faster than 0.12 feet per week.	Protect within basin and downstream habitat and wildlife.
WCA-3A NE	Stage decreased by 0.02 feet	A recession of no faster than 0.05 feet per week.	Conserve water, maintain within basin and downstream habitat and wildlife. Provide suitable depths for aquatic prey and protect against peat soil loss during the dry season.
WCA-3A NW	<u>Stage decreased by 0.22 feet</u>	A recession of no faster than 0.05 feet per week.	
Central WCA-3A S	Stage decreased by 0.05 feet	A recession of no faster than 0.05 feet per week.	Conserve water, maintain within basin and downstream habitat and wildlife. Provide suitable depths for aquatic prey and protect against peat soil loss during the dry season.
Southern WCA-3A S	<u>Stage decreased by 0.22 feet</u>		
WCA-3B	Stage decreased by 0.24 feet	A recession of no faster than 0.12 feet per week.	Protect within basin and downstream habitat and wildlife.
ENP-SRS	Stage decreased by 0.15 feet.	Make discharges to ENP according to COP protocol, considering up/down stream ecological conditions.	Protect within basin and upstream habitat and wildlife.
Taylor Slough	Stage changes ranged from -0.59 feet to -0.06 feet	Move water southward as possible.	When available, provide freshwater to promote water movement.
FB- Salinity	Salinity changes ranged from -11.2 to +14.3	Move water southward as possible.	When available, provide freshwater to promote water movement.