

**Manuscript
EMA # 403**

**Atmospheric Deposition of Phosphorus: Concepts,
Constraints and Published Deposition Rates
for Ecosystem Management**

**(This paper is a major revision of Technical
Publication #360 dated April 1999)**

February, 2002

by

Garth W. Redfield

**Environmental Monitoring & Assessment Department
South Florida Water Management District
3301 Gun Club Road
West Palm Beach, FL 33416**

Atmospheric Deposition of Phosphorus: Concepts, Constraints and Published Deposition Rates for Ecosystem Management

Garth W. Redfield

Environmental Monitoring and Assessment Department,
South Florida Water Management District, 3301 Gun Club Road, West Palm
Beach, FL 33416 gredfiel@sfwmd.gov

This manuscript has been submitted (February 4, 2002) to Freshwater Systems, an electronic web-based journal published by Thescienificworld.com. The paper is a major revision of the District's Technical Publication # 360 of April, 1999, and has been expanded, updated and revised to better reflect the interface between the atmospheric and environmental sciences on atmospheric deposition of phosphorus.

**Atmospheric Deposition of Phosphorus:
Concepts, Constraints and Published Deposition Rates for Ecosystem Management**

Garth W. Redfield*

Environmental Monitoring and Assessment Department,
South Florida Water Management District
3301 Gun Club Road, West Palm Beach, FL 33406
gredfiel@sfwmd.gov

Key Words: atmospheric phosphorus deposition, phosphorus inputs, bulk precipitation, atmospheric particles, conceptual deposition model, dry deposition, nutrient loading, and giant atmospheric particles.

Domains: atmospheric deposition, atmospheric aerosols and particles, ecosystem phosphorus loading, deposition monitoring, bulk precipitation, wet and dry atmospheric deposition, nutrient inputs to ecosystems, phosphorus deposition rates, deposition research planning.

Three figures and five tables in the text.

ABSTRACT

This paper summarizes concepts underlying the atmospheric input of phosphorus (P) to ecosystems, published rates of P deposition, measurement methods, and approaches to future monitoring and research. Phosphorus conveyed through the atmosphere can be a significant nutrient source for many freshwater and marine ecosystems. Particle sources and sinks at the land-air interface produce variation in P deposition from the atmosphere across temporal and spatial scales. Natural plant canopies can affect deposition rates by changing the physical environment and surface area for particle deposition. Land-use patterns can alter P deposition rates by changing particle concentrations in the atmosphere. The vast majority of P in dry atmospheric deposition is conveyed by *coarse* (2.5 to 10 μm) and *giant* (10 to 100 μm)

particles, and yet these size fractions represent a challenge for long-term atmospheric monitoring in the absence of accepted methods for routine sampling. Most information on P deposition is from bulk precipitation and wet/dry sampling, with questionable precision and accuracy. Published annual rates of P deposition using these methods in locations around the globe range from about 5 to well over 100 mg Pm⁻²yr⁻¹, although most inland ecosystems receive between 20 and 80 mg Pm⁻²yr⁻¹. Rates below 30 mg Pm⁻²yr⁻¹ are found in remote areas and near coastlines. Intermediate rates of 30 to 50 mg Pm⁻²yr are associated with forests or mixed land use, and rates of 50 to 100 mg Pm⁻²yr⁻¹ or more are often recorded from urban or agricultural settings. Comparison with other methods suggests that these bulk estimates provide reasonable boundaries around actual P deposition rates for various land uses. However, data screening cannot remove all the positive bias caused by contamination of bucket collectors. As a consequence, continued sampling with these standard collectors in a region will not reduce the large uncertainty in rates derived from existing data. Calibrated surface accumulation methods hold promise as a primary means to estimate P flux in future monitoring. New methods for long-term P deposition monitoring will require an intercomparison of P flux estimates from surrogate surfaces, impactor sampling of particle concentrations combined with deposition models, and “throughfall” estimates for natural canopies. With better sampling methods, P deposition should be included in national atmospheric monitoring networks.

Contents

Introduction

Major Features of the South Florida Environment

Conceptual Basis for P Deposition from the Atmosphere

Components and Size Fractions of P Deposition from the Atmosphere

Major Sources of Variation in P Deposition Rates

P Deposition: Integration from an Ecological Perspective

Published Rates for P Deposition from the Atmosphere

Future Measurements of Atmospheric P Deposition

Wet Deposition of Phosphorus

Dry Deposition of P-Bearing Particles

Beyond Bulk Precipitation and Wet/Dry Deposition Estimates

Conclusions

Acknowledgments

References

INTRODUCTION

The flux, or loading, of nutrients to ecosystems occurs primarily through point source discharges, surface water, groundwater, and atmospheric deposition. The input of nutrients from wastewater discharges has become less of a problem over the last 25 years and can be quantified from discharge monitoring records. Loading by surface water runoff has been measured for many ecosystems and can be the major source of nutrients for aquatic environments, particularly those with large watersheds. Groundwater inputs are more difficult to assess and are often estimated indirectly by mass balance calculations. Nutrients conveyed by rainfall and dry fallout from the atmosphere are a substantial source of nutrient loading to many freshwater environments^{1,2,3,4,5,6}. For the marine environment, Graham and Duce⁷, Jickells⁸ and Guerzoni⁹ have emphasized the overall importance of atmospheric inputs to the oceans, particularly in oligotrophic regions. Conversely, the oceans may contribute substantial amounts of P to continents through aerosols injected into the air from the ocean surface¹⁰.

Compared to other elements conveyed through the atmosphere, P has received relatively little attention. Surprisingly, Gorham's¹ review of the chemistry of inland waters does not include P in the discussion of the chemistry of atmospheric precipitation. Even the multi-agency Great Waters Program¹¹, Environmental Protection Agency (EPA) and National Oceanic and Atmospheric Association (NOAA) dry deposition networks¹², and the National Atmospheric Deposition Program/National Trends Network¹³ do not include total P monitoring from atmospheric sources in spite of the importance of phosphorus to many freshwater systems. The United States Environmental Protection Agency¹⁴, in a thorough review of atmospheric particles, included information on P deposition in just several tables concerning particle composition. In addition, estimates of atmospheric P loading with methods other than bulk and wet/dry bucket collectors are not common because many studies of atmospheric inputs have been focused on aerosols associated with acidic deposition, such as nitrogen and sulfur^{12,15}.

Phosphorus is limiting to algal growth in many freshwater lakes^{16,17}, and P control is often the primary focus of lake management^{18,19}. In large, shallow, freshwater environments, such as the Florida Everglades, atmospheric inputs of P could be very important because a major portion of this expansive marsh ecosystem receives very little water and nutrient input from surface water inflow²⁰. Substantial uncertainty exists in both the precision and accuracy of deposition rates for P in South Florida and elsewhere. Using an extensive 13-station data set from South Florida, Ahn and James²¹ demonstrated high variability in loading estimates for

atmospheric P and concluded that this substantial uncertainty was due to random noise in the data. Irwin and Kirkland²² reported a wide range of values in their review of the historical data on precipitation quality (bulk deposition) in Florida and concluded that these data were insufficient to define baseline conditions for the region. Hendry et al.²³ also reported major variation in P deposition among sampling sites, with values ranging from 17 to 111 mg Pm⁻²yr⁻¹. They found the overall state average deposition rate as bulk precipitation to be 50 mg Pm⁻²yr⁻¹. More recently, Dixon et al.²⁴ reported annual station averages ranging from 52 to 176 mg Pm⁻²yr⁻¹ for bulk deposition in the Tampa Bay area, with a seven-station average of 93 mg Pm⁻²yr⁻¹. Peters and Reese²⁵ also found data variability to be very high (wet values from 0.001 to 0.620 mg P L⁻¹) in dry and bulk bucket-type collectors near Lake Okeechobee in Florida. Such variability is not unique to Florida; published deposition rates for P in other regions^{26,27,28,29} range from less than 10 to well over 150 mg Pm⁻²yr⁻¹. An extensive compilation of these rates is included in this review.

While high variability is common for environmental parameters, it is compounded for P deposition by the lack of information on the accuracy of data used in estimating long-term ecosystem inputs. Standard methods of atmospheric monitoring, particularly those for particle fallout, often measure only a portion of total deposition. Some approaches provide defensible estimates for the deposition of fine aerosols (micro-meteorological methods), but do not quantify the fallout of larger particles³⁰. The common approach to long-term flux measurement using funnels or bucket collectors provides the opposite result. Bucket devices provide some data on the flux of larger particles, but are subject to great uncertainty due to their poor aerodynamic qualities and susceptibility to environmental contamination. Methods for measuring compliance with air quality standards assess ambient concentrations of fine particles that affect human health, but are not suitable for deriving total fallout to the environment³¹. Those studies that have provided rigorous estimates of particle flux used several complementary collection devices simultaneously, but were limited to short-term estimates in only a few environments^{32,33}. The limitations of methods and constraints on study design, as well as other factors to be discussed later in this paper, require the ecosystem scientist or manager to interpret disparate published rates of atmospheric material flux from a large body of literature, with substantial variability and minimal assurance of accuracy.

There are many natural sources of variability in atmospheric deposition³⁴. For wet deposition, rainfall volume per event, interval to last event, season and storm path can alter mass deposited in an event³⁵. Dry deposition is controlled by fundamentally different physical,

chemical, and biological processes than wet deposition. Dry deposition can be affected by the size spectrum of depositing particles, properties of the collecting surface, canopy structure, and season³⁶, as well as wind speed^{37,32}. Another key factor is sample contamination from nonatmospheric natural debris or that associated with birds or insects attracted to the collector device^{25,38,39,40,41,42}. As a result of these factors and others associated with atmospheric turbulence and boundary layer phenomena, dry rates are expected to be more variable than those for wet deposition^{12,34,43}.

The lack of precision and accuracy of dust deposition data in general and of P data in particular presents an immediate challenge for environmental management. For example⁴⁴, the vast majority of the remnant Florida Everglades is very shallow and often displays ambient P levels at or below $10 \mu\text{g P L}^{-1}$. Atmospheric loading in the middle or upper range of published values could be a significant source of enrichment to portions of this ecosystem^{42,45}. The Everglades Forever Act, passed by the Florida Legislature in 1994, requires the South Florida Water Management District, in cooperation with several other state and federal agencies, to restore the remaining Everglades ecosystem. The control of P loading to this oligotrophic, subtropical environment is a cornerstone of the restoration. Based on the literature reviewed for this study and data collected by various organizations in South Florida, a defensible P-loading estimate from the atmosphere, with well-defined precision and accuracy, does not currently exist for any location or for the region as a whole. This lack of definitive information on P inputs from the atmosphere is not atypical for most aquatic ecosystems and provides the fundamental motivation for this study. Gordon⁴⁶, summarizing the findings from a symposium on airborne particles, emphasized the need for atmospheric science to provide improved estimates of nutrient deposition to ecosystems. Likewise, Gardner⁴⁷ recommended more attention to P sources, availability, mechanisms of entrainment and modes of atmospheric deposition to better assess the importance of atmospheric P. Lovett³⁴ called for more attention to atmospheric deposition measurements for nutrients and pollutants in general, emphasizing the need to reconcile methodological uncertainties.

This review attempts to bridge the environmental and atmospheric sciences to convey information on the atmospheric deposition of P. The first objective of this paper is to provide a conceptual framework for information on P deposition in rainfall and dustfall based on published data. The second objective is to summarize P deposition rates in this conceptual framework and provide reasonable bounds for P deposition rates to aquatic ecosystems. Interpretation of this information is bolstered by a review of background deposition rates to

remote areas of the globe. Estimates of P deposition using nontraditional methods are also considered as a source of independent evidence on the accuracy of published values for application to environmental management and to South Florida, specifically. The third objective is to summarize available information on field methods that might be used to improve estimates of atmospheric loading of P to aquatic ecosystems. Finally, this paper concludes with a consideration of the circumstances, under which P deposition monitoring may reduce uncertainty for environmental decision making, considering constraints inherent in its quantification. The information analyzed to meet these objectives should be useful to environmental managers and ecosystem scientists worldwide as they seek to understand the sources and sinks of P in systems where nutrient enrichment is a target of environmental management. The environment of South Florida is used as a case study for illustrating concepts of atmospheric deposition and suspension. Vast areas of the Everglades receive little water directly from surface inflows and may have been influenced over many years by atmospheric deposition.

MAJOR FEATURES OF THE SOUTH FLORIDA ENVIRONMENT

Because examples from South Florida are used frequently in this review, the reader should have a general understanding of the regional ecosystem and its geography. The once vast Everglades landscape of wetland habitats has been reduced to about one-half its original size and is now a compartmentalized, highly modified system. The remnant Everglades is known collectively as the Everglades Protection Area for the purposes of the restorative Everglades Forever Act of 1994 (§373.4592, Florida Statutes) and is composed of four areas (Figure 1). Water Conservation Area 1 (WCA-1) is the Arthur R. Marshall Loxahatchee National Wildlife Refuge (Refuge) and consists of 566 km² of prime wetland habitats. This area receives runoff from the neighboring Everglades Agricultural Area to the northwest and is bordered by suburban development and agriculture to the east. Water from WCA-1 flows southward into Water Conservation Area 2 (WCA-2). This area of 538 km² also receives surface water inflow from the Everglades Agricultural Area and is adjacent to developed areas to the east. WCA-2 has been the subject of intensive eutrophication research and water quality monitoring over the last three decades⁴⁸. It displays a north-south gradient of marsh habitats, from cattail stands in the north, to sawgrass-dominated marshes in the south.

Water Conservation Area 3 (WCA-3; Figure 1), the largest of the WCAs at 2,342 km²,

contains tree islands, wet prairies, and sloughs scattered across a vast sawgrass marsh. WCA-3 has been less impacted by cultural eutrophication than WCA-1 or WCA-2; water from WCA-3 flows southward into Everglades National Park. Everglades National Park contains 5,569 km² of highly valued aquatic sloughs, marl-forming marshes, and mangroves. Vast areas of the park have very low total phosphorus levels in surface water (<10 µg L⁻¹) and receive most water from local rainfall. Atmospheric deposition is expected to be critical to the long-term nutrient status of these marshes.

CONCEPTUAL BASIS FOR P DEPOSITION FROM THE ATMOSPHERE

Components and Size Fractions

Atmospheric deposition is separated into three components (wet, dry and cloud deposition) in the following analysis of relevant concepts³⁴. Wet deposition is associated with directly falling atmospheric precipitation containing gases or particles from the atmosphere. Wet deposition can be expected to display the random variability associated with individual rainfall events, spatial patterns of precipitation and processes involved in scavenging of materials from the air. Jordan et al.³⁵ explained much of the temporal variation in wet deposition by considering season, rainfall volume, and elapsed time since the last event in a regression model of a long-term data set in the Chesapeake Bay area.

The separation between dry and wet deposition is defined operationally by the common Aerochem Metrics[®], Inc. wet/dry collection system that opens the wet bucket only when rain is falling in substantial amounts. Hicks et al.¹² define dry deposition as “the gravitational settling of large particles not associated with falling precipitation, and the turbulent exchange of small particles and trace gases to the surface.” This definition implies that variations in dry deposition are under the control of extremely different physical, chemical and biological factors than wet deposition. Dry deposition rates are determined by wholly different mechanisms from wet deposition, including particle size and concentration in the atmosphere, settling velocity and the nature of the collecting surface. Dry deposition is fundamentally more complex and variable than wet⁴³, and is very important in conveying particles to the land surface³⁴; dry deposition is discussed in greater detail below.

Cloud deposition occurs when fog or clouds contact surfaces and is important in elevated coastal or mountainous environments. Deposition from clouds and fog may be difficult to

separate into wet and dry components; on a practical basis, however, it is measured by dry techniques. Although fogs do occur sporadically across the South Florida landscape, it is unlikely that fog/cloud deposition is a significant part of long-term phosphorus input to this low-elevation subtropical environment. Cloud deposition will not be considered further in this analysis as a distinct process. Any approach to sampling the dry and wet components assumes that materials conveyed by clouds and fog are captured. Also, dew will be considered as a factor altering dry deposition through surface wetting, as suggested by Lovett³⁴, rather than as a separate form of deposition.

Phosphorus is conveyed through the atmosphere almost entirely by suspended particles. However, there is a gaseous component to the P cycle involving phosphine (PH₃), diphosphine (P₂H₄) and phosphorus pentoxide (P₂O₅) that may be an important vehicle for atmospheric P transport under some circumstances. Phosphine gas can be emitted from anaerobic environments in which energy and phosphate are available, such as sewage plants, marsh and river sediments, landfills, or composting soils⁴⁹. Devai et al.⁵⁰ found that up to 50 % of the P mass balance deficit for stabilization lakes associated with two sewage treatment plants in Hungary could be explained by the emission of phosphine gas into the atmosphere. Measurements in marsh sediments, including samples from the Everglades, confirmed that phosphine can be emitted from natural marsh sediments under strong reducing conditions, and emission rates can be enhanced when given additional P and an energy source⁵¹. Newman²⁷ mentions another potential gaseous pathway when plant P is oxidized to P₂O₅ (phosphorus pentoxide) and vaporized during biomass burning. All these compounds will form phosphoric acid when they contact water in the air and will deliver P to an ecosystem as phosphate, a biologically available form. While existing evidence suggests that the gaseous phase of the P cycle is ubiquitous, the quantitative importance of PH₃, P₂H₄ and P₂O₅ to atmospheric deposition is not known. Gaseous forms of P are worthy of additional attention, particularly in the vicinity of large sources, such as landfills, eutrophic marshes and sewage treatment plants.

The dry fallout of P is due almost entirely to particles greater than 2 μm in diameter^{52,7}. Phosphorus can be conveyed in large particles derived from soils and dominated by other crustal elements, such as calcium, silicon, iron and magnesium. Particles containing P may also be biological in origin, including pollen, spores, waxes, resins and microbes^{53,54}. More recently, several studies have demonstrated clearly that dry deposition flux of crustal elements is almost entirely derived from large atmospheric particles^{32,55,56,57,33}. Although the following discussion is organized around the three common size fractions of atmospheric particles (fine, coarse and

giant; Table 1), a conceptual framework for P dry deposition must concentrate on the nature and behavior of larger coarse and giant atmospheric particles. General discussion of the size distribution and sources of atmospheric particles can be found in Finlayson-Pitts and Pitts⁵⁸, Harrison and Perry⁵⁹, Chow³¹ and United States Environmental Protection Agency⁵⁴.

Fine particles (<2.5 μm in diameter) (Table 1) have the potential to travel long distances in association with stratospheric or tropospheric air masses and may have residence times in the atmosphere ranging from weeks to months, depending on mixing and removal processes⁶⁰. Condensation, coagulation and nucleation alter the size distribution of fine particles in the atmosphere due to the formation of secondary particles from smaller ones, decreasing the relative proportion of the smallest particles^{58,59}. Simultaneously, coarse and giant particles are lost from the atmosphere at higher rates. The deposition of aerosols thousands of kilometers from sources reflects the net effect of these processes. Intercontinental dust transport is a classic example of long distance particle movement by fine and smaller coarse-sized particles (Table 4). Studies of such transport suggest that factors other than gravitational settling maintain particle suspension in the atmosphere and allow long distance movement of particles^{61,62}.

Giant particles (10 to 100 μm , Table 1) have high settling rates based on the balance between gravitational settling and viscous atmospheric drag predicted by Stokes' Law; Noll and Fang⁶³, Lin et al.⁵⁶ and Caffrey et al.⁵⁷ provide a guide to information on settling rates. It is tempting to conclude that such high loss rates would preclude giant particles from traveling far from sources. However, other evidence suggests that particle transport is a function of complex and poorly understood atmospheric processes, with the net result of longer suspension and transport than Stokes' Law behavior or more sophisticated computational methods would suggest. An excellent example of giant particles in atmospheric deposition is provided by Betzer et al.⁶² for the North Pacific Ocean. They reported giant (> 75 μm) atmospheric particles at remote marine locations, and established that mass flux from these particles was the dominant component of flux associated with a dust storm event. Betzer et al.⁶² could not explain how such particles could remain suspended and travel over 10,000 km from China, but they suspect that resuspension by convective storms may be involved. Ellis and Merrill⁶¹ also found Stokes' Law predictions did not accurately predict long-range transport to Barbados during certain times of the year. The Barbados case suggests depositional velocities are less than Stokes' Law predictions. Through analyses using a model to predict dry deposition rates, Noll and Fang⁶³ concluded that the tendency for particles to remain suspended is a complex function of both inertial and gravitational forces. Their model predicted that larger (ca. 15 μm) particles tend to

respond more to increased wind speeds than smaller particles and therefore have the potential to remain in the atmosphere longer than might be predicted by Stokes' Law settling velocities. There is other evidence from *in situ* measurements that larger particles stay aloft long enough to dominate dry deposition far from sources. For example, Holsen et al.⁵⁵ found that over 98 % of dry deposition to Lake Michigan was due to particles greater than 6.5 μm . Analysis of data from Chen et al.⁶⁴ (Table 5) revealed that over 85 % of P was conveyed by either coarse or giant particles in maritime breezes.

Davidson et al.^{65,66}, Noll and Fang⁶³, Lin et al.⁵⁶ and Zufall et al.³³ provide overwhelming evidence that the vast majority of dry deposition mass flux is conveyed by giant particles between about 10 and 100 μm . In addition, Brook et al.⁶⁷ found that particles greater than 10 μm represented about 44 % of total suspended particles for 14 urban monitoring stations across Canada, while Lundgren et al.⁶⁸ noted substantial particle mass occurring in sizes greater than about 15 μm in five U.S. cities. Paode et al.⁶⁹ provide evidence that the dry deposition of lead, copper and zinc, traditionally thought to be conveyed by fine particles, is dominated by particle flux from coarse and giant particles. Although giant particles are usually present in low concentrations, their high sedimentation rates (according to Lin et al.⁵⁶, a 10- μm particle falls at about 1.0 cm sec^{-1}) and high mass per particle (one 10- μm particle has the mass of 1000 one- μm particles) provide for high deposition flux. There is no reason these findings on the importance of giant dust particles to dry deposition rates should not apply equally to P-bearing particles. Giant particles (10 to 100 μm) must be quantified in any valid estimate of P deposition rate.

Defining which particle size is truly atmospheric is not a straightforward process. The dilemma, as emphasized by Hicks et al.¹², is how to decide what particulate matter is a natural part of atmospheric deposition and what is caused by contamination of the sampler or by local, nonatmospheric inputs. The objective of measuring P deposition for environmental management or research is to estimate total input of P to an ecosystem. Any pragmatic size cutoff point for atmospheric particles consider account inputs that could be missed as a result of the criterion. Studies dating from the 1950s suggest that natural organic matter from the air may be a very important source of carbon or other elements^{1,70}. Ecological studies have reported that particulate debris up to 1 mm in diameter can represent an important source of depositional P^{71,72,5}. The importance of debris inputs may be scale-dependent and small aquatic ecosystems surrounded by forest may be exceptional in the amount of material delivered via air transport of large particulate detritus. In fact, there are estimates of substantial deposition from airborne

particles greater than 1mm in diameter, known as *large particulate organic matter* in the ecological literature^{73,74}.

Natural detritus or very large dust particles in the air are difficult to separate from sample contamination by bird excrement, insects or organic matter from the area immediately surrounding the sampler. For larger ecosystems, the input of localized, large particulate organic matter is less important, and sample censoring, screening, filtration and other deposition methods that do not sample particles greater than 100 μm will affect overall deposition rates to a relatively minor extent. For most systems, atmospheric transport should be considered an important mechanism for particles of 100 μm or less in diameter. Some bulk deposition sampling programs preclude all larger materials from samples by physical means: van Ek and Draaijers⁴³ and Anttila et al.⁷⁵ used filtration to remove debris from samples; Brown and Lund⁷⁶ used cotton in funnels to keep larger materials out of samples. Other sampling programs use netting to avoid all particles greater than about 1 mm^{2,77}. Most particles above 100 μm in size will travel relatively short distances. Using 1-mm screening or collection methods that do not sample particles greater than 100 μm in size is appropriate to avoid local debris without interfering with the input of true atmospheric fallout.

Major Sources of Variation in P Deposition Rates

Many processes lead to natural changes in atmospheric particle concentrations and deposition rates (Table 1). Changes in the location and magnitude of sources for P-bearing particles can provide variation in atmospheric P inputs to ecosystems. Phosphorus-bearing particles in deposition from the atmosphere are derived largely from manufacturing, fugitive soil particles, pollen, insects and other biological debris, fertilizers, ashes due to volcanic activity, combustion of coal and other fossil fuels, organophosphate pesticides, and aerosols from the ocean^{26,27}. Phosphorus can be conveyed through exhaust from vehicles or incinerators when many common materials are burned, including some complex organic compounds, such as organophosphoric acid triesters used in plastics, polyvinylchlorides and lubricants⁷⁸. Many sources of atmospheric P are altered by human activities, and it is, therefore, not surprising that long-term variations in dust concentrations in the atmosphere can be traced to land-use changes; Prospero et al.⁷⁹ even consider emitted dust as a global pollutant.

Unfortunately, unnatural variation in rates from local contamination is a very common problem and can dominate values generated by long-term sampling with surrogate surface collectors, particularly in productive wetland environments. Screening or sample filtration

cannot prevent occasional contamination, and censoring of samples that have been compromised by contamination is essential to estimating deposition rates. Using data from wet/dry bucket monitoring in South Florida, Ahn^{38,39} and Ahn and James^{40,41,21} provide quantitative means for detecting outlying data, filling data gaps through statistical modeling and assessing uncertainty in data compromised by contamination. These authors used a multiple-level culling approach to identify contamination in over one-third of both wet and dry samples. However, following statistical treatment of both wet and dry deposition data, no differences were visible between stations with widely differing surroundings or in monitoring from different years. Even though their mean deposition rate was quite reasonable, the general lack of temporal or spatial signals in the post-censored data is not expected based on concepts of particle deposition. The absence of a clear signal in the final data set suggests that the statistical removal of contaminated data was not able to clarify the underlying deposition signal.

Walker⁴² also used wet/dry bucket data from South Florida to derive estimates of P deposition to the Arthur R. Marshall Loxahatchee National Wildlife Refuge in the Northern Everglades (Figure 1) using a different approach to data analysis. With over five years of weekly data from four stations, Walker⁴² assessed the sensitivity of both wet and dry data to various sources of contamination. He found that over one-quarter of wet deposition samples and one-half of dry deposition samples could not be analyzed due to gross contamination or equipment failure. With the remaining data, he conducted a thorough analysis of effects of contamination on deposition estimates from each of the four stations and found that droppings, vegetation, insect debris and spider webs were common sources of contamination from within the marsh environment. Even with many contaminated samples removed, rates from weekly data still displayed variation over about two orders of magnitude. His evaluation demonstrated that deposition rates from collectors, such as wet/dry buckets, are altered dramatically depending on how contamination is treated and censored in data analysis. The four stations examined by Walker⁴² in the Refuge ranged from less than 20 to over 100 mg Pm⁻²yr⁻¹ and rates appear to be more associated with the characteristics of the sampling sites, rather than a reflection of the incoming atmospheric deposition. The analyses of both Walker⁴² and Ahn and James²¹ document that it is not possible to completely separate or remove the effects of contamination from data sets in wetland environments. As a result, any long-term data from bulk, wet/dry or other surrogate surface devices yield data with some unrepresentative high values and rate estimates that are likely to be biased high. This bias should be borne in mind in a review of published deposition rates to be considered in the next section of this paper. Screening

bulk collectors, keeping careful field notes during sample collection and removing extraneous material from samples may help reduce the frequency of gross contamination, but these practices cannot eliminate the problem. Likewise, *post hoc* statistical treatment removes some variation generated by extraneous material, but is not able to effectively remove noise and leave a contaminant-free signal.

Another important source of variation in P deposition flux is the elemental composition of depositing particles. The P content of atmospheric dust particles ranges from about 0.02 to over 2.0 % of the mass, and varies with particle source. An extensive review by Chow³¹ revealed that P in particles from road dust, soil, vehicles and power plants occurs at levels between 0.1 and 1 %. Particles of crustal origin can be expected to contain P at about 0.1 to 0.2 %²⁶; Prospero et al.⁷⁹ found 0.105 % to be a defensible average for estimating P transport from data on dust. Avila et al.⁸⁰ reported the P content of red rain dusts from the soils of North Africa to be 0.063 %. The P content of sedimentary rocks is from 0.02 to 0.07 %, and for igneous and surficial rocks from 0.06 to 0.13 %, according to Newman²⁷. Fine particles in exhaust from gasoline-powered cars were found to contain 0.11 % P, and diesel exhaust contained 0.06 % P⁸¹. Olmez et al.⁸² provided data showing high P levels in particles from industrial sources. For the fine fraction, they found concentrations of 0.45 to 1.5 %, and for coarse particles concentrations of 0.22 to 2.3 %; particles from coal-fired power plants contained P at significant concentrations of 1.0 to 2.3 %. Other biological particles have higher P content than crustal materials; 13 types of pollen showed an average of 1.6 % P in a study by Banks and Nighswander⁶. For purposes of this study a conservative value of 0.1 % will be used to estimate P deposition from data on total particle mass. However, the reader should recognize that particles of anthropogenic or biological origin could be an order of magnitude higher in P content.

Surface effects can be very important to the deposition rates measured by various methods and can generate variation at the landscape scale. Both Cole et al.⁷² and Jassby et al.⁵ used wet vessels to estimate dry deposition to a lake surface and found that the wetted surface captured much more P than the same containers without water. Caffrey et al.⁵⁷, working on Lake Michigan, suggest that the lake surface can be expected to have higher rates of dry deposition than a flat surface due to surface roughness, whitecap formation and humidity gradients. Zufall et al.^{83,84} used modeling and wind tunnel measurements to confirm that wavy surfaces generally collect more particles than flat surfaces, and that giant particles may be less affected than smaller particles by the presence of waves. For elements such as P being conveyed in coarse and giant particles, water surfaces provide for less resuspension and bounce-off, resulting in greater

net flux than a dry, flat surface. However, Hicks et al.¹² argued that the open water surface does not necessarily trap particles as efficiently as might be suggested by data from containers. Air flow over a wetland or lake water surface may tend to be stratified during midday, when dry deposition rates are highest, and although exchange with the air may be greater at night due to convective mixing, the air layer that is cleansed of particles may be quite shallow. Also, as noted by Holsen et al.⁵⁵, water surfaces cool and heat more slowly than land so that air movements in spring and summer may be more restricted, while in the cooler months warmer waters may enhance vertical mixing of air masses. The net effect of all these processes remains uncertain, particularly when compared to the complex effects of landscape structure. Clearly, more definitive information on deposition to water surfaces is needed to predict deposition rates for very large water surfaces, such as Florida's Lake Okeechobee and the Florida Everglades, with reasonable certainty.

Davidson et al.³⁷ stressed the importance of the collecting surface to deposition rates, and noted earlier work suggesting that deposition rates can vary over several orders of magnitude based on fine structure or geometry of the collecting surface. Thus, it would not be surprising to find significant differences in dry deposition rates in the same area within different plant canopies. The canopies tend to act as a sink for coarse and giant dust particles from the atmosphere⁸⁵, but may be a source of particles through resuspension and biological processes. However, vegetation processing within a canopy can alter the apparent deposition rate either through uptake or leaching, as discussed below concerning the measurement of canopy throughfall.

Edge effects occur both within and outside vegetation canopies. There are edge effects in forest stands so that deposition rates within five stand-heights of the edge are higher than those in the interior of the stand (reviewed by Erisman et al.⁸⁶). Weathers et al.⁸⁷ reported that cloud deposition was reduced by 50 % at 28 m from forest edges. Conversely, as one moves away from a canopy source, the deposition of plant-generated materials can be expected to decline greatly with distance from the stand edge^{88,70,74,72,6}. This same phenomenon can be seen at a large scale in atmospheric P concentrations moving away from the coast; levels near Bermuda about 1000 km offshore are approximately an order of magnitude less than P concentrations at the North American coastline⁷. For Southern Lake Huron, Delumyea and Petel⁸⁹ found across-lake gradients in P-bearing particles when the wind was moving offshore, and used these gradients in concentration to calculate deposition velocities for P. The challenge is how to estimate net deposition to large and complex natural surfaces, such as the Florida Everglades. The Leaf Area

Index has been found to be a quantitative means of scaling data from surrogate surfaces to estimate deposition to the forest floor⁹⁰. Scaling factors clearly demonstrate that the capture of particles by plant canopies is important for any landscape-level estimate of deposition.

Processes responsible for particle concentrations in the atmosphere result in daily and seasonal variations in deposition rates. Deposition flux was found to be 65 % lower at night due to lower wind velocity, lower particle concentrations, and a shift toward smaller particles at night^{91,32,92}. Eisenreich et al.⁹³ (Lake Michigan area, U.S.A.) and Pratt et al.⁹² (Minnesota, U.S.A.) found that rates were highest in summer. Shaw et al.⁹⁴ found that nitrogen (N) and P deposition rates were highest in May and remained relatively high during the summer months in Central Alberta, Canada. They also noted that monthly P loading from the atmosphere did not correlate closely to precipitation; N loading corresponded more closely to rainfall amounts. Jordan et al.³⁵ reported that both N and P deposition rates were highest in the spring in Maryland, U.S.A., mimicking the pattern of precipitation. In the Czech Republic, Kopacek et al.⁹⁵ found deposition rates for P and N to be two-to-three times higher during the period from April through July than during the winter months. Yang et al.⁹⁶ found that both wet and dry deposition rates for P were highest in the spring, followed by the summer quarter for three sites in Connecticut, U.S.A. Brook et al.⁶⁷ also noted that air concentrations of particulates tended to be higher in the summer than in the winter for Canadian cities. However, these same authors also provided data demonstrating that the relative amounts of fine, coarse and large particles vary greatly between sites and sampling periods. Finally, Prospero et al.⁷⁹ cited data on dust concentrations in Barbados with higher levels in summer and long-term changes associated with climatic cycles in North Africa.

When all these factors are combined (Table 1), the net effect is that fine particles tend to deposit gradually and more continually than larger fractions. However, this component of deposition is a trivial contributor to P flux due to very low mass per particle and low settling rates. The coarse and giant particle fluxes are more variable and event-related, since high settling rates require greater atmospheric turbulence to keep particles in suspension. Holsen et al.⁵⁵ concluded that Lake Michigan received most of its dry deposition during periods of strong southerly winds from the urban areas to the south of the lake. Lin et al.⁵⁶ provided direct evidence of high deposition rates associated with strong winds. Prospero et al.⁷⁹ concluded that dust transport varies with geographic sources and meteorological conditions, particularly transporting winds. These factors produced clear seasonal and annual signals in particle concentrations spanning 27 years of sampling on the island of Barbados. The P-deposition data

of Bergametti et al.⁹⁷ also demonstrated episodic and seasonal signals for the Northwestern Mediterranean Sea. Large particulate organic matter will generate variability in deposition rates stemming from transport during windy, high-energy events and/or pulses of biological production, such as pollen release, insect emergence, leaf abscission, epidemics of plant disease, or other natural occurrences. Quantifying the coarse and giant fractions (Table 1) will require sampling strategies capable of capturing rapidly settling particles while minimizing the bias associated with artificial surfaces and contamination. Wind speed and direction and other meteorological data may prove useful in reducing noise and accounting for variation in dry deposition rates.

P Deposition: Integration from an Ecological Perspective

A conceptual model for South Florida provides some insight into the sources, sinks and processes across the landscape (Figure 2) and shows the dynamic nature of atmospheric particle concentrations and deposition rates spanning multiple temporal and spatial scales. Intercontinental and regional inputs are augmented by local sources, particularly those associated with urban or agricultural land uses, as air masses travel across the landscape. Fine particles may coalesce and form larger aggregates. As this particulate matter is carried over a large natural area or water surface, such as the Everglades, larger particles are deposited at high settling velocities. Particle concentrations gradually decline as deposition exceeds suspension. Convective storms promote scavenging of particles from the troposphere. Some particles will be emitted from natural sources, and resuspension of particles is possible. Several studies of atmospheric deposition have suggested that materials can spiral across terrestrial systems, with the atmosphere as the agent of transport^{66,98}. To the extent that this phenomenon occurs in atmospheric transport, it may involve active resuspension or simply the rebounding of larger particles³⁷.

The United States Environmental Protection Agency¹⁴ provides an excellent summary of the scales of phenomena that affect particle concentrations. Residence time in the atmosphere links time and space scales for sources and sinks of particles and is strongly dependent on particle size and density. Obviously, the timing of rainfall events will dictate when wet deposition occurs, and convective storms are most common after midday from June through October over large areas of North America. Longer-lasting frontal storms tend to occur in the fall and winter and may bring with them strong winds carrying terrestrial or marine particles. Similarly, for dry

deposition weather patterns and long-term climatic cycles will alter the trajectory of atmospheric dusts from particle sources spread unevenly around the globe.

Considering the number of factors involved in particle production, transport, and sedimentation (Table 1; Figure 2), temporal and spatial variability in P deposition rates are guaranteed. The studies reviewed next in this paper tend to smooth out some of this inherent variability through biweekly or monthly composite sampling and by averaging data from local sites in an attempt to derive representative annual rates. Data on time and space variation in concentrations of particles in the ambient air provide direct confirmation of the relative scales of variation that will ultimately affect the annual P deposition rates summarized in Tables 2 and 3. Graham and Duce²⁶ reported that P concentrations in the atmosphere over cities range from 30 to 1450 ng P m⁻³, with an average of about 150 ng P m⁻³, while values in coastal environments tend to be lower, in the range of 10 to 20 ng P m⁻³. They also found oceanic values to be highly variable; higher values (ca. 5 to 55 ng) occurred near continental sources, and very low concentrations, sometimes below 1 ng P m⁻³, occurred in the open marine environment. Low P levels of 1.3 and 0.2 ng P m⁻³ were also reported by these authors for remote continental areas: Northwest Territories, Canada and the South Pole, respectively. For 33 samples from the western North Atlantic Ocean, Graham and Duce⁷ found values ranging from 0.6 to 21.9 ng P m⁻³, with an average of 6.4 ng P m⁻³. Brook et al.⁶⁷ reported median concentrations for Canadian cities ranging from 10 to 36 ng P m⁻³ for particles less than 10 μm; concentrations for total suspended particles can be expected to be about twice these amounts. Prospero et al.⁷⁹ summarized monthly P air concentration data from Barbados from 1965 to 1992 and documented order of magnitude variations associated with seasons and climatic cycles. In essence, concentration data tend to follow expectations based on particle sources and transport: higher values in developed areas and low concentrations in undisturbed, remote environments. Phosphorus concentrations in combination with meteorological conditions set the stage for spatial and temporal variations in P deposition rates from the atmosphere.

Capturing the variability of P deposition associated with landscape patchiness remains a challenge. Plant canopies in aquatic or terrestrial systems can be expected to increase particle capture, but this effect might be less significant for giant particles than for pollutants conveyed by fine aerosols. In any case, scaling factors to translate data from surrogate surface collections into rates for various cover types in a regional landscape mosaic are not currently available. Quantifying spikes in deposition is also a challenge. For dry P deposition driven by coarse and giant particles (Table 1), variability generated by rare, windy atmospheric events and associated

high suspension, transport, and deposition rates must be quantified. Annual rates could be more affected by episodic inputs than by more common deposition rates during fair weather.

PUBLISHED RATES FOR P DEPOSITION FROM THE ATMOSPHERE

The second objective of this study was to summarize P deposition rates from the scientific literature in light of the conceptual framework provided above and to assess these estimates for applicability to environmental management and to South Florida specifically. There are hundreds of annual values for atmospheric P deposition published in the literature. This investigation seeks to build on earlier literature reviews^{99,26,93,29,28,27} and provide a synoptic view of annual rates to bracket long-term values that can be expected in aquatic ecosystems surrounded by various land uses.

The vast majority of published rates of P deposition are estimates from *bulk* measurements in which the sampler, often a funnel, is left exposed to both wet and dry deposition for extended periods, usually one to two weeks. Atmospheric scientists are often critical of deposition estimates from bulk precipitation, especially for any substance conveyed by fine particles. Bulk collectors and the common Aerochem Metrics® wet/dry collectors are not designed aerodynamically, so that smaller particles will not be sampled efficiently, and captured particles may be resuspended from dry buckets. All surrogate surface sampling can be influenced to varying degrees by local contamination. This constraint, plus problems associated with the geometry and aerodynamics of the collector, must be recognized as these data are considered. While highlighting these considerations, Hicks et al.¹² note that bulk collectors or other artificial surfaces become more efficient as the particle size increases into coarse and giant particles controlled by sedimentation from the atmosphere. While the drawbacks of bulk collections are undeniable¹⁰⁰, the lack of practical alternatives for long-term sampling of coarse and giant particles encourages many investigators to continue publishing bulk deposition data or values from the common wet/dry bucket approach. The evidence compiled in this study will establish that many published bulk rates for P deposition are in reasonable agreement with data derived from other independent approaches to estimating deposition, and tend to be internally consistent in range and average magnitude for various land uses. Data from bulk or wet/dry collections are not reliable for any material transported primarily by fine particles (lead, sulfur or nitrogen) or as gases (O₃, SO₄, NO₄). Until estimates of P deposition are available in a variety of ecosystems using other methods, bulk or wet/dry deposition data provide the best available estimates of

long-term atmospheric inputs for most management and research purposes.

Annual mean rates of P deposition for specific locations and using bulk and Aerochem Metrics[®] wet/dry measurements range from 12.4 to 93.3 $\text{Pm}^2\text{yr}^{-1}$ (Table 2). Regional mean P deposition rates reflected in earlier literature summaries on the subject range from 23.8 to 65.7 $\text{Pm}^2\text{yr}^{-1}$. Although the complete range of published annual values in the cited references for both tables is from 3.6 to 200 $\text{Pm}^2\text{yr}^{-1}$, most published deposition rates fall between 20 and 80 $\text{mg Pm}^2\text{yr}^{-1}$. An earlier summary of 45 published values by Gibson et al. yielded a smaller range in annual rates (5 to 60 $\text{mg Pm}^2\text{yr}^{-1}$). His summary had a mean of 26 $\text{mg Pm}^2\text{yr}^{-1}$, which is very close to the mean value of 34.8 $\text{mg Pm}^2\text{yr}^{-1}$ in Table 2 of this study. Newman²⁷ (1995) scrutinized published bulk deposition values from funnel collectors where technically sound methods were documented. Compared with local values in this study (Table 2), Newman's data resulted in a higher mean of 65.7 $\text{mg Pm}^2\text{yr}^{-1}$, with a nearly identical range of 7 to 170 $\text{mg Pm}^2\text{yr}^{-1}$. Only two sources of data were used in both Newman²⁷ and this study: Graham and Duce²⁶ and Schindler et al.²

In marine environments, deposition tends to occur at relatively low rates in the offshore environment despite the P-enriched surface microlayer that can be a source of atmospheric P¹⁰. For example, Graham and Duce²⁶ reported a value of 7 $\text{mg Pm}^2\text{yr}^{-1}$ from coastal Rhode Island, U.S.A., and in Florida, Hendry et al.²³ (1981) reported a low deposition rate of about 17 $\text{mg Pm}^2\text{yr}^{-1}$ for a maritime station in the Florida Keys. Guerzoni et al.¹⁰¹ concluded that P deposition to the Mediterranean Sea ranges from about 20 to 40 $\text{mg Pm}^2\text{yr}^{-1}$, with higher values near the shore.

Inland sites associated with northern forests of the Midwestern United States or Canada, and one site from a tropical dry forest in Mexico, tend to have deposition rates of less than about 40 $\text{mg Pm}^2\text{yr}^{-1}$. Rates in the middle of the range (30 to 50 $\text{mg Pm}^2\text{yr}^{-1}$) represent a variety of aquatic systems surrounded by mixed land uses (Tables 2 and 3). Values of 50 or more tend to be associated with productive ecosystem sites or agricultural and urban land uses. The mean total N:P mass ratio in annual values of Table 2 is 25, and the N:P ratios tend to decrease as P deposition rates increase. Thus, P limitation of primary productivity will tend to be maintained or enhanced by atmospheric nutrient inputs in less-disturbed forested landscapes. Jassby et al.⁵ (1994) concluded that the N:P relationship in atmospheric deposition to Lake Tahoe (CA/NV, U.S.A.) was consistent with the observed shift from N to P limitation in this oligotrophic subalpine lake.

Together, the information on mean rates and ranges in Tables 2 and 3 provides estimates of deposition useful for assessing the relative importance of atmospheric inputs into ecosystems. However, it is vital to recognize that the values compiled in Tables 2 through 5 are published arithmetic means and represent upper bound deposition rates for the various locations being studied. The use of geometric means or medians in the original data would reduce these values, particularly in those data sets subject to high variability. Considering this caveat, and using South Florida as a case study, we should expect values less than $20 \text{ mg Pm}^{-2}\text{yr}^{-1}$ for sparsely populated areas near the Florida coast receiving mostly onshore winds. Inland rates will vary with land use and distance from sources. High P inputs (60 to $100 + \text{ mg Pm}^{-2}\text{yr}^{-1}$) can be anticipated near urban or agricultural areas, dropping off as one moves into open-water areas of the Everglades; Hendry et al.²³ reported the high values for urban ($48 \text{ mg Pm}^{-2}\text{yr}^{-1}$) and agricultural ($66 \text{ mg Pm}^{-2}\text{yr}^{-1}$) areas of Florida.

A spatially integrated rate for the remnant Everglades marshes must consider urban inputs from the east, agricultural sources to the north and east, and maritime areas to the south and west (Figure 1). It is reasonable to expect low deposition rates over large areas of Everglades National Park and Water Conservation Area 3, distant from local sources, and higher rates in the northern parts of the remnant system, including Water Conservation Areas 1 and 2. With these facts in mind, a spatially integrated, long-term average input of atmospheric P to Everglades marshes, known collectively as the Everglades Protection Area, should fall between 30 and $40 \text{ mg Pm}^{-2}\text{yr}^{-1}$, based on published values in the literature. McDowell et al.¹⁰² also arrived at this range based on their analysis of P deposition into South Florida. The upper bound of this range is close to the $40.9 \text{ mg Pm}^{-2}\text{yr}^{-1}$ 13-station mean of Ahn and James²¹. The values developed by Walker⁴² in the Arthur R. Marshall Loxahatchee National Wildlife Refuge (WCA-1, Figure 1) bracket rates expected as background (ca. $20 \text{ mg Pm}^{-2}\text{yr}^{-1}$) and as levels associated with high anthropogenic inputs (ca. $100 + \text{ mg Pm}^{-2}\text{yr}^{-1}$). The earlier $37\text{-mg Pm}^{-2}\text{yr}^{-1}$ estimate used by Walker¹⁰³ for the Northern Everglades appears reasonable in light of the values summarized in this study. Finally, a value of $30 \text{ mg Pm}^{-2}\text{yr}^{-1}$ was used by Fitz and Sklar¹⁰⁴ in a spatially explicit ecosystem model of Water Conservation Area 2A in the Northern Everglades. Their model was found to provide good agreement with observed data, including data on P in WCA-2A.

P deposition rates for remote areas of the globe illustrate that there is no place in the world where one would expect the rate of P deposition from the atmosphere to be zero (Table 4). There is evidence that areas thousands of kilometers from Asian or African sources of dust still

receive from about 1 to 5 mg Pm²yr⁻¹ from atmospheric transport. Although these estimates seem small relative to local recycling, they can be significant over long periods in oligotrophic tropical settings¹⁰⁵. The two largest background values are based on long-term sediment P accumulation rates from the Hawaiian Islands and the Amazon Basin, receiving P loading almost entirely from the atmosphere by long-distance transport (Table 4). The rates at the bottom of Table 4 are derived from sediment P accumulation rates and overlap with the range of values seen in remote marine and forested settings over the last few decades (Tables 2 and 3).

The sediment P accumulation values for the unenriched sections of Florida Everglades marshes in Craft and Richardson¹⁰⁶ and Reddy et al.¹⁰⁷ are higher at about 70 and 150 mg Pm²yr⁻¹, respectively. These rates were not included as background values because they have been influenced to some uncertain, but significant, extent by water management and associated inputs of nutrient-rich stormwater. Walker¹⁰³ used these P accumulation data to derive a relationship between P accretion and distance from surface inflows in WCA-2 of the Northern Everglades. As P concentrations fall to near-background levels over 10 km from inflows, Walker's model predicts P accretion rates of about 75 mg Pm²yr⁻¹. The proportion of nonatmospheric loading in this rate is uncertain, but atmospheric P deposition clearly must be substantially less than this value over vast areas of the Everglades far from the influence of surface inflows. Everglades P accretion data are within the range of bulk deposition values established for other ecosystems in regions with mixed land uses. Because the rates are a composite of atmospheric and surface water loading, the 30 to 40 mg Pm²yr⁻¹ overall deposition rate for the remnant Everglades is reasonable.

The bulk deposition values (Tables 2 and 3) are also consistent with rates derived from techniques using concentration data and size-specific deposition velocities, inverted Frisbee collectors or Petri dish samples combined with funnel sampling of rain (Table 5). Data from urban Chicago sites range from 47 to 135 mg Pm²yr⁻¹, rural sites range from 14.6 to 32.5 and aquatic sites range from 5.9 to 15 (both rates are biased low)(Table 5). Most of these data for concentrations and settling velocities to flat surfaces are for short sampling periods, and extrapolation to annual values is tenuous. Overall, the values of Tables 4 and 5 support the conclusion that inherent uncertainties in estimates from bulk or Aerochem Metrics[®] wet/dry collections do not preclude the use of this approach for setting bounds on atmospheric P deposition rates expected for various land uses. The vast majority of these data was derived from lake or forested environments in which contamination may have less influence on average values than would be the case for wetlands.

FUTURE MEASUREMENTS OF ATMOSPHERIC P DEPOSITION

Are the values compiled in this study useful as defensible estimates for decision making in ecosystem management or research? To address this question, each manager will have to assess the potential importance of atmospheric inputs to decide technical effort to give in reducing the uncertainty of P inputs from the atmosphere. In oligotrophic systems with low hydraulic loading rates, atmospheric influences are more important and may be greatly affected by land use changes. Regarding South Florida, the majority of P loading to the pristine portions of the Everglades ecosystem comes directly from the atmosphere, and the remnant system is, as a whole, very oligotrophic. In fact, there is strong evidence that a P water quality criterion for the ecosystem of about $10 \mu\text{g PL}^{-1}$ is scientifically defensible⁴⁴, and major changes in P deposition could alter water column P concentrations in this system. Based purely on the important role of P in this ecosystem, more accurate and precise information would be desirable. However, the need for better information must be tempered by practicalities. From a resource-management perspective, most atmospheric non-point source inputs of P-bearing particles are not readily abatable, at least on a regional level. If the rates compiled to date are found to be unrepresentative and overly conservative for localized sources from industry, agriculture, or urban areas through mass balance studies or water quality monitoring, then further quantification might be considered for the Everglades or any other managed ecosystem. The second practical consideration is the challenge of improving upon existing bulk or wet/dry data, with methods suitable for long-term quantification; this aspect is considered next.

When additional measurements are warranted for a particular ecosystem, how should they be done? Published bulk precipitation or Aerochem Metrics[®] wet/dry measurements provide rough rate estimates, but the drawbacks of this approach, particularly contamination and inherent characteristics of bucket collectors, suggest that additional data will be of limited value in improving the predictability of P deposition rates beyond currently available information. New options for long-term monitoring of coarse and giant particles are limited, however, because recent methodological advances in long-term atmospheric environmental measurements have focused on acid precipitation and other pollutants with gaseous forms or in fine aerosols³⁴. Many of the extant monitoring programs reflect a justifiable interest in determining the deposition of N, sulfur, ozone and other constituents with potential impacts on the environment¹². Unfortunately, this emphasis has produced elegant techniques primarily

applicable to gases or aerosols in the fine size fraction of atmospheric particles. Some long-term programs have looked at particle inputs, but new approaches have not been tested for P. There are promising methods for quantifying coarse and giant particles, but none have been used on a multiyear basis or perfected for application in remote locations.

The analysis that follows attempts to guide the environmental manager through available methods to arrive at recommendations on how to improve estimates of P deposition rates from the atmosphere, particularly for the dry component. The evaluation is organized around the recommendations of Hicks et al.¹², Erisman et al.⁸⁶, Lovett³⁴ and McDowell et al.¹⁰². These authors emphasize that wet and dry deposition should be monitored separately; dry deposition conveys most atmospheric loading; and estimates of P deposition will be most improved with several complementary techniques used simultaneously.

Wet Deposition of Phosphorus

Automatic wet/dry deposition collectors are recommended for wet deposition sampling⁸⁶. The Aerochem Metrics[®], Inc. device is used commonly for measuring wet deposition and occasionally for measuring dry deposition. These samplers consist of a wet and dry bucket, each 29.4 cm in diameter. The wet bucket is uncovered only during periods of rainfall, while the dry bucket is left open all the time, except for periods of active precipitation. For wet deposition, wet/dry collectors appear to provide representative samples. The National Atmospheric Deposition Program (NADP) recommends the Aerochem Metrics[®] 301 Collector for its nationwide network of wet deposition sites, and the device has been used widely in other studies of wet deposition^{23,5,35,25,42,21}. The wet bucket collects rainfall in amounts very close to those attained by independent rain gauges¹⁰⁸, and there are no cases of systematic bias in these samplers for rainfall water quality and quantity reported in the deposition literature. Sample preservation and holding time can be important issues for many water quality parameters. Total phosphorus should not be affected by these concerns for the usual weekly sampling routine.

Challenges remain, however, in using Aerochem Metrics[®], Inc., collectors for wet deposition. In active wildlife areas, there will always be some contaminated samples from biological debris that could dissolve in the collector and go undetected in field processing. Still, standard procedures for field sampling and detailed field notes can help reduce and identify unrepresentative samples. Careful selection of sites is important to avoid local contamination, and mounting collectors above the ground could help to avoid large local particles and minimize

biofouling (McDowell et al.¹⁰² recommend sampling at a 10-m height; however, 2 m is common). Also, procedures are needed for data screening to deal objectively and consistently with contaminated wet samples; Ahn and James²¹ and Walker⁴² provide examples of statistical approaches to screening deposition data mentioned earlier. When staff resources permit, short-term or manual deployment of samplers for rainfall events should avoid most contamination, but this is impractical for long-term studies.

Dry Deposition of P-Bearing Particles

Common methods for estimating the dry deposition of particles include three general approaches to measurement: micro-meteorological methods, surface accumulation methods and inferential techniques. Reviews by Hicks⁹¹, Nicholson¹⁰⁹, Erisman et al.⁸⁶, Hicks et al.¹², and Lovett³⁴ provide detailed comparisons of various methods for assessing gaseous, aerosol, and particulate deposition. The following summary is intended only to highlight key facts on each technique, with emphasis on particulate measurements applicable to particles conveying P, and with consideration of the potential for routine quantification for future environmental studies.

I. Micrometeorological methods have been successful in quantifying the submicron aerosol component of long-term ecosystem inputs for such pollutants as ozone, sulfate, methane, and ammonia^{30,34}. These methods can be applied *in situ*, provide continuous information, and do not disturb the sampling site. Micrometeorological methods use several approaches to estimate flux rates from meteorological data. Dry flux is estimated from detailed data on air concentrations above the target surface and various relationships with parameters associated with turbulent exchange processes^{91,30,86}. Baldocchi et al.³⁰ and Nicholson¹⁰⁹ give a summary of the fundamental theory for these approaches.

Unfortunately, these techniques as currently developed are not suitable for particles, even fine particles, above one micron in size^{91,86}. Highly sensitive rapid-response instruments are needed to quantify turbulence and directional concentrations. Sampling gradients and directional fluxes of larger, rapidly settling particles is extremely difficult or impossible¹⁰⁹. Even for fine aerosols, continuous measurement of air concentrations and meteorological parameters requires intense sampling and cutting-edge instrumentation. Documenting no more than a few percent of dry P deposition with current micrometeorological methods from very fine (submicron) particles cannot justify the investment needed for measurements. This conclusion might not hold true for cases in which the deposition of gaseous P must be measured. Micrometeorological methods are

scientifically appealing because they attempt to directly quantify atmospheric exchange processes, allowing estimates at the landscape level. They might become more useful if a measurable scalar, such as aerosol SO₂, can be determined to be applicable to important size fractions of particulate P in the air. Also, advances in instrumentation might make them more widely applicable to P-bearing supermicron particles and routine deposition monitoring.

II. Surface accumulation methods provide accumulation rates from mass changes directly on surfaces, natural or surrogate^{91,109,86}. These methods hold the greatest promise for long-term monitoring of P deposition. Surface-accumulation studies have proven more useful for coarse (2.5 to 10 μm) and giant particles (> 10 μm), whose deposition is governed by gravitational forces^{65,55,56}. Most sizes of atmospheric particles accumulate on downward- and upward-facing surfaces due to the concurrent effects of inertia and gravity³². As a result of this complexity, interpreting data for a particular surface area and collector shape is not straightforward. Combinations of scaled surrogate surface accumulation measurements might eventually be appropriate for environmental monitoring; calibration to estimate P deposition onto complex natural surfaces is needed.

Net Throughfall: Differences between bulk deposition in an open area and under a canopy estimate net dry deposition to a natural plant community in what is termed “net throughfall.” Throughfall measurements reflect the sum of wet, dry, cloud, and dew deposition onto natural plant communities^{34,86}. This technique has been used successfully to estimate net deposition fluxes to forest canopies^{110,90,76,111}. Vegetative processes (foliar leaching or uptake) can affect data interpretation, and P enrichment of precipitation moving through tree canopies has been documented^{112,113,114}. Such canopy processes are likely to confound throughfall P measurements, and this source of variation, plus the physical difficulty with *in situ* canopy and bulk measurements in wetland settings, make net throughfall an unlikely candidate for P deposition monitoring. If throughfall studies are found to be useful for P, their primary purpose will be to calibrate other approaches to long-term data collection for particular canopy types. Application to emergent and floating aquatic plant canopies has not been attempted.

Frisbee-shaped Collectors: To overcome aerodynamic interference with depositing particles, inverted Frisbee-shaped airfoils have been used for particle flux measurements^{115,116}. Net mass accumulation to flat surfaces inside the sampler is quantified over specified periods of time, and these Frisbee-shaped collectors have been used effectively in combination with other sampling devices in estimating dry deposition of a diverse suite of parameters^{57,33}. These aerodynamically shaped devices offer advantages for deposition monitoring, and they have been

used in the United Kingdom for dust monitoring¹¹⁷. Resulting data give reasonable estimates of deposition rates, as long as coated surfaces are used; these estimates are viewed as minimal rates to complex natural surfaces. Wind losses, changes in surface stickiness, and contamination events are potential drawbacks.

Flat Plate Sampler: Another device that has good aerodynamic qualities is the flat plate sampler with a wind vane developed by Noll et al.³². This plate sampler provides reproducible flux estimates to a flat, greased surface and is well suited to both chemical and numerical studies of particles⁶³. Using coatings in a dry collector increases capture efficiency and might provide better estimates for aquatic systems. In productive aquatic settings, however, either of these devices will still be subject to contamination by insects and birds. Field trials are needed to compare and calibrate methods for use in routine monitoring of P deposition, but the cited studies suggest that these collectors offer promise for routine deposition monitoring.

Integrated Throughfall, Surrogate Surface and Bulk Sampling: Long-term monitoring of chemical deposition to forests has been accomplished using artificial surfaces designed to simulate a canopy (Teflon string), with calibration by throughfall collections and bulk deposition¹¹⁸. It uses the method developed by Ferm and Hultberg¹¹⁹ for forest deposition monitoring in Sweden. The dry deposition of an ion is calculated by multiplying the ratio of sodium deposited on the string device to that in throughfall by the amount of the ion deposited on the string collector. This technique is attractive in that it assesses inputs at the canopy level without using electricity. With multiple sites, it is suitable for long-term data collection at the regional level to assess critical loads of pollutants. Variations of this integrated approach using various other artificial surfaces have been applied to forest deposition¹¹⁸.

This integrated method is cost effective once the artificial surface has been shown to collect a target chemical species; it has provided results for N and sulfur deposition comparable to other estimates. Determining the utility of the method for coarse and giant particles would require additional research. Contamination issues for bulk measurements and throughfall sampling remain significant uncertainties, particularly in productive aquatic environments. Covering the surrogate surface collector might help with sampler contamination, but its effect on particle capture would need to be studied. Even with these challenges, the concept of using a convenient surrogate collector calibrated by throughfall studies has potential for large-scale long-term monitoring of P deposition.

Bulk Funnels, Wet/Dry Buckets and Other Artificial Surfaces: Bulk and wet/dry

Aerochem Metrics[®], Inc. collectors are the most common monitoring devices for atmospheric deposition of P; many references are given in Tables 2 and 3. Reviews of deposition methods provide examples of other diverse surfaces that have been used to estimate bulk atmospheric inputs^{91,34,86}. As particle size increases, these collectors give better estimates of flux^{91,86}, although bulk measurements tend to underestimate inputs of ions conveyed by vapor or fine particles into forest canopies¹¹⁰. When contamination is not frequent and detectable, these devices are useful to provide a conservative estimate of deposition rates for coarse and giant particulate materials. Davidson et al.⁶⁵ and Lovett³⁴ provide a guide to the literature on attempts to use many other artificial surfaces in deposition studies, and they note that artificial surfaces might be the best available method to assess deposition rates of large particles.

Some inherent deficiencies exist in surrogate surface collectors, such as the common Aerochem Metrics[®], Inc. dustfall buckets for collecting quantitative samples of dry deposition. Particle resuspension and bounce-off are always possible for artificial surfaces. If exposed to the atmosphere for up to a week at a time, surrogate surfaces are subject to persistent contamination by bird and insect excrement and miscellaneous debris that, as discussed previously, may or may not be considered part of atmospheric deposition. Birds are a particular challenge to deposition measurements in marshes where any object used as a perch can be subject to extreme levels of contamination that could preclude interpretation of deposition rates from many samples, and even entire sampling sites^{39,40,42}. The geometry of dustfall buckets is also a problem, according to Hicks⁹¹ and Davidson et al.⁶⁵; dustfall buckets have a high surface:bottom area ratio of 4.8, compared to vessels with smaller sides, such as the common Petri dish, with a ratio of 1.8. This high surface:bottom ratio makes dry deposition data difficult to interpret as deposition per unit surface area of a canopy or the ground. In addition, bucket geometry creates an aerodynamic environment that decreases collection efficiency, particularly for smaller particles and under windy conditions¹⁰⁰. The distortion of airflow by the bucket rim causes particles to change their trajectories and escape capture by the bucket¹²⁰. As pointed out by Harrison and Perry⁵⁹, even larger particles tend to fall at very small angles and can easily be diverted away from surfaces by airflow characteristics of the bucket rim. Further, the chemical and physical nature of the surface can also be a factor in observed mass accumulations^{121,122}.

Together, these considerations make the interpretation of dry deposition data from dustfall buckets and bulk collectors difficult^{109,100,102}. The evidence reviewed in this study suggests these uncertainties are not large enough to preclude using bulk deposition and data from wet/dry collectors to bracket deposition rates for P to various land cover types (Tables 2 through 5).

Because these collectors provide more accurate estimates for deposition rates for substances, such as P, conveyed primarily by giant particles ($> 10 \mu\text{m}$), this general correspondence is expected. For study areas having infrequent contamination events, and where the need for a rough estimate of P deposition has been established, wet/dry collectors can still be considered a viable option until other routine methods are available. Adding water to the dry bucket seems reasonable to avoid resuspension and to get estimates closer to those experienced at the air-water interface^{72,5}.

Surface Accumulation Directly on Vegetation or Water Surfaces: Mass accumulation over a period of several days on leaf surface has been combined with surrogate samplers to estimate coarse particle deposition to forest canopies¹²³. Leaf surface provides a direct estimate of dry deposition but is subject to large uncertainties. Cloud/dew droplets can change rates. Foliar interactions with deposited P may be a substantial interference for estimates to a system with a plant canopy. Phosphorus accumulation on the surface of vegetation is probably not appropriate for routine monitoring of atmospheric deposition, but could be considered for validation and calibration purposes. Yi et al.¹²⁴ used a water surface sampler to measure dry deposition of sulfate. This surface sampler deserves further study, although it will be difficult to deploy in remote areas and will be subject to interference from strong winds and local contamination.

III. Inferential Techniques estimate deposition rates from air concentration data used in combination with direct particle enumeration or models to estimate size-specific deposition velocities⁹¹. Inferential techniques vary greatly in their level of detail and resolution. They can provide reliable estimates for aerosols, gases, and fine particles for which air concentrations can be measured. Deposition velocities can be estimated by models driven by meteorological data^{125,126,92}. Other variations on the inferential approach use direct particle enumeration and quantification of air concentrations for coarse and giant particles; these designs hold great promise as means of estimating dry deposition of particles conveying P into natural systems.

Many devices are available to collect fine and smaller coarse particles for compliance with ambient air standards³¹, and air concentration data can be used to yield crude deposition estimates when combined with average velocities. The difficulty is getting air concentration and deposition velocity data for specific size classes, particularly coarse and giant particle classes. Various impactor samplers allow quantification of the concentration of particles from below 1 to about $100 \mu\text{m}$ ^{63,56}. Combining rates of particle accumulation from a standardized surface, such as a flat plate sampler³², with air concentrations from impactor sampling can yield good

estimates of dry deposition for P, as has been done by Davidson et al.⁶⁶ and Lin et al.⁵⁶ for other components of atmospheric deposition.

Low Volume Filter Packs and the NOAA, DRY DEP 2 Model: Air concentrations of submicron particles and gases derived from filter pack samples and meteorological parameters are widely used to model deposition rates^{125,92}. Deposition models are dependent on multiple estimates of resistance, with many inherent assumptions. Models have been calibrated extensively for gaseous deposition^{126,125}; validation of modeled parameters remains a major uncertainty³⁴. The approach appears to produce conservative deposition estimates and is being used in the 's Dry Deposition Monitoring Program. While it is tempting to consider filter pack monitoring data for estimating P deposition, the method does not apply to deposition of supermicron particles. By design, filter packs do not quantify larger coarse and giant particles. Consequently, the P deposition rates estimated with filter pack data and deposition models by Yang et al.⁹⁶ and Hu et al.¹²⁷ were unrealistically low, representing less than 10 % of likely deposition rates. Overall, the approach will measure only a small fraction of dry P deposition, and it is not useful for monitoring total P inputs unless larger particles can be sampled and modeled more effectively.

Rotary Impactors and Flat Plates: Large volumes of air are sampled by a multistage impactor, and change in mass of each stage yields size-specific concentrations. The device is used in combination with flux measurement to flat plates to derive depositional velocities for particle size classes^{128,63,32}. The Noll Rotary Impactor provides a distribution of mass in 10 size classes up to about 100 μm and associated concentrations. A flat plate is used to estimate size-specific mass flux to a coated surface. Supermicron particle concentrations are quantified and, together with flux rates from plate sampling, data from this approach should provide mass flux for size classes that can be added to estimate total atmospheric P deposition. This sampling approach can be used to analyze for chemical tracers of particle sources and could be configured as upwind/downwind stations⁸⁹. Caffrey et al.⁵⁷ and Paode et al.⁶⁹ combine the Noll Rotary Impactor for air concentration data for coarse and giant particles and surrogate surface (Frisbee or flat plate) methods to estimate deposition velocities to derive dry deposition estimates. This approach has great potential for P deposition studies and routine monitoring.

BEYOND BULK PRECIPITATION AND WET/DRY DEPOSITION ESTIMATES

Lovett³⁴ was referring to many airborne nutrients and pollutants when he concluded: "A

major challenge for the future will be to reconcile differences among existing measurement methods for cloud and dry deposition, and to arrive at a suite of generally agreed-upon methods that are applicable to a broad range of sites and chemical substances.” His words are true today for atmospheric P deposition.

If the uncertainty in our current understanding of P deposition based largely on bulk and wet/dry collectors is too great for a particular circumstance, then a combination of the approaches discussed above is needed to improve estimates of dry deposition rates. Because the goal is typically to measure total P deposition to an ecosystem, there is currently no single technique that can give all the necessary information for determining and validating deposition rates for the size classes of particles conveying P over diel, seasonal and annual cycles. The only way to estimate total P deposition is by combining and comparing data from several surrogate surface and inferential approaches, at least until far more data are available to allow calibration and simplification of deposition sampling.

The South Florida Water Management District recently contracted with a group of experts in atmospheric science to investigate the science of P deposition relevant to the Florida Everglades. An important end product of this effort was a study design that would reduce uncertainty in rates and begin the process of developing a new generation of monitoring methods. The final, integrated product of their research was a work plan for a one-year field study to: (1) establish upper and lower bounds of dry P deposition; (2) inter-compare methods of estimation; and (3) calibrate fluxes to a vegetated Everglades canopy⁴⁵. The proposed work plan would lead to a more focused and longer-term study to refine methods into procedures suitable for long-term P monitoring.

The research plan of Pollman et al.⁴⁵ uses three complementary data streams, with *state of the science* methods (Figure 3). For one data stream, surrogate surface samplers (flat plates and Frisbee-shaped samplers) would provide flux estimates for various size classes of P to flat surfaces. These estimates are viewed as setting the lower bound for P deposition rates. For the second data stream, air concentrations for more than 12 different size classes of particles would be measured through impactor sampling (Micro-orifice Uniform Deposit Impactor and Noll Rotary Impactor). These impactor data would be compared with concurrent collections by several other common air sampling devices. Data on wind speed/direction, temperature, solar radiation, and humidity would be combined with aerosol concentration data in modeling deposition velocities and estimating dry deposition flux from the second data stream. To validate and calibrate rates derived from fluxes to surrogate surfaces and particle concentrations in the

air, the third component of the study would use throughfall measurements of P deposition to provide information on canopy scaling. Throughfall estimates in marsh plant communities would also be developed for a conservative tracer of P deposition, probably sodium. This element of the proposed study would effectively “ground-truth” throughfall estimates and, ultimately, validate rates from surrogate surfaces and aerosol concentrations and inferential models. Wet P deposition would be determined with Aerochem Metrics[®], Inc. wet/dry collectors. All three data streams of the study plan would be conducted at a primary study site, and flat plate and wet deposition data from a suite of satellite sites. Together the sampling methods and sites would provide detailed data to constrain estimates of particle fluxes in relation to local sources and meteorology.”

The Pollman et al.⁴⁵ study design dramatically illustrates that there is no simple or single means to improve upon extant P deposition data. The proposed study would require intense collaboration for one year of field sampling, moving well beyond most other studies involving multiple approaches to dry deposition that have been conducted for relatively brief periods of time^{55,57,33}. The proposed approach is also noteworthy in addressing uncertainty in P deposition in a holistic and balanced manner using methods at the cutting edge of deposition-related science. This work plan and the approach it represents will hopefully inspire further study of atmospheric P deposition to enable moving beyond the uncertainty of existing information.

CONCLUSIONS

This review has attempted to bridge the disparate disciplines of environmental and atmospheric science to improve interdisciplinary understanding of phosphorus deposition from the atmosphere. The study’s conclusions build upon and consolidate earlier synthetic work on P deposition to provide information supporting ecosystem management. By melding the information from the disciplinary literature referenced in this study, the author has come to the following conclusions and recommendations:

1. The vast majority of P in dry deposition is conveyed by coarse (2.5 to 10 μm) and giant (10 to 100 + μm) particles (Table 1). While fine particles (< 2.5 μm) also contain P, their minute mass and slow settling rate makes them responsible for only a few percent of total dry deposition. The roles of phosphine and other forms of gaseous P remain highly uncertain. The coarse and giant components of deposition have been quantified far less than those in the fine size range, because other priority airborne pollutants (sulfur, nitrogen, metals) tend to be conveyed in the fine fraction.

2. Processes operating across temporal and spatial scales at the land-air interface produce variation in P deposition from the atmosphere (Figure 2). Dynamics in particle sources lead to variation in particle concentrations and P content and are strongly associated with land use. Important sources of P in airborne particles are fugitive soils, biological debris and pollen, ash, combustion products of fossil fuels, emissions from manufacturing, and aerosols from the surface of marine and freshwater ecosystems. Natural plant canopies capture particles and lead to variation in deposition rates across the landscape mosaic.
3. The only substantial information base on P deposition rates is derived from bulk precipitation and wet/dry bucket collectors. These methods are subject to constraints in both precision and accuracy. A compilation of annual published rates (Tables 2 and 3) reveals that many freshwater aquatic ecosystems receive a mean deposition of about 20 and 80 mg Pm⁻²yr⁻¹. The overall range of published annual rates is from 3.6 to 200 mg Pm⁻²yr⁻¹. The median value for local rates is 30.1 mg Pm⁻²yr⁻¹ (Table 2), and for regional mean rates is 33.8 (Table 3).
4. P deposition occurs in measurable amounts, even in remote locations of the globe (Table 4), and P can be transported thousands of kilometers from sources in Asia and North Africa. The magnitudes of values given in Tables 4 and 5 suggest that the inherent limitations in bulk or wet/dry sampling do not preclude their use in estimating the upper bounds of annual P deposition from the atmosphere associated with various land uses.
5. There is no reason that the common wet/dry bucket sampling should not be used to sample wet deposition of P. Even with stringent quality assurance measures in the field and laboratory; however, contamination will continue to produce positive bias wet deposition estimates.
6. Improving the precision and accuracy of dry deposition rates will require an intercomparison of methods and an intense data collection effort. The likelihood of success is greatest using a comparative approach, such as that of Pollman et al.⁴⁵, in which P flux in deposition is estimated directly using surrogate surfaces, and inferentially using aerosol concentrations from impactor sampling and models to estimate deposition velocities. Flux estimates can be validated and calibrated using throughfall measurements to *ground truth* and scale rates to natural canopies⁴⁵ (Figure

- 3).
7. Contamination of samples in long-term monitoring will continue to cause some positive bias in estimates, even with field, laboratory, and statistical screening. As new methods are tested for P deposition sampling, reducing the frequency of contamination and increasing the likelihood of detecting unrepresentative samples must be primary considerations.
8. For South Florida's Everglades, information in this review leads to the conclusion that future sampling with wet/dry bucket collectors will not result in improved precision or accuracy in wet or dry P_t deposition estimates compared to those derived from existing data by Ahn and James²¹ and Walker⁴². The substantial investment in operating this network of stations cannot be justified by uncertain information to be gained by future monitoring in these wetlands.
9. The comparative study designed by Pollman et al.⁴⁵ could constrain deposition estimates from the wide range found in this review and lead to new ways of conducting long-term, cost-effective monitoring. Studies of this type are worth pursuing in a variety of ecosystems for the purposes of both management and research. With improved methods of estimating deposition over the long term, better information could be gathered at regional and national levels. Phosphorus deposition deserves assessment through large-scale atmospheric monitoring programs, such as the National Atmospheric Deposition Program (NADP), Atmospheric Integrated Research Monitoring Network (AIRMoN), and National Atmospheric Deposition Program-National Trends Network (NADP-NTN).

Greater precision and accuracy in deposition rate estimates are fundamentally important to the certainty of decision making in eutrophication management worldwide and are most needed for environmental management of oligotrophic systems. In South Florida, as surface water nutrient loading to the Florida Everglades is reduced through implementation of the Everglades Forever Act and the recently authorized Comprehensive Everglades Restoration Plan, atmospheric loading will retake its historical place as a primary source of nutrients to most areas of South Florida's "river of grass." Hopefully, the conceptual model, literature review and evaluation of methods presented above will contribute to an improved understanding of, and interest in, the transport and impact of airborne P in the Everglades and other aquatic ecosystems.

ACKNOWLEDGEMENTS

This study could not have been completed without the support of the South Florida Water Management District, West Palm Beach, Florida; I deeply appreciate this opportunity and challenge. I gratefully acknowledge the expert guidance provided by Drs. Tim Bechtel, Michael Chimney, Carl Fitz, Thomas Fontaine, Susan Gray, Karl Havens, R. Thomas James, Dave Rudnick, and Alan Steinman on the earlier versions of the manuscript. Many thanks to Carrie L. Trutwin and Felicia Berger, who used their technical editing and grammatical expertise to fine-tune this document. The study benefited greatly from a synthesis of information on P deposition conducted for our agency by Drs. Curtis Pollman, Kenneth Noll, William Landing, and Steven Lindberg. Rachelle Grein and Christopher King collaborated to create Figure 1. Christopher King provided graphics expertise for Figure 2.

REFERENCES

1. Gorham, E. (1961) Factors influencing supply of major ions to inland waters, with reference to the atmosphere. *Bull. Geol. Soc. Amer.* **72**, 795-840.
2. Schindler, D.W., Newbury, R.W., Beaty, K.G. and Campbell, P. (1976) Natural water and chemical budgets for a small Precambrian lake basin in Central Canada. *J. Fish. Res. Bd. Canada.* **33**, 2526-2543.
3. Lewis, W.M., Jr. (1983) Collection of airborne materials by a water surface. *Limnol. Oceanogr.* **28**, 1242-1246.
4. Psenner, R. (1984) The proportion of empneuston and total atmospheric inputs of carbon, nitrogen and phosphorus in the nutrient budget of a small mesotrophic lake (Piburger See, Austria). *Int. Revue Ges. Hydrobiol.* **69**, 23-39.
5. Jassby, A.D., Reuter, J.E., Axler, R.P., Goldman, C.R. and Hackley, S.H. (1994) Atmospheric deposition of nitrogen and phosphorus in the annual nutrient load of Lake Tahoe (California-Nevada). *Water Resources Res.* **30**, 2207-2216.
6. Banks, H.H. and Nighswander, J.E. (2000) Relative contribution of hemlock pollen to the phosphorus loading of the Clear Lake ecosystem near Minden, Ontario. In *U.S.D.A. Forest Service, General Technical Report GTR-NE-267*. McManus, K.A., Shields, K.S. and Souto, D.R., Eds. Proceedings: Symposium on Sustainable Management of Hemlock Ecosystems in Eastern North America, Delaware, OH, pp. 168-174.
7. Graham, W.F. and Duce, R.A. (1982) The atmospheric transport of phosphorus to the Western North Atlantic. *Atmosph. Environ.* **16**, 1089-1097.
8. Jickells, T. (1995) Atmospheric inputs of metals and nutrients to the oceans: Their magnitude and effects. *Marine Chem.* **48**, 199-214.
9. Guerzoni, S. and co-authors (1999) The role of atmospheric deposition in the biogeochemistry of the Mediterranean Sea. *Progress in Oceanogr.* **44**, 147-190.
10. Graham, W.F., Piotrowicz, S.R. and Duce, R.A. (1979) The sea as a source of atmospheric phosphorus. *Marine Chem.* **7**, 325-342.
11. United States Environmental Protection Agency. (1994) Deposition of Air Pollutants to the Great Waters. *First Report to Congress*. Office of Air Quality, U.S. Environmental Protection Agency, Research Triangle Park, N.C. EPA-453/R-93-055, 89 pp.
12. Hicks, B., McMillen, R., Turner, R.S., Holdren, G.R. and Strickland, T.C. (1993) A national critical loads framework for atmospheric deposition effects assessment: III. Deposition characterization. *Environ. Management.* **17**, 343-353.
13. National Atmospheric Deposition Program. (1996) *NADP/NTN wet deposition in the United States, 1995*. National Atmospheric Deposition Program, Colorado State University, Fort Collins, CO.

14. United States Environmental Protection Agency. (1996) *Air Quality Criteria for Particulate Matter, Vol. I, Chapter 6*. Office of Research and Development, Washington, D.C. EPA/600/P-95/001aF.
15. Puckett, L.J. (1995) Identifying the major sources of nutrient water pollution. *Environ. Sci. Tech.* **29**, 408-414.
16. Hecky, R.E. and Kilham, P. (1988) Nutrient limitation of phytoplankton in freshwater and marine environments: A review of recent evidence on the effects of enrichment. *Limnol. Oceanogr.* **33**, 796-822.
17. Smith, V.H. (1990) Phytoplankton responses to eutrophication in inland waters. In *Introduction to Applied Phycology*. Akatsuka, I., Ed. SPB Academic Publishing. The Hague, Netherlands, pp. 231-249.
18. Welch, E.B. and Cooke, G.D. (1987) Lakes. In *Restoration Ecology – a synthetic approach to ecological research*. Jordan, W.R., Gilpin, M.E. and Aber, J.D., Eds. Cambridge University Press, Cambridge, MA, pp. 109-129.
19. Smith, V.H., Tilman, G.D. and Nekola, J.C. (1999) Eutrophication: Impacts of excess nutrients on freshwater, marine and terrestrial ecosystems. *Environ. Poll.* **100**, 179-196.
20. Davis, S.M. (1994) Phosphorus inputs and vegetation sensitivity in the Everglades. In *Everglades – The Ecosystem and its Restoration*. Davis, S.M and Ogden, J.C., Eds. St. Lucie Press, Lantana, FL, pp. 357-378.
21. Ahn, H. and James, R.T. (2001) Variability, uncertainty, and sensitivity of phosphorus deposition load estimates in South Florida. *Water, Air, and Soil Poll.* **126**, 37-51.
22. Irwin, G.A. and Kirkland, R.T. (1980) *Chemical and physical characteristics of precipitation at selected sites in Florida*. U.S. Geological Survey, Water Resources Investigations, 80-81, 70 pp.
23. Hendry, C.D., Brezonik, P.L. and Edgerton, E.S. (1981) Atmospheric deposition of nitrogen and phosphorus in Florida. In *Atmospheric Pollutants in Natural Waters*. Eisenreich, S.J., Ed. Ann Arbor Science, Ann Arbor, MI, pp. 199-215.
24. Dixon, L.K., Murray, S., Perry, J.S., Minotti, P.J., Henry, M.S. and Pierce, R.H. (1996) Assessment of Bulk Atmospheric Deposition to the Tampa Bay Watershed. *Tampa Bay National Estuary Program Technical Publication #08-96*, September 1996, by Mote Marine Laboratory, Sarasota, FL, 232 pp.
25. Peters, N.E. and Reese, R.S. (1995) Variations of weekly atmospheric deposition for multiple collectors at a site on the shore of Lake Okeechobee, Florida. *Atmosph. Environ.* **29**, 179-187.
26. Graham, W.F. and Duce, R.A. (1979) Atmospheric pathways of the phosphorus cycle. *Geochimica et Cosmochimica Acta.* **43**, 1195-1208.

27. Newman, E.I. (1995) Phosphorus inputs to terrestrial ecosystems. *J. of Ecology*. **83**, 713-726.
28. Likens, G.E. and Bormann, F.H. (1995) *Biogeochemistry of a forested ecosystem. 2nd edition*. Springer Verlag, New York, 159 pp.
29. Gibson, C.E., Wu, Y. and Pinkerton, D. (1995) Substance budgets of an upland catchment: The significance of atmospheric inputs. *Freshwater Biol.* **33**, 385-392.
30. Baldocchi, D.D., Hicks, B.B. and Meyers, T.P. (1988) Measuring biosphere-atmosphere exchanges of biologically related gases with micrometeorological methods. *Ecology*. **69**, 1331-1340.
31. Chow, J.C. (1995) Measurement methods to determine compliance with ambient air quality standards for suspended particles. *J. of Air and Waste Management Assoc.* **45**, 320-381.
32. Noll, K.E., Fang, K.Y.P. and Watkins, L.A. (1988) Characterization of the deposition of particles from the atmosphere to a flat plate. *Atmosph. Environ.* **22**, 1461-1468.
33. Zufall, M.J., Davidson, C.I., Caffrey, P.F. and Ondov, J. (1998) Airborne concentrations and dry deposition fluxes of particulate species to surrogate surfaces deployed in Lake Michigan. *Environ. Sci. and Techn.* **32**, 1623-1628.
34. Lovett, G.M. (1994) Atmospheric deposition of nutrients and pollutants in North America: An ecological perspective. *Ecological Applications*. **4**, 629-650.
35. Jordan, T.E., Correll, D.L. Weller, D.E. and Goff, N.M. (1995) Temporal variation in precipitation chemistry on the shore of the Chesapeake Bay. *Water, Air, and Soil Poll.* **83**, 263-284.
36. Wu, Z.Y., Han, M., Lin, Z.C., and Ondov, J.M. (1994) Chesapeake Bay atmospheric deposition study, Year 1: Sources and dry deposition of selected elements in aerosol particles. *Atmosph. Environ.* **28**, 1471-1486.
37. Davidson, C.I., Miller, J.M. and Pleskow, M.A. (1982) The influence of surface structure on predicted particle dry deposition to natural grass canopies. *Water, Air, and Soil Poll.* **18**, 25-43.
38. Ahn, H. (1999a) Statistical modeling of total phosphorus concentrations measured in South Florida rainfall. *Ecological Modeling*. **116**, 33-44.
39. Ahn, H. (1999b) Outlier detection in total phosphorus concentration data from South Florida rainfall. *J. Amer. Water Resources Assoc.* **35**, 301-310.
40. Ahn, H. and James, R.T. (1999) Outlier detection in phosphorus dry deposition rates measured in South Florida. *Atmosph. Environ.* **33**, 5123-5131.
41. Ahn, H. and James, R.T. (2000) Statistical modeling of phosphorus dry deposition rates measured in South Florida. *Water, Air and Soil Poll.* **121**, 43-57.

42. Walker, W.W. (1999) *Analysis of water quality monitoring data from Arthur R. Marshall Loxahatchee National Wildlife Refuge*. Prepared for the Department of the Interior by William W. Walker, Environmental Engineer, Concord, MA. Available at: <http://www2.primushost.com/~wwwwalker/>.
43. van Ek, R. and Draaijers, G.P.J. (1994) Estimates of atmospheric deposition and canopy exchange for three common tree species in the Netherlands. *Water, Air, and Soil Poll.* **73**, 61-82.
44. McCormick, P.V., Rawlik, P.S., Lurding, K., Smith, E.P. and Sklar, F.H. (1996) Periphyton-water quality relationships along a nutrient gradient in the Northern Florida Everglades. *J. N. Amer. Benthol. Soc.* **15**, 433-449.
45. Pollman, C., Landing, W., Lindberg, S., Mulholland, P. and Noll, K. (2000) *Atmospheric deposition of phosphorus in the Everglades: Proof of concepts and methods. Work Plan of Phase II Studies*. Report to South Florida Water Management District, West Palm Beach, FL., June 5, 2000, 38 pp.
46. Gordon, G.E. (1991) Airborne particles on regional and global scales. *Environ. Sci. and Tech.* **25**, 1822-1828.
47. Gardner, L.R. (1990) The role of rock weathering in the phosphorus budget of terrestrial watersheds. *Biogeochemistry.* **11**, 97-110.
48. South Florida Water Management District (2001) *2001 Everglades Consolidated Report, Chapters 2 and 3*, January 1, 2001. South Florida Water Management District, West Palm Beach, FL. Available at: www.sfwmd.gov/org/ema/everglades.
49. Glindeman, D., Stottmeister, U. and Bergmann, A. (1996) Free phosphine from the anaerobic biosphere. *Environ. Sci. and Poll. Res.* **3**, 17-19.
50. Devai, I., Felfody, L., Wittner, I. and Plosz, S. (1988) Detection of phosphine: New aspects of the phosphorus cycle in the hydrosphere. *Nature.* **333**, 343-345.
51. Devai, I. and Delaune, R.D. (1995) Evidence for phosphine production and emission from Louisiana and Florida marsh soils. *Org. Chem.* **23**, 277-279.
52. Lawson, D.R. and Winchester, J.W. (1979) Sulfur, potassium and phosphorus associations in aerosols from South American tropical rain forests. *J. Geophys. Res.* **84**, 3723-3727.
53. Casareto, B.E., Suzuki, Y., Okada, K. and Morita, M. (1996) Biological microparticles in rainwater. *Geophysical Research Letters.* **23**, 173-176.
54. United States Environmental Protection Agency (1996) *Air Quality Criteria for Particulate Matter, Vol. 1, Chapter 5*. Office of Research and Development, Washington, D.C. EPA/600/P-95/001aF.
55. Holsen, T.M., Noll, K.E., Fang, G.C., Lee, W.J., Lin J.M. and Keeler, G.J. (1993) Dry deposition and particle size distributions measured during the Lake Michigan urban air toxics study. *Environ. Sci. and Tech.* **27**, 1327-1333.

56. Lin, J.J., Noll, K.E. and Holsen, T.M. (1994) Dry deposition velocities as a function of particle size in the ambient atmosphere. *Aerosol Sci. and Tech.* **20**, 239-252.
57. Caffrey, P.F., Ondov, J.M., Zufall, M.J. and Davidson, C.I. (1998) Determination of size-dependent dry particle deposition velocities with multiple intrinsic elemental tracers. *Environ. Sci. and Tech.* **32**, 1615-1622.
58. Finlayson-Pitts, B.J. and Pitts, J.N. (1986) *Atmospheric Chemistry: Fundamentals and Experimental Techniques*. John Wiley and Sons, New York.
59. Harrison, R.M. and Perry, R. (1986) *Handbook of Air Pollution Analysis, Chapter 4 (2nd Edition)*. Chapman and Hall, Ltd., London.
60. Waser, N.A.D. and Bacon, M.P. (1995) Wet deposition fluxes of cosmogenic ³²P and ³³P and variations in the ³³P/³²P ratios at Bermuda. *Earth & Planetary Science Letters*. **133**, 71-80.
61. Ellis, W.G. and Merrill, J.T. (1995) Trajectories for Saharan dust transported to Barbados using Stokes' Law to describe gravitational settling. *J. of Applied Meteorology*. **34**, 1716-1726.
62. Betzer, P.G., and 11 co-authors (1988) Long-range transport of giant mineral aerosol particles. *Nature*. **336**, 568-571.
63. Noll, K.E. and Fang, K.Y.P. (1989) Development of a dry deposition model for atmospheric coarse particles. *Atmosph. Environ.* **23**, 585-594.
64. Chen, L., Arimoto, R. and Duce, R.A. (1985) The sources and forms of phosphorus in marine aerosol particles and rain from Northern New Zealand. *Atmosph. Environ.* **19**, 779-787.
65. Davidson, C.I., Lindberg, S.E., Schmidt, J.A., Cartwright, L.G. and Landis, L.R. (1985a) Dry deposition of sulfate onto surrogate surfaces. *J. of Geophysical Res.* **90**, 2123-2130.
66. Davidson, C.I., Goold, W.D., Mathison, T.P., Wiersma, G.B., Brown, K.W. and Reilly, M.T. (1985b) Airborne trace elements in the Great Smoky Mountains, Olympic and Glacier National Parks. *Environ. Sci. and Tech.* **19**, 27-35.
67. Brook, J.R., Dann, T.F. and Burnett, R.T. (1997) The relationship among TSP, PM_{2.5}, PM₁₀ and inorganic constituents of atmospheric particulate matter at multiple Canadian locations. *J. of Air and Waste Management Assoc.* **47**, 2-19.
68. Lundgren, D.A., Hausknecht, B.J. and Burton, R.M. (1984) Large particle size distribution in five U.S. cities and the effect on a new ambient particulate matter (PM) standard. *Aerosol Sci. and Tech.* **7**, 467-473.
69. Paode, R., Sofuoglu, S.C., Sivadechathep, J., Noll, Kenneth E., Holsen, T.M. and Keeler, G. (1998) Dry deposition fluxes and mass size distribution of Pb, Cu, and Zn measured in Southern Lake Michigan during AEOLOS. *Environ. Sci. and Tech.* **32**, 1629-1635.

70. Gasith, A. and Hasler, A.D. (1976) Airborne litterfall as a source of organic matter in lakes. *Limnol. Oceanogr.* **21**, 253-258.
71. Likens, G.E. (1985) The aquatic ecosystem and air-land-water interactions. In *An Ecosystem Approach to Aquatic Ecology*. Likens, G.E., Ed. Springer Verlag, New York, pp. 430-435.
72. Cole, J.J., Caraco, N.F. and Likens, G.E. (1990) Short-range atmospheric transport: A significant source of phosphorus to an oligotrophic lake. *Limnol. Oceanogr.* **35**, 1230-1237.
73. Rau, G.H. (1976) Dispersal of terrestrial plant litter into a subalpine lake. *OIKOS*. **27**, 153-160.
74. Psenner, R. (1984) The proportion of empenuston and total atmospheric inputs of carbon, nitrogen and phosphorus in the nutrient budget of a small mesotrophic lake (Piburger See, Austria). *Int. Revue Ges. Hydrobiol.* **69**, 23-39.
75. Anttila, P., Paatero, P., Tapper, U. and Jarvinen, O. (1995) Source identification of bulk wet deposition in Finland by positive matrix factorization. *Atmosph. Environ.* **29**, 1705-1718.
76. Brown, A.D., and Lund, L.J. (1994) Factors controlling throughfall characteristics at a high elevation Sierra Nevada site, California. *J. Environ. Qual.* **23**, 844-850.
77. Grant, M.C. and Lewis, W.M. (1982) Chemical loading rates from precipitation in the Colorado Rockies. *Tellus*. **34**, 74-88.
78. Takimoto, K., Hirakawa, T., Ito, K., Mukai, T. and Okada, M. (1999) Source and transport of tricresyl phosphate (TCP) isomers in the Kurose River Basin. *Atmosph. Environ.* **33**, 3191-3200.
79. Prospero, J.M., Barrett, K., Church, T., Dentener, F., Duce, R.A., Galloway, J.N., Levy, H. II, Moody, J. and Quinn, P. (1996) Atmospheric deposition of nutrients to the North Atlantic Basin. *Biogeochemistry*. **35**, 27-73.
80. Avila, A., Alarcon, M. and Queralt, I. (1998) The chemical composition of dust transported in red rains – its contribution to the biogeochemical cycle of a Holm Oak forest in Catalonia (Spain). *Atmosph. Environ.* **32**, 179-191.
81. Watson, J.G., Chow, J.C., Lowenthal, D.H., Pritchett, L.C., Frazier, C.A., Neuroth, G.R. and Robbins, R. (1994) Differences in the carbon composition of source profiles for diesel- and gasoline-powered vehicles. *Atmosph. Environ.* **28**, 2493-2505.
82. Olmez, I., Sheffield, A.E., Gordon, G.E., Houck, J.E., Pritchett, L.C., Cooper, J.A., Dzubay, T.G. and Bennett, R.L. (1988) Compositions of particles from selected sources in Philadelphia for receptor modeling applications. *J. Air Poll. Control Assoc.* **38**, 1392-1402.
83. Zufall, M.J., Weiping, D., Davidson, C.I. and Etyemezian, V. (1999a) Dry deposition of particles to wave surfaces: I. Mathematical modeling. *Atmosph. Environ.* **33**, 4273-4281.

84. Zufall, M.J., Weiping, D. and Davidson, C.I. (1999b) Dry deposition of particles to wave surfaces: II. Wind tunnel experiments. *Atmosph. Environ.* **33**, 4283-4290.
85. Lindberg, S.E. and Lovett, G.M. (1983) Application of surrogate surface and leaf extraction methods to estimation of dry deposition to plant canopies. In *Precipitation Scavenging, Dry Deposition and Resuspension. Vol. 2, Dry Deposition and Resuspension*. Pruppacher, H.R., Semonin, R.G. and Slinn, W.G.N., Eds. Elsevier Sciences Publishing Co., New York, pp. 837-846.
86. Erisman, J.W., Beier, C., Draaijers, G. and Lindberg, S. (1994) Review of deposition monitoring methods. *Tellus*. **46B**, 79-93.
87. Weathers, K.C., Lovett, G.M. and Likens, G.E. (1995) Cloud deposition to a spruce forest edge. *Atmosph. Environ.* **29**, 665-672.
88. Richerson, P.J., Moshiri, G.A. and Godshalk, G.L. (1970) Certain ecological aspects of pollen dispersion in Lake Tahoe (California-Nevada). *Limnol. Oceanogr.* **15**, 149-153.
89. Delumyea, R. and Petel, R.L. (1979) Deposition velocity of phosphorus-containing particles over Southern Lake Huron, April - October 1975. *Atmosph. Environ.* **13**, 287-294.
90. Lindberg, S.E., Lovett, G.M., Schaefer, D.A. and Bredemeier, M. (1988) Dry deposition velocities and surface-to-canopy scaling factors for aerosol calcium from forest canopy throughfall. *J. Aerosol Sci.* **19**, 1187-1190.
91. Hicks, B.B. (1986) Measuring dry deposition: A reassessment of the state of the art. *Water, Air, and Soil Poll.* **30**, 75-90.
92. Pratt, G.C., Orr, E.J., Bock, D.C., Strassman, R.L., Fundine, D.W., Twaroski, C.J., Thornton, J.D. and Meyers, T.P. (1996) Estimation of dry deposition of inorganics using filter pack data and inferred deposition velocity. *Environ. Sci. and Tech.* **30**, 2168-2177.
93. Eisenreich, S.J., Emmling, P.J. and Beeton, A.M. (1977) Atmospheric loading of phosphorus and other chemicals to Lake Michigan. *J. Great Lakes Res.* **3**, 291-304.
94. Shaw, R.D., Trimbee, A.M., Minty, A., Fricker, H. and Prepas, E.E. (1989) Atmospheric deposition of phosphorus and nitrogen in Central Alberta with emphasis on Narrow Lake. *Water, Air, and Soil Poll.* **43**, 119-134.
95. Kopacek, J., Prochazkova, L., Hejzlar, J. and Blazka, P. (1997) Trends and seasonal patterns of bulk deposition of nutrients in the Czech Republic. *Atmosph. Environ.* **31**, 797-808.
96. Yang, X, Miller, D.R., Xu, X., Yang, L.H., Chen, H.M. and Nikolaidis, N.P. (1996) Spatial and temporal variations of atmospheric deposition in interior and coastal Connecticut. *Atmosph. Environ.* **30**, 3801-3810.

97. Bergametti, G., Remoudaki, E., Losno, R., Sterner, E. and Chatenet, B. (1992) Source, transport and deposition of atmospheric phosphorus over the northwestern Mediterranean. *J. of Atmosph. Chemistry*. **14**, 501-513.
98. Steen, B. (1986) Dry deposition of 4- to 50- μ m dolomite particles on vegetation, flat surfaces and deposition gauges. *Atmosph. Environ.* **20**, 1597-1604.
99. Ahl, T. (1980) Eutrophication in relation to the load of pollution. *Progress in Water Technol.* **12**, 49-61.
100. Lodge, J.P. (1989) Method 502. *Methods of Air Sampling and Analysis, Third Edition*. Lodge, J.P., Ed. Lewis Publishers, Chelsea, MI.
101. Guerzoni, S. and co-authors. (1999) The role of atmospheric deposition in the biogeochemistry of the Mediterranean Sea. *Progress in Oceanogr.* **44**, 147-190.
102. McDowell, W.H., Gatz, D.F., Ondov, J.M., Pollman, C.J. and Winchester, J.W. (1997) *Atmospheric deposition into South Florida: Advisory panel final report*. South Florida Water Management District, P.O. Box 24680, West Palm Beach, FL, 33416-4680. December 24, 1997, 37 pp.
103. Walker, W.W. (1995) Design basis for Everglades Stormwater Treatment Areas. *Water Resources Bull.* **31**, 671-685.
104. Fitz, H.C. and Sklar, F.H. (1999) Ecosystem analysis of phosphorus impacts and altered hydrology in the Everglades: A landscape modeling approach. In *Phosphorus Biogeochemistry in Subtropical Ecosystems*. Reddy, K.R., O'Connor, G.A. and Schelske, C.L., Eds. Lewis Publishers, Boca Raton, FL, pp. 585-620.
105. Chadwick, O.A., Derry, L.A., Vitousek, P.M., Huebert, B.J. and Hedin, L.O. (1999) Changing sources of nutrients during four million years of ecosystem development. *Nature*. **397**, 491-497.
106. Craft, C.B. and Richardson, C.J. (1993) Peat accretion and phosphorus accumulation along a eutrophication gradient in the northern Everglades. *Biogeochemistry*. **22**, 133-156.
107. Reddy, K.R., DeLaune, R.D., DeBusk, W.F. and Koch, M.S. (1993) Long-term nutrient accumulation rates in the Everglades. *J. Soil Science Society of Amer.* **57**, 1147-1155.
108. South Florida Water Management District, (unpubl. data).
109. Nicholson, K.W. (1988) The dry deposition of small particles: A review of experimental measurements. *Atmosph. Environ.* **22**, 2653-2666.
110. Lindberg, S.E., Lovett, G.M., Richter, D.D. and Johnson, D.W. (1986) Atmospheric deposition and canopy interactions of major ions in a forest. *Science*. **231**, 141-145.
111. Butler, T.J. and Likens, G.E. (1995) A direct comparison of throughfall plus stemflow to estimates of dry and total deposition of sulfur and nitrogen. *Atmosph. Environ.* **29**, 1253-1265.

112. Li, Y.C., Alva, A.K., Calvert, D.V. and Zhang, M. (1997) Chemical composition of throughfall and stemflow from citrus canopies. *J. of Plant Nutrition*. **20**, 1351-1360.
113. McDowell, W.H. (1998) Internal nutrient fluxes in a Puerto Rican rain forest. *J. of Trop. Ecology*. **14**, 521-536.
114. Clark, K.L., Nadkarni, N.M., Schaefer, D. and Gholz, H.L. (1998) Atmospheric deposition and net retention of ions by the canopy in a tropical montane forest, Monteverde, Costa Rica. *J. of Trop. Ecology*. **14**, 27-45.
115. Davidson, C.I. and Wu, Y.L. (1990) Dry deposition of particles and vapors. In *Acidic Precipitation, Volume 3. Sources, deposition and canopy interactions*. Lindberg, S.E., Page, A.L. and Norton, S.A., Eds. Springer-Verlag, New York, pp. 103-216.
116. Hall, D.J., Upton, S.L. and Marsland, G.W. (1994) Designs for a deposition gauge and a flux gauge for monitoring ambient dust. *Atmosph. Environ.* **28**, 2963-2979.
117. Vallack, H.W. (1995) A field evaluation of Frisbee-type dust deposit gauges. *Atmosph. Environ.* **29**, 1465-1469.
118. Ferm, M. and Hultberg, H. (1999) Dry deposition and internal circulation of nitrogen, sulphur and base cations to a coniferous forest. *Atmosph. Environ.* **33**, 4421-4430.
119. Ferm, M. and Hultberg, H. (1995) Method to estimate atmospheric deposition of base cations in coniferous throughfall. *Water, Air, and Soil Poll.* **85**, 2229-2234.
120. Dombrowski, N., Fourny, E.A., Ingham, D.B. and Qi, Y.D. (1995) Design of wind-independent deposition gauges. *Atmosph. Environ.* **29**, 767-779.
121. Dasch, J.M. (1983) A comparison of surrogate surfaces for dry deposition monitoring. In *Precipitation Scavenging, Dry Deposition and Resuspension*. Semonin, R.G., Pruppacher, H.R. and Slinn, W.G., Eds. Elsevier Pubs., New York, pp. 883-899.
122. Dasch, J.M. (1985) Direct measurement of dry deposition to a polyethylene bucket and various surrogate surfaces. *Environ. Sci. and Tech.* **19**, 721-725.
123. Lindberg, S.E. and Lovett, G.M. (1985) Field measurements of particle dry deposition rates to foliage and inert surfaces in a forest canopy. *Environ. Sci. and Tech.* **19**, 238-244.
124. Yi, S.M., Holsen, T.M. and Noll, K.E. (1997) Comparison of dry deposition predicted from models and measured with a water surface sampler. *Environ. Sci. and Tech.* **31**, 272-278.
125. Meyers, T.P. and Baldocchi, D.D. (1988) A comparison of methods for deriving dry deposition fluxes of O_3 and SO_2 to a forest canopy. *Tellus*. **40B**, 270-284.

126. Hicks, B.B., Baldocchi, D.D., Meyers, T.P., Hosker, R.P. and Matt, D.R. (1987) A preliminary multiple resistance routine for deriving dry deposition velocities from measured quantities. *Water, Air, and Soil Poll.* **36**, 311-330.
127. Hu, H.L., Chen, H.M., Nikolaidis, N.P., Miller, D.R. and Yang, X. (1998) Estimation of nutrient atmospheric deposition to Long Island Sound. *Water, Air, and Soil Poll.* **105**, 521-538.
128. Noll, K.E. and Fang, K. (1986) *A rotary impactor for size selective sampling of atmospheric coarse particles.* 79th Annual Meeting, Air Pollution Control Association, Minneapolis, MN, June 22-27, 22 pp.
129. Malueg et al. (1975) Table 2.
130. Campo et al. (2001) Table 2.
131. Brown et al. (1984) Table 2.
132. Reid and Dillon (1994) Table 2.
133. Swap et al. (1992) Table 4.
134. Ahl, T. (1988) Background yield of phosphorus from drainage area and atmosphere: An empirical approach. *Hydrobiologia.* **170**, 35-44, Tables 3 and 4.
135. Gough et al. (2001) Data from Tables 10 and 11.
136. Dolske and Sievering (1980) (Table 6).

FIGURE CAPTIONS

Figure 1. Major geographic features of South Florida. The Everglades Protection Area encompasses the remnant Florida Everglades and is comprised of three Water Conservation Areas and Everglades National Park. Water resources in the region are the responsibility of the South Florida Water Management District, West Palm Beach, Florida.

Figure 2. A conceptual model of the transport and deposition of phosphorus from the atmosphere. The landscape of South Florida (Figure 1) is used for illustration, although sources, processes, and concepts can be applied to any landscape.

Figure 3. A diagrammatic representation of the experimental plan proposed by Pollman et al. (2000) to improve estimates of P deposition. The plan features three data streams leading to three comparable sets of flux estimates. This figure is a major revision of Figure 3 from Pollman et al.⁴⁵, and the author is solely responsible for its content.

TABLE LEGENDS

Table 1. The characteristics of particle size categories conveying atmospheric P. The information in this table was drawn from various sources in the literature, particularly Finlayson-Pitts and Pitts⁵⁸, Noll and Fang⁶³, Hicks et al.¹², Lovett³⁴, Chow³¹ and United States Environmental Protection Agency⁵⁴.

Table 2. A summary of annual rates of P deposition from the atmosphere. Deposition values are from published data using bulk or wet/dry collectors for periods from about one to over twenty years. These examples are based on data from either multiple years at a site or multiple collectors for at least one year to illustrate typical rates for various locations on an annual basis.

Table 3. A compilation of average rates of atmospheric P deposition from published literature summaries on atmospheric deposition. Each mean value in this table represents multiple years of data summarized by the authors and can be considered as a representative regional or continental average for deposition rates within the last three decades.

Table 4. A summary of rates of atmospheric P deposition from the scientific literature for remote areas of the globe. Each value in this table represents an annual estimate using a variety of methods as described. These values can be viewed as minimal, long-distance, long-term P deposition rates from the atmosphere.

Table 5. A summary of rates P deposition estimated using methods other than bulk deposition or wet/dry bucket collections. Most estimates are derived from variations of the inferential technique using air concentration data and deposition velocities. Note that

several of the rates for P are conversions from total particle mass using a value of P mass = 0.001 total mass, a conservative average for crustal materials.

Characteristic	Particle Size Category*		
	Fine Particles ($< 2.5 \mu\text{m}$)**	Coarse Particles ($2.5 - 10 \mu\text{m}$)**	Giant Particles ($10 - 100+ \mu\text{m}$)
Key Attributes for P Deposition			
Scale of Transport	Often global/regional; long atmospheric suspension possible	Often regional; inter-continental transport of dusts well documented	Often regional/local; global episodically; mechanisms of suspension unclear
Common Sources	Combustion, air transformations, organic compounds, microbes	Fugitive dusts, biotic processes, sea spray, volcanic ash	Soil, coal, vehicular and industrial dusts, organic debris, sea spray
Land-use Association	Low density; low landscape disturbance	Urban, industrial or agriculture	Dependent on local groundcover and biological communities
P Content	High, emission sources vary (0.1 - 1.0 % P)	Crustal levels likely (0.1 - 0.2 % P)	Variable based on particle source
Transport and Deposition Mechanisms	Follow local air eddies; Brownian diffusion and surface adhesion can be important	Less dependent on eddies; interception and inertial effects important	Gravitational and density effects dominate settling and surface adhesion
Deposition Velocity	Low, ca. $0.01 - 0.1 \text{cm s}^{-1}$	Moderate (ca. $0.1 - 1.0 \text{cm s}^{-1}$)	High (ca. $1 - > 10 \text{cm s}^{-1}$); density and shape effects possible
Surface Effects	Surface fine structure and wetness are important	Variable effects, wetness important, inertia involved	Inertial and gravitational forces dominate
Canopy Effects	Important as sink; biochemical release uncertain source; leaf area alters deposition	Variable; resuspension, creation and bounce-off possible	Important as source, esp. biotic particles; canopy structure less important
General linkage to P Deposition Rate	Overall, minor contributor to P flux; may support back-ground deposition	Important component of total P flux; diel, seasonal and climatic cycles	Dominates P flux; diel, seasonal and episodic variation; desert storms

* Size of groups of particles is usually expressed as aerodynamic diameter; the diameter of a sphere with a settling velocity at a density of 1g cm^{-3} equal to that being considered.

** The fine and coarse size categories follow those developed for particulate matter monitoring and assessment by the U.S. Environmental Protection Agency³⁴. The giant particle category is not relevant for air pollution assessment based on human health considerations and cannot be measured by most high-volume compliance samplers that have a cutoff point of about $10 \mu\text{m}$.

Table 1

P Deposition Rate (mgPm ⁻² yr ⁻¹) Mean, Range, Sampling Period	N Deposition Rate (mgNm ⁻² yr ⁻¹)	Sampling Location and Cover Type	Notes on Sampling Design and Techniques	Sources
12.4 8.8 - 17.6 6 years	795	Shagawa Lake, Minnesota; forested, small- town setting	Bulk precipitation samples at a single stations in the city of Ely, Minn.; Bulk concentrations X monthly precipitation on lake	Malueg, et al. ¹²⁹ (1975)
14. 6 - 22 5 years	490	Northern inland, various forested sites	Bulk deposition (assumed), eight stations, five-year period, 1971-1976	Ahl ⁹⁹ (Table 1)
15. 12 - 18 15 years	1020	Slapy Reservoir, Czech Republic; forested	Bulk collector, one station, sampled after precipitation events (site I), 4m above maximum water level	Kopacek et al. ⁹⁵ (1997)
15.6 6.9 - 22.8 6 years	None	Chamela tropical dry forest, Mexico	Bulk collectors with funnels, six stations in cleared areas, glass wool in funnels, storm event sampling	Campo et al. ¹³⁰ (2001)
19. None 1 year	None	Coastal shrub habitat, South Africa	Five replicate collectors with 1mm screens, 1.5m above soil surface, weekly sampling interval; local soils very low in phosphorus	Brown et al. ¹³¹ (1984)
20.3 17.8 - 23.7 3 years	424	Narrow Lake, Alberta, Canada; forested	Three-to-five bulk collectors, sampled after events and dry periods, funnels on collectors	Shaw et al. ⁹⁴ (1989)
23.8 11.1 - 74.9 16 years	1103	Lakes in Central Ontario, Canada; forested	Various sites, 1976-1992, bulk collections with window screens, sampling median 8 days	Reid and Dillon ¹³² (1994)
24. 12 - 31 13 years	1380	Rimov Reservoir, Czech Republic; farmland, village	Bulk collector, single station, sampled after precipitation events (site II), 14 m above maximum water level	Kopacek et al. ⁹⁵ (1997)
27.7 16.9 - 55.8 (1.5 years)	None	Lake Michigan area; urban settings	Data from ref. Figure 2; 15 sites sampled monthly for 18 months, bulk collections	Eisenreich et al. ⁹³ (1977)

Table 2

P Deposition Rate (mgPm ² yr ⁻¹) Mean, Range, Sampling Period	N Deposition Rate (mgNm ² yr ⁻¹)	Sampling Location and Cover Type	Notes on Sampling Design and Techniques	Sources
32.5 24 - 53 4 years	625	Rawson Lake, Ontario, Canada; forested	Single station on island in lake, 4 years of data, bulk collections with 1.5-mm screening	Schindler et al. ² (1976)
35.8 No range 1 year	587	Piburger See, Austria; Forested	Five stations on lake for large material + 10 for fine; biweekly, bulk samples, large samplers + 1-mm screen, small w/o screens	Psenner ⁷⁴ (1984)
40.1 38.3 - 42.0 2.5 years	None	Mediterranean Sea, Corsica Island; marine	Single station on the coast, 10-m tower, 2.5 years of data, bulk collections, sampling period 15 days	Bergametti, et al. ⁹⁷ (1992)
40.7 11.3 - 76.6 5 years	None	Various land uses, Southern Florida	Thirteen stations with Aerochem ^R wet/dry collectors, extensive data screening and uncertainty analysis; Std. Dev. = 33 mg P; range reflects site means for 5 years	Ahn and James ²¹ (2001)
41.0 27.4 - 70.6 12 years	1176	Chesapeake Bay; rural, forested, wetland	Single-site event sampling since 1978 for TN and 1980 for TP, bulk collections, 13-m tower (Baltimore/Washington D.C. airshed)	Jordan et al. ³⁵ (1995)
50 17 - 96 1 year	750	Florida, coastal, rural, agricul. and urban	24 sites within the state, sampled biweekly, bulk collections using buckets for 12 months in 1978/79	Hendry et al. ²³ (1981)
59 24 - 96 1 year	910	Florida,urban, maritime, rural and agricul.	Sites at 4 locations in Florida, sampled biweekly, wet/dry buckets, Aerochem ^R wet/dry collectors, 12 months 1978/79	Hendry et al. ²³ (1981)
62.0 16 - 118 5 years	None	Northern Everglades marsh sites	Four marsh sites with Aerochem ^R wet/dry collectors, weekly samples, extensive contamination screening; range applies to site averages	Walker ⁴² (1999)
93.3 52 - 176 1 year	859	Tampa Bay Watershed; misc. land uses	Seven stations, sampled biweekly, Bulk collections using funnel samplers; value is mean of 7 stations	Dixon et al. ²⁴ (1996)

P Deposition Rate (mgPm ⁻² yr ⁻¹) Mean, Range	Sampling Location and Cover Type	Notes on Sampling Design and Techniques; Misc. Comments	References (Figures or Tables)
23.8 3.6 - 60	Mean for 17 forests in United States and Western Europe	Various annual mass balance studies of key elements, primarily bulk collections	Likens and Bormann ²⁸ (1995) (Table 24)
26. 5 - 60 +	45 literature values for P inputs worldwide	Weighted mean based on ref. Figure 4; some values above 60 not included in figure; based on summary tables or cases published in the literature; one-year value of 22 from Gibson et al. not included	Gibson et al. ²⁹ (1995) (Figure 4)
30. 7 - 150	North America, continental average	Twenty-five authors summarized in ref. Table 3; average given in ref. Table 6, various approaches to measurements, 1960-1976	Graham and Duce ²⁶ (1979) (Table 3)
33.8 8.9 - 102	Great Lakes, regional mean	Nine annual means, 6 studies, bulk deposition samples, monthly data	Eisenreich et al. ³³ (1977)
35.9 12.4 - 93.3	Typical value for aquatic ecosystems	Average of 17 values, each representing multiple annual measurements of bulk deposition for a site or region; data in Table 2, this study	This study (Table 2)
50. 20 - 200	Western Europe, regional average.	Average from 16 authors reviewed in ref. Table 4, average from ref. Table 6; bulk deposition data 1951-1976	Graham and Duce ²⁶ (1979) (Table 6)
65.7 7 - 170	P input to ecosystems worldwide	Mean deposition rate for 20 sites compiled after scrutinizing for soundness; most from North America and Western Europe; bulk measurements with funnels	Newman ²⁷ (1995) (Table 5)

Table 3

P Deposition Rate (mgPm ⁻² yr ⁻¹)	Location, Sources and Other Notes	Notes on Sampling Design and Techniques; Misc. Comments	References (Figures or Tables)
0.9	P input from dust transported from Asia to the Hawaiian Islands	Based on the application of geochemical tracers to separate atmospheric dust from Hawaiian basalt soils; dust is transported over 6000 km from Asia.	Chadwick et al. ¹⁰⁵ (1999)
1.1 - 4.7	Saharan dust delivered to the Amazon Basin	Trajectory analysis of air masses used with concentration data and percent mass of Saharan dust to estimate a deposition rate for P into the Amazon Basin, South America.	Swap et al. ¹³³ (1992) (Table 4)
3.2	P deposition to N.E. Spain in "red rains" from North Africa	Based on data from Catalonia, Spain collected over 11 years; isentropic back trajectories used to identify sources and P averaged 0.063% of North African dust	Avila et al. ⁸⁰ (1998)
5 - 6	Deposition rate for remote areas in Northern Hemisphere	Based on data from United States, Canada, Finland, Russia, Sweden, and other sites.	Ahl (1988) ¹³⁴ (Tables 3 and 4)
13	P accumulation rate for raised bogs in Southern Sweden	Based on recent rates in the upper four cm of sediment cores; rate 4-to-6 fold higher than that for pre-industrial period (2-to-3 mgP/m ² /yr)	Ahl (1988) ¹³⁴ (Table 5)
21.5	Pre-industrial P accumulation rate at site TS7 in the Taylor Slough drainage of the Southern Florida Everglades	Based on mean accumulation rate for sections of a sediment core below 6 cm, dated at 1892; one high value at 12.5cm excluded. Other nearby sites had pre-1900 mean accumulation rates from 17 to 78 mgP/m ² /yr. William Orem (pers. com.) recommended TS7 as an undisturbed site supported almost solely by rainfall. Upper two sections of core, post-industrial period, averaged 26.1 mgP/m ² /yr.	Gough et al. (2001) ¹³⁵ (data from Tables 10 and 11)

Table 4

P Deposition Rate (mgPm ⁻² yr ⁻¹)	Location, Sources and Other Notes	Notes on Sampling Design and Techniques; Misc. Comments	References (Figures or Tables)
32.5 (a) 15.6 (b)	North Yorkshire, England, rural setting	Rates from 17 monthly dust collections; mean data from duplicate wet inverted Frisbees. Dust mass assumed to contain 0.001P. Stations near Barlow (a) and Smeaton (b), England.	Vallack ¹¹⁷ (1995) (Tables 1 (a) and 2 (b))
135.	Urban site, Chicago, Illinois	Based on particle mass and size data from a flat plate with Mylar strips and concentration/size data from Wide-Range Aerosol Classifier. Multi-step calculation method used with 21 days of summer data; Rate assumes particles contain P= 0.001mass; dry deposition only.	Lin et al. ⁵⁶ (1994) (Table 3)
94.6	Urban site, Chicago, Illinois	Based on 17 summer samples, both day and night; mass flux by size from a flat deposition plate with Mylar strips and concentration data from a Noll Rotary Impactor. Rate assumes P = 0.001 mass. With upper and lower surfaces included, deposition rate would be 126.1 mgPm ⁻² yr ⁻¹ ; dry deposition only.	Noll et al. ³² (1988)
15.	Ninety Mile Beach, coast of North Island, New Zealand	Based on three-month study in 1983: dry deposition to a 10-cm Petri dish on 11-m tower from five, 50-hr samples; wet deposition via funnel in seven events. Dry data biased low due to smooth surface and summer collections only with onshore winds.	Chen et al. ⁶⁴ (1985)
47 (a) 14.6 (b) 9.9 (c)	a. urban, Chicago b. rural, South Haven c. aquatic, Lake Michigan	Based on three sampling periods in July and August, 1991 in Illinois with Noll Rotary Impactor for concentration and flat plate sampler for size-specific flux. Multi-step method used to calculate mass flux; converted at P=0.001mass; rates conservative due to summer sampling and no wet deposition data.	Holsen et al. ⁵⁵ (1993)
5.9	Aquatic site, Southern Lake Michigan	Data from 3-to-5 days per month from May through September 1977; high-volume sampler and three-stage cascade impactor, plus meteorological data. Mean deposition velocity and concentration used for rates; rate biased low due to the exclusion of giant particles, low deposition velocities, summer data only, and no wet deposition data; probable rate = >15 mgPm ⁻² yr ⁻¹ .	Dolske and Sievering ¹³⁶ (1980) (Table 6)

Table 5




Figure 1




Figure 3

