

PESTICIDE SURFACE WATER AND SEDIMENT QUALITY REPORT

SEPTEMBER 2015 SAMPLING EVENT



Richard J. Pfeuffer
South Florida Water Management District
MSC 4441
3301 Gun Club Road
West Palm Beach, FL 33406

Table of Contents

Summary.....2
Background and Methods.....2
Results3
Usage and Water Quality Impacts4
Quality Assurance Evaluation.....7
Figure 1. South Florida Water Management District Pesticide Monitoring Network.....8
Table 1. Method detection limits (MDLs) and practical quantitation limits (PQLs) for June 2015 sampling event.....9
Table 2. Summary of pesticide residues ($\mu\text{g/L}$) detected above the method detection limit in surface water samples collected by SFWMD in September 2015.....10
Table 3. Summary of pesticides residues ($\mu\text{g/Kg}$) detected above the method detection limit in sediment samples collected by SFWMD in September 2015.....11
Table 4. Selected properties of pesticides detected during the September 2015 sampling event.....12
Table 5. Toxicity of pesticides detected during the September 2015 sampling event to freshwater aquatic invertebrates and fishes ($\mu\text{g/L}$).13
Glossary14
References15

Summary

As part of the South Florida Water Management District’s (SFWMD) quarterly ambient monitoring program, unfiltered water and sediment samples were collected September 21 to 24, 2015, and analyzed for over 70 pesticides and/or products of their degradation.

The herbicides 2,4-D, ametryn, atrazine, bentazon, diuron, and metribuzin, along with the insecticide/degradate imidacloprid, were detected in one or more of these surface water samples. No harmful impacts are expected from the detected pesticides.

The insecticides/degradates chlordane, DDD, DDE, DDT, dieldrin, beta endosulfan, endosulfan sulfate, and permethrin, along with three PCB compounds were found in the sediment at several locations. One chlordane and one DDE compound sediment concentrations were of a magnitude considered to have a harmful effect to freshwater sediment-dwelling organisms or wildlife. No harmful impacts are expected from the other detected pesticides.

The compounds and concentrations found are typical of those expected from an area of intensive historical and contemporary agricultural activity.

Background and Methods

The SFWMD pesticide monitoring network includes sites designated in the Everglades Settlement Agreement, the Lake Okeechobee Protection Act Permit, and the non-Everglades Construction Project (non-ECP) permit. The canals and marshes depicted in **Figure 1** are protected as Florida

Administrative Code (F.A.C.) 62-302 Class III (fishable and swimmable) waters, while Lake Okeechobee is protected as a Class I drinking water supply. Water Conservation Area 1 (WCA-1) and the Everglades National Park are also designated as Outstanding Florida Waters, to which anti-degradation standards apply. Surface water and sediment are sampled quarterly and semiannually, respectively, upstream at each structure identified in the permit or agreement. Sediment samples are collected using a petite Ponar® dredge.

Seventy-four pesticides and degradation products were analyzed in samples from 25 of the network 26 sites (**Figure 1**). The analytes, their respective method detection limits (MDLs), and practical quantitation limits (PQLs) are listed in **Table 1**. All the analytical work is performed by the Florida Department of Environmental Protection (FDEP) Central Laboratory in Tallahassee, Florida. Analytical method details can be found at the following location: <http://www.dep.state.fl.us/labs/cgi-bin/sop/chemsop.asp>.

To evaluate the potential impacts on aquatic life, the observed concentration is compared to the appropriate criterion outlined in F.A.C. 62-302.530. If a pesticide compound is not specifically listed, acute and chronic toxicity criterion are calculated as one-third and one-twentieth, respectively, of the amount lethal to 50% of the test organisms in 96 hours, using the lowest technical grade effective concentration 50 (EC₅₀) or lethal concentration 50 (LC₅₀) reported in the summarized literature for the species significant to the indigenous aquatic community (F.A.C. 62-302.200). Each pesticide's description and possible uses and sites of application described herein are taken from Hartley and Kidd (1987). Sediment concentrations are compared to freshwater sediment quality assessment guidelines (MacDonald Environmental Sciences, Ltd., and United States Geological Survey, 2003). A value below the threshold effect concentration (TEC) should not have a harmful effect on sediment-dwelling organisms. Values above the probable effect concentration (PEC) demonstrate that harmful effects to sediment-dwelling organisms are likely to be frequently or always observed. This summary covers surface water and sediment samples collected from September 21 to September 24, 2015.

Results

At least one pesticide was detected in surface water at 24 of the 26 sites (**Table 2**) and in sediment at 15 of the 17 sites (**Table 3**). The non-ECP permit requires sampling at S142 only during discharge or flow events. For this sampling event, no sample was obtained due to the lack of discharge at the time of sample collection. A minor modification of the Lake Okeechobee Water Control Structure Operations Permit (#0174552-010, dated December 18, 2011) eliminated sediment sampling at S65E, S191, and FECSR78. Additionally, sediment sampling was reduced to an annual frequency at S2, S3, and S4 for only ametryn, chlordane, DDD, DDE, and DDT analysis, which was not performed during this sampling event. Sediment samples are not collected at S333, S356-334, and TAMBR105, due to no requirement in the respective mandate. All of these compounds except for permethrin, have previously been detected in this monitoring program.

The above findings must be considered with the caveat that pesticide concentrations in surface water and sediment may vary significantly in relation to the timing and magnitude of pesticide application, rainfall events, pumping and other factors, and that this was only one sampling event. The possible

acute and chronic toxicity and environmental fate impacts are reported based on the single sampling event and do not take into account previous monitoring data.

Usage and Water Quality Impacts

2,4-D: 2,4-D is a selective systemic herbicide used for the post-emergence control of annual and perennial broad leaf weeds in terrestrial (grassland, established turf, sugarcane, rice, and on non-crop areas) as well as aquatic areas. Environmental fate and toxicity data in **Tables 4 and 5** indicate that 2,4-D (1) has minimum loss from soil by surface adsorption, with a moderate loss by leaching and surface solution; (2) is slightly toxic to mammals and relatively non-toxic to fish; and (3) does not bioaccumulate significantly. The highest 2,4-D concentration in surface water was detected at S6 (2.8 micrograms per liter [$\mu\text{g/L}$]) (**Table 2**). Using these criteria, this observed level should not have an acute or chronic effect on fish or aquatic invertebrates. 2,4-D was not detected in the sediment.

Ametryn: Ametryn is a selective terrestrial herbicide registered for use on sugarcane, bananas, pineapple, citrus, corn, and non-crop areas. Most algal effects occur at concentrations greater than ($>$) 10 $\mu\text{g/L}$ (Verschueren, 1983). Environmental fate and toxicity data in **Tables 4 and 5** indicate that ametryn (1) is lost from soil relatively easily by leaching, surface adsorption, and in surface solution; (2) is relatively non-toxic to mammals and fish; and (3) does not bioconcentrate significantly. Additional fish toxicity data include a 96-hour LC_{50} of 14.1 milligrams per liter (mg/L) for goldfish (Hartley and Kidd, 1987). The ametryn surface water concentrations found in this sampling event ranged from 0.031 to 0.073 $\mu\text{g/L}$ (**Table 2**). Using these criteria, these observed surface water concentrations should not have an acute, detrimental impact on fish or aquatic invertebrates. Ametryn was not detected in the sediment.

Atrazine: Atrazine is a selective systemic herbicide registered for use on pineapple, sugarcane, corn, rangelands, ornamental turf and lawn grasses, and non-crop areas. Environmental fate and toxicity data in **Tables 4 and 5** indicate that atrazine (1) is easily lost from soil by leaching and in surface solution, with moderate loss from surface adsorption; (2) is relatively non-toxic to mammals and fish; and (3) does not bioconcentrate significantly. Additional fish toxicity data include a 96-hour LC_{50} of 76 mg/L for carp, 16 mg/L for perch, and 4.3 mg/L for guppies (Hartley and Kidd, 1987). Also, in a flow-through bioassay, the maximum acceptable toxicant concentration (MATC) of atrazine was 90 and 210 $\mu\text{g/L}$ for bluegill and fathead minnow, respectively (Verschueren, 1983). The draft ambient aquatic life water quality criterion identifies a one-hour average concentration that does not exceed 1,500 $\mu\text{g/L}$ more than once every three years on the average (United States Environmental Protection Agency [U.S. EPA], 2003a). The atrazine surface water concentrations found in this sampling event at 11 of the 25 sampling locations, ranged from 0.010 to 0.16 $\mu\text{g/L}$ (**Table 2**). Using these criteria, these observed surface water concentrations should not have an acute or chronic detrimental impact on fish or invertebrates. Atrazine was not detected in the sediment.

Bentazon: Bentazon is a contact herbicide used for post-emergence control of many annual broad-leaved weeds in beans, peas, rice, and established turf. Environmental fate and toxicity data in

Tables 4 and 5 indicate that bentazon (1) is easily lost from soil by leaching, with moderate loss from surface solution, and minimum loss by surface adsorption; (2) is relatively non-toxic to mammals and fish; and (3) does not bioconcentrate significantly. The highest detected concentration of 0.68 µg/L at S5A and S7 (**Table 2**), is below any level that would have an acute or chronic detrimental impact on fish or aquatic invertebrates.

Chlordane: Chlordane is a chlorinated hydrocarbon previously used as a contact insecticide. Chlordane consists of 72% and 23% of the stereoisomers alpha and gamma chlordane, respectively (U.S. EPA, 1997). Environmental fate and toxicity data in **Tables 4 and 5** indicate that chlordane (1) is moderately toxic to mammals and highly toxic to fish; and (2) has the potential for significant bioconcentration. Freshwater sediment quality assessment guidelines identified a TEC of 3.2 µg/Kg and PEC of 18 µg/Kg for chlordane. The detected sediment residue at S5A (18 µg/Kg, respectively) are at a concentration where harmful effects to sediment-dwelling organisms may be frequently or always observed (**Table 3**). While the use of this compound has been discontinued in recent years, its persistence and tendency to accumulate in sediments makes chlordane a compound of concern. Chlordane was not detected in the surface water.

DDD, DDE, DDT: DDE is an abbreviation of **dichlorodiphenyldichloroethylene** [2, 2-bis (4-chlorophenyl)-1, 1-dichloroethene]. DDE is an environmental dehydrochlorination product of DDT (**dichlorodiphenyltrichloroethane**), a popular insecticide for which the U.S. EPA cancelled all uses in 1973. The large volume of DDT used, the persistence of DDT, DDE and another metabolite, DDD (**dichlorodiphenyldichloroethane**), and the high K_{oc} of these compounds account for the frequent detections in sediments. The large hydrophobicity of these compounds also results in a significant bioconcentration factor (**Table 4**). In sufficient quantities, these residues have reproductive effects in wildlife and carcinogenic effects in many mammals.

The detected DDD sediment concentration at S5A (8.6 µg/Kg) falls between the TEC (4.9 µg/Kg) and PEC (28 µg/Kg) (**Table 3**). This concentration may have the possibility for harmful effects on freshwater sediment-dwelling organisms. DDD was not detected in the surface water.

The TEC is 3.2 µg/Kg and the PEC is 31 µg/Kg for DDE in freshwater sediments. The concentration of DDE detected at S5A (**Table 3**) exceeded the PEC and frequently or always has the possibility for impacting sediment-dwelling organisms. The DDE concentrations detected at S178, S177, and S6, falls between the TEC and PEC. These concentrations may have the possibility for harmful effects on freshwater sediment-dwelling organisms. DDE was not detected in the surface water.

None of the DDT concentrations detected (**Table 3**) exceeded the TEC (4.2 µg/Kg). No DDT was detected in the surface water.

Dieldrin: Dieldrin is a non-systemic insecticide with all uses canceled in the United States. Environmental fate and toxicity data in **Tables 4 and 5** indicate that dieldrin (1) is highly toxic to mammals and fish; and (2) bioconcentrates significantly due to this compounds hydrophobicity. The high K_{oc} and low water solubility accounts for dieldrin's affinity for sediment. The dieldrin

concentration at S5A (0.84 µg/Kg) (**Table 3**) is less than the TEC (1.9 µg/Kg) and should not have an impact on sediment-dwelling freshwater organisms. Dieldrin was not detected in the surface water.

Diuron: Diuron is a selective, systemic terrestrial herbicide registered for use on sugarcane, bananas, and citrus. Environmental fate and toxicity data in **Tables 4 and 5** indicate that diuron (1) is easily lost from soil in surface solution, with moderate loss from leaching or surface adsorption; (2) is relatively non-toxic to mammals and fish; and (3) does not bioconcentrate significantly. Additional fish toxicity data include a 96-hour LC₅₀ of 25 mg/L for guppies (Hartley and Kidd, 1987). Crustaceans are affected at lower concentrations with a 48-hour LC₅₀ of 1.4 mg/L for water fleas and a 96-hour LC₅₀ of 0.7 mg/L for water shrimp (Verschueren, 1983). Most algal effects occur at concentrations > 10 µg/L (Verschueren, 1983). The highest surface water concentration of diuron found during this sampling event was 0.012 µg/L at S140 (**Table 2**). Using these criteria, this concentration should not have an acute, harmful impact on fish, aquatic invertebrates, or algae.

Endosulfan: Endosulfan is a non-systemic insecticide and acaricide registered for use on many crops, including beans, tomatoes, corn, cabbage, citrus, and ornamental plants. Technical endosulfan is a mixture of the two stereoisomeric forms, alpha (α) and beta (β). Endosulfan is highly toxic to mammals, with an acute oral LD₅₀ for rats of 70 mg/Kg (**Table 4**). The Soil Conservation Service (SCS) rates endosulfan with an extra small potential for loss due to leaching, a large potential for loss due to surface adsorption and a moderate potential for loss in surface solution (**Table 4**). Beta endosulfan's water solubility and Henry's law constant (1.91 x 10⁻⁵ atm – m³/mole) (Lyman, et al., 1990) indicate volatilization may be significant in shallow waters. The bioconcentration factors indicate a low to moderate degree of accumulation in aquatic organisms (**Table 4**). Beta endosulfan was detected in the sediment at S177 (0.35 µg/Kg) in the South Miami-Dade County farming area (**Table 3**). However, a sediment quality assessment guideline has not been developed. Endosulfan was not detected in the surface water.

Endosulfan sulfate: Endosulfan sulfate is an oxidation metabolite of the insecticide endosulfan. The water solubility and Henry's law constant (9.63 x 10⁻⁸ atm – m³/mole) (Lyman, et al., 1990) indicate that endosulfan sulfate is less volatile than water and concentrations will increase as water evaporates (**Table 4**). Endosulfan sulfate has a relatively high degree of accumulation in aquatic organisms (**Table 4**). The highest endosulfan sulfate concentration was detected in the sediment at S178 (0.50 µg/Kg) (**Table 3**). However, no sediment quality assessment guideline has been developed for endosulfan sulfate. Endosulfan sulfate was not detected in the surface water.

Imidacloprid: Imidacloprid is a systemic insecticide registered for use on a variety of row crops and turf grass applications as well as for flea control. Environmental fate and toxicity data in **Tables 4 and 5** indicate that imidacloprid (1) is soluble in water; (2) is slightly toxic to mammals and relatively non-toxic to fish; and (3) does not bioconcentrate significantly. The highest detected concentration of 0.053 µg/L at S9 (**Table 2**) is below any level that would have an acute or chronic detrimental impact on fish or aquatic invertebrates.

Metribuzin: Metribuzin is a selective systemic herbicide used on a variety of crops including potatoes, tomatoes, sugarcane, and peas. Environmental fate and toxicity data in **Tables 4 and 5** indicate that metribuzin (1) has a large potential for loss due to leaching, a medium potential for loss in surface solution, and a small potential for loss due to surface adsorption; (2) is relatively non-toxic to mammals and fish; and (3) does not bioaccumulate significantly. The only concentration of metribuzin detected was 0.026 µg/L at S2 (**Table 2**). Using these criteria, this surface water concentration should not have an acute impact on fish or aquatic invertebrates. Metribuzin was not detected in the sediment.

Permethrin: Permethrin is a broad spectrum insecticide used on fruit, vegetable, ornamental and potatoes. Environmental fate and toxicity data in **Tables 4 and 5** indicate that permethrin (1) has a small potential for loss due to leaching and surface solution and a medium potential for loss due to surface adsorption; (2) is highly toxic to fish; and (3) does not bioconcentrate significantly. The only concentration of permethrin was detected at S190 at 3.2 µg/Kg (**Table 2**). However, no sediment quality assessment guideline has been developed for permethrin. Permethrin was not detected in the surface water.

PCBs: Polychlorinated biphenyls (PCBs) is the generic term for a group of 209 congeners that contain a varying number of substituted chlorine atoms on one or both of the biphenyl rings. PCB-1242, PCB-1254, and PCB-1260 is a commercial grade mixture containing 42, 54, and 60, respectively, percent chlorine by weight. Production of PCBs was banned in 1978 and closed system uses are being phased out. In natural water systems, PCBs are found primarily absorbed to suspended sediments due to the very low solubility in water (Callahan et al., 1979). The tendency of PCBs for adsorption increases with the degree of chlorination and with the organic content of the adsorbent. While the production ban, phase out of uses, and stringent spill clean-up requirements have significantly reduced environmental loadings in recent years, the persistence and tendency to accumulate in sediment and bioaccumulate in fish, make this class of organochlorine compounds especially problematic. The TEC and PEC are 60 µg/Kg and 680 µg/Kg, respectively, for total PCBs. None of the detected sediment residues would have an impact freshwater sediment-dwelling organisms, since the concentrations are all below the TEC (**Table 3**). None of the PCB congeners were detected in the surface water.

Quality Assurance Evaluation

No pesticide analytes were detected in the equipment blanks or field blank performed at S191, US41-25, S8, and S5A. All of the collected samples were shipped and all bottles were received.

Pesticide Monitoring Program Report: September 2015 Sampling Event

Table 1. Method detection limits (MDLs) and practical quantitation limits (PQLs) for September 2015 sampling event.

Pesticide or metabolite	Water: range of MDLs - PQLs (µg/L)	Sediment: range of MDLs - PQLs (µg/Kg)	Pesticide or metabolite	Water: range of MDLs - PQLs (µg/L)	Sediment: range of MDLs - PQLs (µg/Kg)
2,4-D	0.002 - 0.2	9.0 - 310	endrin aldehyde	0.0035 - 0.016	0.37 - 17
2,4,5-T	0.002 - 0.01	9.0 - 310	ethion	0.0090 - 0.04	2.3 - 110
2,4,5-TP (silvex)	0.002 - 0.01	9.0 - 310	ethoprop	0.0045 - 0.02	1.1 - 54
acifluorfen	0.002 - 0.01	9.0 - 310	fenamiphos	0.027 - 0.12	4.6 - 220
alachlor	0.054 - 0.24	17 - 770	fonofos	0.0090 - 0.04	1.1 - 54
aldrin	0.0017 - 0.008	0.092 - 4.3	heptachlor	0.0017 - 0.008	0.046 - 2.2
ametryn	0.0090 - 0.04	2.3 - 110	heptachlor epoxide	0.0017 - 0.008	0.092 - 4.3
atrazine	0.0090 - 0.04	2.3 - 110	hexazinone	0.027 - 0.12	6.9 - 320
atrazine desethyl	0.0090 - 0.04	N/A	imidacloprid	0.002 - 0.01	N/A
atrazine desisopropyl	0.0090 - 0.04	N/A	linuron	0.004 - 0.02	1.4 - 65
azinphos methyl (guthion)	0.018 - 0.08	6.9 - 320	malathion	0.0090 - 0.04	2.3 - 110
bentazon	0.002 - 0.01	N/A	metalaxyl	0.036 - 0.16	N/A
α-BHC (alpha)	0.0017 - 0.008	0.046 - 2.2	methamidophos	N/A	9.2 - 430
β-BHC (beta)	0.0017 - 0.013	0.046 - 2.2	methoxychlor	0.0087 - 0.04	0.046 - 2.2
δ-BHC (delta)	0.0017 - 0.008	0.030 - 1.4	metolachlor	0.054 - 0.24	14 - 650
γ-BHC (gamma) (lindane)	0.0017 - 0.008	0.046 - 2.2	metribuzin	0.019 - 0.088	4.6 - 220
bromacil	0.036 - 0.16	14 - 650	mevinphos	0.0090 - 0.04	2.3 - 110
butylate	0.018 - 0.08	N/A	mirex	0.0035 - 0.016	0.092 - 4.3
carbophenothion (trithion)	0.0052 - 0.024	0.14 - 6.5	monocrotophos	N/A	2.8 - 130
chlordane	0.017 - 0.08	0.46 - 2.2	naled	0.036 - 0.16	9.2 - 430
chlorothalonil	0.0069 - 0.032	0.18 - 8.6	norflurazon	0.027 - 0.12	6.9 - 320
chlorpyrifos ethyl	0.0094 - 0.04	2.3 - 110	parathion ethyl	0.018 - 0.08	2.3 - 110
chlorpyrifos methyl	0.0090 - 0.04	2.3 - 110	parathion methyl	0.0090 - 0.04	2.3 - 110
cypemethrin	0.010 - 0.048	0.23 - 11	PCB-1016	0.017 - 0.08	0.92 - 43
DDD-P,P'	0.0035 - 0.016	0.046 - 2.2	PCB-1221	0.017 - 0.08	0.92 - 43
DDE-P,P'	0.0035 - 0.016	0.030 - 1.4	PCB-1232	0.017 - 0.08	0.92 - 43
DDT-P,P'	0.0035 - 0.016	0.11 - 5.2	PCB-1242	0.017 - 0.08	0.92 - 43
demeton	0.022 - 0.096	2.8 - 130	PCB-1248	0.017 - 0.08	0.92 - 43
diazinon	0.0090 - 0.04	2.3 - 110	PCB-1254	0.017 - 0.08	0.92 - 43
dicofol (kelthane)	0.021 - 0.096	0.092 - 4.3	PCB-1260	0.017 - 0.08	0.92 - 43
dieldrin	0.0017 - 0.008	0.18 - 8.6	permethrin	0.0087 - 0.04	0.11 - 5.2
disulfoton	0.0045 - 0.02	1.1 - 54	phorate	0.0045 - 0.02	1.1 - 54
diuron	0.002 - 0.01	1.4 - 65	prometon	0.018 - 0.08	N/A
α-endosulfan (alpha)	0.0017 - 0.016	0.18 - 8.6	prometryn	0.018 - 0.08	4.6 - 220
β-endosulfan (beta)	0.0017 - 0.016	0.18 - 8.6	simazine	0.0090 - 0.04	2.3 - 110
endosulfan sulfate	0.0035 - 0.016	0.092 - 4.3	toxaphene	0.087 - 0.4	4.6 - 220
endrin	0.0035 - 0.016	0.18 - 8.6	trifluralin	0.0069 - 0.032	0.074 - 3.4

N/A = not analyzed

Pesticide Monitoring Program Report: September 2015 Sampling Event

Table 2. Summary of pesticide residues ($\mu\text{g/L}$) detected above the method detection limit in surface water samples collected by SFWMD in September 2015.

Date	Location	Flow	2,4-D	ametryn	atrazine	bentazon	diuron	imidacloprid	metribuzin	Number of compounds detected at location
9/21/2015	S191	Y	0.082	-	-	0.0023 I	-	0.0062 I	-	3
	S2	N	0.58	0.038 I	0.13	0.034	-	-	0.026 I	5
	S3	N	0.11	0.035 I	0.073	0.034	-	-	-	4
	S18C	Y	-	-	-	0.029	-	0.0053 I	-	2
	S178	Y	-	-	-	0.12	-	0.040	-	2
	S177	N	-	-	-	0.0087 I *	-	0.0068 I *	-	2
	S4	N	0.38	0.058	0.16	0.046	0.0050 I	-	-	5
	FEC SR78	Y	-	-	-	-	0.0024 I	0.010	-	2
	S332DX	Y	-	-	-	0.010	-	-	-	1
	S65E	Y	0.0076 I	-	0.011 I	-	-	0.017	-	3
9/22/2015	S331	Y	-	-	-	0.0092 I	-	-	-	1
	S31	N	0.17	-	0.058	0.22	0.0037 I	0.038	-	5
	S356-334	Y	0.018	-	0.011 I	0.42	-	-	-	3
	S333	N	0.0062 I	-	0.010 I	0.31	-	-	-	3
	S12A	N	0.0060 I	-	-	0.010	-	-	-	2
	US41-25	N	-	-	-	-	-	-	-	0
9/23/2015	TAMBR105	Y	-	-	-	-	-	-	-	0
	S140	Y	0.0065 I	-	-	0.012	0.012	0.020	-	4
	S190	Y	0.40	-	-	-	-	-	-	1
	L3BRS	Y	0.027	-	-	0.0087 I	-	-	-	2
9/24/2015	S8	Y	0.27	-	-	0.018	-	0.0064 I	-	3
	S9	Y	0.19	-	0.067	0.060	-	0.053	-	4
	S7	Y	0.024	0.031 I	0.022 I	0.68	-	-	-	4
	S6	Y	2.8	0.045	0.13	0.32	-	-	-	4
Total number of compound detections	S5A	Y	0.80	0.073	0.024 I	0.68	-	-	-	4
				17	6	11	20	4	10	1

N = no, Y = yes, R = reverse

- = denotes that the result is below the method detection limit

I = value reported is less than the practical quantitation limit, and greater than or equal to the method detection limit

* = value is the average of replicate samples

Pesticide Monitoring Program Report: September 2015 Sampling Event

Table 3. Summary of pesticides residues ($\mu\text{g/Kg}$) detected above the method detection limit in sediment samples collected by SFWMD in September 2015.

Date	Location	Flow	chlordan	DDD-P,P'	DDE-P,P'	DDT-P,P'	dieldrin	beta endosulfan	endosulfan sulfate	permethrin	PCB-1242	PCB-1254	PCB-1260	Number of compounds detected at location
9/21/2015	S18C	Y	-	-	0.59 *	-	-	-	-	-	-	-	-	1
	S178	Y	4.8	-	5.4	-	-	-	0.50 I	-	-	-	-	3
	S177	N	1.2 I	0.24 I	13	0.43 I	-	0.35 I	0.20 I	-	-	-	-	6
	S332DX	Y	-	0.16 I	2.9	-	-	-	-	-	-	1.7 I	-	3
	S331	Y	-	0.092 I	1.6	-	-	-	-	-	-	-	-	2
9/22/2015	S31	N	-	0.063 I	0.46	-	-	-	-	-	1.4 I	-	2.3 I	4
	S12A	N	-	-	1.4 I	-	-	-	-	-	-	-	-	1
	US41-25	N	-	-	0.74	-	-	-	-	-	-	-	-	1
9/23/2015	S190	Y	-	-	0.15 I	-	-	-	-	3.2	-	-	-	2
	L3BRS	Y	3.7 I	0.21 I	1.0	-	-	-	-	-	5.4 I	-	-	4
	S8	Y	-	-	0.097 I	-	-	-	-	-	-	-	-	1
9/24/2015	S9	Y	-	0.086 I	1.9	-	-	-	-	-	-	5.5 I	-	3
	S7	Y	-	-	0.32	-	-	-	-	-	-	-	-	1
	S6	Y	6.4	3.7	16	0.61	-	-	-	-	-	-	1.5 I	5
	S5A	Y	18	8.6	65	2.5	0.84 I	-	-	-	-	18	-	6
Total number of compound detections			5	8	15	3	1	1	2	1	2	3	2	43

N = no, Y = yes, R = reverse

- = denotes that the result is below the method detection limit

I = value reported is less than the practical quantitation limit, and greater than or equal to the method detection limit

* = value is the average of replicate samples

Values in bold, italicized font are at a concentration that harmful effects to sediment-dwelling organisms are likely to be frequently or always observed.

Pesticide Monitoring Program Report: September 2015 Sampling Event

Table 4. Selected properties of pesticides detected during the September 2015 sampling event.

Common Name	Surface Water Standards F.A.C. 62-302 (µg/L)	Acute Oral LD ₅₀ For Rats (mg/Kg) (1)	Bioconcentration Factor (2)	Volatility from Water (2)	Soil Conservation Service (SCS) rating (3)			K _{oc} (mL/g) (3, 4)	Soil Half-life (days) (3, 4)	Water Solubility (WS) (mg/L) (3, 4)	U.S. EPA Carcinogenic Potential (5)
					LE	SA	SS				
2,4-D (acid)	(100)	375	13	I	M	S	M	20	10	890	D
ametryn	-	1,110	33	I	M	M	M	300	60	185	D
atrazine	-	3,080	86	I	L	M	L	100	60	33	C
bentazon	-	1,100	19	I	L	S	M	34	20	500	C
chlordane	0.0043	365 - 590	3,141	I	-	-	-	3,800	-	0.056	B2
DDD-P,P'	-	3,400	3,173	I	-	-	-	239,900	-	0.055	B2 ⁽⁸⁾
DDE-P,P'	-	880	2,887	S	-	-	-	243,220	-	0.065	B2 ⁽⁷⁾
DDT-P,P'	0.001	113	15,377	I	-	-	-	140,000	-	0.00335	B2 ⁽⁸⁾
diuron	-	3,400	75	I	M	M	L	480	90	42	D
dieldrin	0.0019	37 - 87	1,873	I	-	-	-	10,000 est.	-	0.14	B2
endosulfan	0.056	70	884	S	XS	L	M	12,400	50	0.53	-
endosulfan sulfate	-	-	2,073	I	-	-	-	-	-	0.117	-
imidacloprid	-	424 ⁽⁶⁾	18	I	-	-	-	178 ⁽⁶⁾	520 ⁽⁶⁾	510 ⁽⁶⁾	E
metribuzin	-	2,200	11	I	L	S	M	41	30	1,220	D
permethrin	-	8,900 ⁽⁹⁾	11,069	I	S	M	S	100,000	32	0.006	C ⁽⁹⁾
PCB's	0.014	-	-	-	-	-	-	-	-	-	B2

- = No data available

FDEP F.A.C. 62-302 surface water standards (7/2012) for Class III waters except Class I noted in ()

Bioconcentration Factor (BCF) calculated as $BCF = 10^{(2.71 - 0.564 \log WS)}$ (2); Volatility from water: R = rapid, I = insignificant, S = significant

SCS ratings are pesticide loss due to leaching (LE), surface adsorption (SA) or surface solution (SS) and grouped as large (L), medium (M), small (S), or extra small (XS)

B2 = probable human carcinogen; C = possible human carcinogen; D = not classified; E = evidence of non-carcinogen for humans (5)

(1) Hartley and Kidd (1987)

(3) Goss and Wauchope (1992)

(5) U.S. EPA (1996)

(7) U.S. EPA (1999)

(9) U.S. EPA (2009)

(2) Lyman, et al. (1990)

(4) Montgomery (1993)

(6) U.S. EPA (1994a)

(8) U.S. Department of Health (1994)

Pesticide Monitoring Program Report: September 2015 Sampling Event

Table 5. Toxicity of pesticides detected during the September 2015 sampling event to freshwater aquatic invertebrates and fishes (µg/L).

Pesticide Common Name	Water Flea (<i>Daphnia magna</i>)			Fathead Minnow # (<i>Pimephales promelas</i>)			Bluegill (<i>Lepomis macrochirus</i>)			Largemouth Bass (<i>Micropterus salmoides</i>)			Rainbow Trout # (<i>Oncorhynchus mykiss</i>)			Channel Catfish (<i>Ictalurus punctatus</i>)		
	48 hour EC ₅₀	Acute Toxicity (*)	Chronic Toxicity (*)	96 hour LC ₅₀	Acute Toxicity	Chronic Toxicity	96 hour LC ₅₀	Acute Toxicity	Chronic Toxicity	96 hour LC ₅₀	Acute Toxicity	Chronic Toxicity	96 hour LC ₅₀	Acute Toxicity	Chronic Toxicity	96 hour LC ₅₀	Acute Toxicity	Chronic Toxicity
2,4-D	25,000 (5)	8,333	1,250	133,000 (5)	44,333	6,650	180,000 (6) 900 (48 hr) (4)	60,000 -	9,000 -	-	-	-	100,000 (2) 110,000 (5)	33,333 36,667	5,000 5,500	-	-	-
ametryn	28,000 (5)	9,333	1,400	16,000 (7)	5,333	800	4,100 (2)	1,367	205	-	-	-	8,800 (2) 3,600 (7)	2,933 1,200	440 180	-	-	-
atrazine	6,900 (5)	2,300	345	15,000 (5)	5,000	750	16,000 (2)	5,333	800	-	-	-	8,800 (2) 5,300 (8)	2,933 1,767	440 265	7,600 (2)	2,533	380
bentazon	>100,000 (12)	33,333	5,000	-	-	-	>100,000 (12)	33,333	5,000	-	-	-	>100,000 (12)	33,333	5,000	-	-	-
chlordane	-	-	-	-	-	-	70 (3)	23	3.5	-	-	-	90 (3)	30	4.5	-	-	-
DDD-P,P'	3,200 (4)	1,067	160	4,400 (1)	1,467	220	42 (1)	14	2.1	42 (1)	14	2.1	70 (1)	23	4	1,500 (1)	500	75
DDE-P,P'	-	-	-	-	-	-	240 (1)	80	12	-	-	-	32 (1)	11	2	-	-	-
DDT-P,P'	-	-	-	19 (3)	6	1	8 (3)	2.7	0.4	2 (3)	0.7	0.1	7 (3)	2	0.4	16 (3)	5.3	0.8
dieldrin	-	-	-	16 (3)	5.3	0.8	8 (3)	2.7	0.4	-	-	-	10 (3)	3.3	0.5	4.5 (3)	1.5	0.2
diuron	1,400 (5) 1,400 (10)	467 467	70 70	14,200 (5) 14,000 (10)	4,733 4,667	710 700	5,900 (2)	1,967	295	-	-	-	5,600 (2)	1,867	280	-	-	-
endosulfan	166 (5)	55	8	1 (1)	0.33	0.05	1 (1) 2 (13)	0.33 0.67	0.05 0.10	-	-	-	1 (1) 3 (14) 1 (13) 0.3 (3) 0.8 (15)	0.33 1.00 0.33 0.10 0.27	0.05 0.15 0.05 0.02 0.04	1 (1) 1.5 (5)	0.33 0.50	0.05 0.08
imidacloprid	85,200 (11)	28,400	4,260	-	-	-	-	-	-	-	-	-	83,000 (11)	27,667	4,150	-	-	-
metribuzin	4,200 (5) 4,200 (9)	1,400 1,400	210 210	-	-	-	80,000 (2) 75,900 (9)	26,667 25,300	4,000 3,795	-	-	-	64,000 (2) 76,770 (9)	21,333 25,590	3,200 3,839	100,000 (5)	33,333	5,000
permethrin	-	-	-	-	-	-	0.79 (16)	0.26	0.04	-	-	-	-	-	-	-	-	-

- = No data available

(*) Florida Administrative Code (F.A.C.) 62-302.200, for compounds not specifically listed, acute and chronic toxicity standards are calculated as one-third and one-twentieth, respectively, of the amount lethal to 50% of the test organisms in 96 hours, where the 96 hour LC₅₀ is the lowest value which has been determined for a species significant to the indigenous aquatic community.

(#) Species is not indigenous. Information is given for comparison purposes only.

- | | | | |
|-------------------------------|---------------------------------|-----------------------|----------------------|
| (1) Johnson and Finley (1980) | (5) U.S. EPA (1991) | (9) U.S. EPA (1998) | (13) Sneider (1979) |
| (2) Hartley and Kidd (1987) | (6) Mayer and Eilersieck (1986) | (10) U.S. EPA (2003b) | (14) U.S. EPA (1977) |
| (3) Montgomery (1993) | (7) U.S. EPA (2005) | (11) U.S. EPA (1994a) | (15) U.S. EPA (2002) |
| (4) Verschuere (1983) | (8) U.S. EPA (2006) | (12) U.S. EPA (1994b) | (16) U.S. EPA (2009) |

Glossary

Bioconcentration Factor: The ratio of the concentration of a contaminant in an aquatic organism to the concentration in water, after a specified period of exposure via water only. The duration of exposure should be sufficient to achieve a near steady-state condition.

EC₅₀: A concentration necessary for 50 percent of the aquatic species tested to exhibit a toxic effect short of mortality (e.g., swimming on side or upside down, cessation of swimming) within a short (acute) exposure period, usually 24 to 96 hours.

Henry's law constant (H): Relates the concentration of a compound in the gas phase to its concentration in the liquid phase. The constant is calculated from the formula: $H = P_{vp}/S$ where P_{vp} is pressure in units of atmospheres and S is solubility in units of moles/meter³ for a compound.

K_{oc}: The soil/sediment partition or sorption coefficient normalized to the fraction of organic carbon in the soil. This value provides an indication of the chemical's tendency to partition between soil organic carbon and water.

LC₅₀: A concentration which is lethal to 50 percent of the aquatic animals tested within a short (acute) exposure period, usually 24 to 96 hours.

LD₅₀: The dosage which is lethal to 50 percent of the terrestrial animals tested within a short (acute) exposure period, usually 24 to 96 hours.

Method Detection Limits (MDLs): The minimum concentration of an analyte that can be detected with 99 percent confidence of its presence in the sample matrix.

Practical Quantitation Limits (PQLs): The lowest level of quantitation that can be reliably achieved within specified limit of precision and accuracy during routine laboratory operating conditions. The PQLs are further verified by analyzing spike concentrations whose relative standard deviation in 20 fortified water samples is < 15 percent. In general, PQLs are 2 to 5 times larger than the MDLs.

Probable Effect Concentration (PEC): The probable effect concentration is intended to identify concentrations above which harmful effects to sediment-dwelling organisms are likely to be frequently or always observed.

Soil or water half-life: The time required for one-half the concentration of the compound to be lost from the water or soil under the conditions of the test.

Threshold Effect Concentration (TEC): The threshold effect concentration is intended to identify concentrations below which harmful effects to freshwater sediment-dwelling organisms are unlikely to be observed.

References

Callahan, M.A., M.W. Slimak, N.W. Gabel, I.P. May, C.F. Fowler, J.R. Freed, P. Jennings, R.L. Durfee, F.C. Witmore, B. Maestri, W.R. Mabey, B.R. Holt, and C. Gould. (1979). *Water-Related Environmental Fate of 129 Priority Pollutants, Volume I*. USEPA 440/4-79-029a.

Goss, D. and R. Wauchope. (Eds.) 1992. *The SCS/ARS/CES Pesticide Properties Database: II Using It With Soils Data In A Screening Procedure*. Soil Conservation Service. Fort Worth, TX.

Hartley, D. and H. Kidd. (Eds.) 1987. *The Agrochemicals Handbook*. Second Edition, The Royal Society of Chemistry. Nottingham, England.

Johnson, W.W. and M.T. Finley. 1980. *Handbook of Acute Toxicity of Chemicals to Fish and Aquatic Invertebrates*. U.S. Department of the Interior, Fish and Wildlife Service Resource Publication 137. Washington, DC.

Lyman, W.J., W.F. Reehl, and D.H. Rosenblatt. 1990. *Handbook of Chemical Property Estimation Methods*. American Chemical Society, Washington, DC.

MacDonald Environmental Sciences, Ltd. and United States Geological Survey. 2003. Development and Evaluation of Numerical Sediment Quality Assessment Guidelines for Florida Inland Waters. Report to Florida Department of Environmental Protection. Tallahassee, FL.

Mayer, F.L. and M.R. Ellersieck. 1986. *Manual of Acute Toxicity: Interpretation and Database for 410 Chemicals and 66 Species of Freshwater Animals*. United States Fish and Wildlife Service Publication No. 160.

Montgomery, J.H. 1993. *Agrochemicals Desk Reference: Environmental Data*. Lewis Publishers. Chelsea, MI.

Schneider, B.A. (Ed.) 1979. *Toxicology Handbook, Mammalian and Aquatic Data, Book 1: Toxicology Data*. U.S. Environmental Protection Agency. U.S. Government Printing Office. Washington, DC. EPA-5400/9-79-003.

U.S. Department of Health 1994. 4,4-DDT, 4,4'-DDE, 4,4'-DDD Update. Public Health Service, Agency for Toxic Substances and Disease Registry

U.S. EPA 1977. *Silvacultural Chemicals and Protection of Water Quality*. Seattle, WA. EPA-910/9-77-036.

U.S. EPA 1991. Pesticide Ecological Effects Database. Ecological Effects Branch, Office of Pesticide Programs, Washington, DC.

U.S. EPA 1994a. Pesticide Fact Sheet: Imidacloprid

U.S. EPA 1994b. Reregistration Eligibility Decision (RED) Bentazon. EPA 738-R-94-029 September 1994.

U.S. EPA 1996. *Drinking Water Regulations and Health Advisories*. Office of Water. EPA 822-B-96-002.

U.S. EPA 1997. Toxicological Review of Chlordane (Technical) In Support of Summary Information on the Integrated Risk Information System (IRIS), Washington, DC.

U.S. EPA 1998. Reregistration Eligibility Decision (RED) Metribuzin, EPA 738-R-37-006 February 1998.

U.S. EPA 1999. Integrated Risk Information System (IRIS) on p,p-Dichlorodiphenyldichloroethylene. National Center for Environmental Assessment. Office of Research and Development. Washington, DC.

U.S. EPA 2002. Reregistration Eligibility Decision for Endosulfan. EPA 738-R-02-013 November 2002.

U.S. EPA 2003a. Ambient Aquatic Life Water Criteria for Atrazine. Revised Draft EPA-822-R-03-023. October 2003.

U.S. EPA 2003b. Reregistration Eligibility Decision (RED) Diuron. September 30, 2003.

U.S. EPA 2005. Reregistration Eligibility Decision (RED) for Ametryn; EPA 738-R-05-006 September 2005.

U.S. EPA 2006. Decisions Document for Atrazine.

U.S. EPA 2009. Reregistration Eligibility Decision (RED) for Permethrin; EPA 738-R-09-306 May 2009.

Verschueren, K. 1983. *Handbook of Environmental Data on Organic Chemicals*. Second Edition, Van Nostrand Reinhold Co. Inc. New York, NY.