

Disclaimer: Information contained in the report addresses environmental conditions only and is not the official South Florida Water Management District operations recommendation or decision.

M E M O R A N D U M

TO: John Mitnik, Chief, Operations, Engineering and Construction Bureau
Paul Linton, Chief, Operations Section

FROM: SFWMD Staff Environmental Advisory Team

DATE: November 27, 2018

SUBJECT: Weekly Environmental Conditions for Systems Operations

Summary

Weather Conditions and Forecast

Turning dry and cool; some scattered shower activity returning this weekend. A cold front is pushing through the southern end of the peninsula this morning and a second push of dry air will move through the area during the day today. Dry conditions and cool temperatures will persist over the area tonight through Thursday before temperatures moderate on Friday. The next cold front is forecast to stall across Florida's panhandle on Saturday and then return north. This boundary will bring some moisture and instability to the area and allow some scattered showers to pop up mainly over the interior Saturday afternoon. Widely scattered shower activity is then forecast to develop mainly over the interior each day Sunday and Monday. The next cold front has the potential to produce some moderate rainfall Wednesday of next week.

Kissimmee

Tuesday morning stages were 57.0 feet NGVD (1.0 feet below schedule) in East Lake Toho, 54.7 feet NGVD (0.3 feet below schedule) in Toho, and 49.5 feet NGVD (3.0 feet below schedule) in Kissimmee-Cypress-Hatchineha; headwater stages were 46.4 feet NGVD at S-65A and 25.9 feet NGVD at S-65D. Tuesday morning discharges were N/A at S-65, 283 cfs at S-65A, and 312 cfs at S-65E. Dissolved oxygen concentration in the Kissimmee River averaged 5.5 mg/L for the week. Kissimmee River mean floodplain depth on Sunday was 0.12 feet. No new recommendations.

Lake Okeechobee

Lake Okeechobee stage is 13.22 feet NGVD, falling 0.13 feet from the previous week and 0.61 feet over the past 30 days. Lake stages are the lowest they have been for this time of year since 2010 and are now 1.28 feet below the bottom of the preferred ecological envelope, which varies seasonally from 12.5 – 15.5 feet NGVD. However, given potential for heavy rainfall associated with El Niño conditions this winter and the poor condition of SAV and EAV in the nearshore zone, these lower lake stages are ideal for vegetation recovery. Cyanobacterial bloom potential decreased in the south but remains slightly elevated in the southwest region of the lake based on the latest NOAA image (November 21).

Estuaries

Total inflow to the St. Lucie Estuary averaged 157 cfs over the past week with 0 cfs coming from Lake Okeechobee. Surface salinity increased by 2 to 3 points at HR1, US1 and A1A Bridge over the last week. The seven-day average salinity at the US1 Bridge is in the good range for adult eastern oysters. Total inflow to the Caloosahatchee Estuary averaged 1,272 cfs over the past week with 685 cfs coming from the Lake. Surface salinity increased by 1 to 2 points at Val I-75 and downstream over the last week. The 30-day moving average surface salinity is 1.9 at Val I-75 and 7.1 at Ft. Myers.

Salinity conditions between Val I-75 and Ft. Myers are good for tape grass. Salinity conditions are in the good range for adult eastern oysters at Cape Coral and Shell Point.

Stormwater Treatment Areas

Over the past week, the STAs received approximately 6,100 acre-feet of Lake releases. The total amount of inflows to the STAs in WY2019 (since May 1, 2018) is approximately 1,173,000 acre-feet, which includes approximately 317,600 acre-feet of Lake releases. Most STA cells are at or above target depths. Operational restrictions are in place for construction related activities in STA-1W (all flow-ways). STA-5/6 Flow-ways 2 and 3 are offline for initiation of a Restoration Strategies project to grade non-effective treatment areas. This week, if Lake releases are sent to the WCAs and conditions allow, releases will be sent to STA-1E and STA-2.

Everglades

Over the last week water depths declined on average across the Everglades at a rate similar to previous weeks. Conditions within the Everglades are drying out unseasonably early but have remained stable over the last few weeks. Stages remain below the respective regulation lines in the WCAs with the exceptions of WCA-2A and 2B. WCA-2A is now receding parallel to its regulation line. WCA-3A North and northern WCA-1 continue to dry out as indicated by the WDAT model output. At the gauges located in WCA-3A North, in the northwest stages remain essentially unchanged over the last three weeks but fell in the northeast by 0.35 feet. Stages continued to decrease this week throughout Taylor Slough. However, stages in the northern reaches remain above average for this time of year while stages in the south are below average. Discharge from the five creeks is now below the long-term average. Salinities in Florida Bay increased on average this past week, and conditions at the western stations remain higher than their historic averages for this time of year.

KESSIMMEE BASIN

Kissimmee Basin Rainfall

The Upper Kissimmee Basin received 0.03 inches of rainfall in the past week and the Lower Basin received 0.10 inches (SFWMD Daily Rainfall Report 11/23/2018).

Upper Kissimmee Basin

Stages and departures in the Kissimmee Chain of Lakes (KCOL) are shown in **Table 1**. KCOL stage hydrographs with respective regulation schedules and rainfall are shown in Figures 1-7.

Table 1. Average discharge (cfs) for the preceding seven days, one-day stage (feet NGVD), and departures from KCOL flood regulation (R) or temporary schedules (T, A, or S). Provisional, real-time data are from SFWMD.
Report Date: 11/27/2018

Water Body	Structure	7-day Average Discharge (cfs) ¹	Stage Monitoring Site ²	Lake Stage (feet)	Schedule Type ³	Schedule Stage (feet)	Daily Departure (feet)						
							11/25/18	11/18/18	11/11/18	11/4/18	10/28/18	10/21/18	10/14/18
Lakes Hart and Mary Jane	S-62	0	LKMJ	60.0	R	61.0	-1.0	-1.0	-1.0	-1.1	-0.9	-0.7	-0.3
Lakes Myrtle, Preston, and Joel	S-57	0	S-57	61.1	R	62.0	-0.9	-0.8	-0.8	-0.8	-0.7	-0.4	-0.1
Alligator Chain	S-60	0	ALLI	63.0	R	64.0	-1.0	-0.9	-0.9	-0.9	-0.7	-0.4	-0.2
Lake Gentry	S-63	0	LKGT	61.5	R	61.5	0.0	0.1	0.0	-0.1	0.0	0.1	0.1
East Lake Toho	S-59	0	TOHOE	57.0	R	58.0	-1.0	-1.0	-0.9	-1.0	-0.9	-0.6	-0.3
Lake Toho	S-61	0	TOHOW, S-61	54.6	R	55.0	-0.4	-0.4	-0.4	-0.6	-0.5	-0.3	0.0
Lakes Kissimmee, Cypress, and Hatchineha	S-65	346	KUB011, LKIS5B	49.5	R	52.5	-3.0	-2.9	-2.8	-2.7	-2.5	-2.1	-1.4

¹ Seven-day average of weighted daily means through midnight.

² Names of in-lake monitoring sites and structures used to determine lake stage; if more than one site is listed, an average is reported.

³ T = temporary schedule, R = USACE flood control schedule, S = temporary snail kite schedule, A = projected ascension line, N/A = not applicable or data not available.

DATA ARE PROVISIONAL

Lower Kissimmee Basin

Discharges at Lower Basin structures are shown in Table 2. SFWDAT depth maps for the Phase I restoration area are shown in Figure 8. Kissimmee River floodplain stages at selected stations are shown in Figure 9.

Table 2. One-day and seven-day averages of discharge at S-65x structures, of dissolved oxygen concentration in the Phase I area river channel, and water depth in the Phase I area floodplain. Data are provisional real-time data from SFWMD.

Report Date: 11/27/2018

Metric	Location	1-Day Average		Average for the Preceding 7-Days ¹								9/23/18
		11/25/2018	11/25/18	11/18/18	11/11/18	11/4/18	10/28/18	10/21/18	10/14/18	10/7/18	9/30/18	
Discharge (cfs)	S-65	347	346	349	336	350	480	1,592	1,559	1,542	1,485	1,560
Discharge (cfs)	S-65A ²	248	254	256	252	288	383	1,394	1,382	1,391	1,416	1,532
Discharge (cfs)	S-65D ²	313	315	321	360	404	1,024	1,461	1,521	1,646	1,982	2,221
Headwater Stage (feet NGVD)	S-65D ²	26.04	26.20	26.66	27.15	27.62	27.69	27.78	27.89	27.81	27.81	27.75
Discharge (cfs)	S-65E ²	329	335	317	370	405	1,058	1,535	1,598	1,684	2,062	2,296
Discharge (cfs)	S-67	0	0	0	0	0	0	0	0	67	310	288
DO (mg/L) ³	Phase I river channel	5.5	5.5	5.7	5.2	5.3	4.3	4.6	4.7	4.3	3.3	2.8
Mean depth (feet) ⁴	Phase I floodplain	0.12	0.13	0.16	0.22	0.22	0.36	0.52	0.56	0.64	0.75	0.80

¹ Seven-day average of weighted daily means through Sunday midnight.

² S-65A discharge combines S-65A with auxiliary structures; S-65D discharge combines discharge at S-65D, S-65DX1, and S-65DX2; S-65D stage averages stage at S-65D and S-65DX1; S-65E discharge combines S-65E and S-65EX1.

³ DO is the average for sondes at PC62 and PC33.

⁴ 1-day spatial average from South Florida Water Depth Assessment Tool (SFWDAT).

DATA ARE PROVISIONAL; N/A indicates that data were not available.

KCOL Hydrographs (through Sunday midnight)




Figure 1.




Figure 2.




Figure 3.




Figure 4.




Figure 5.




Figure 6.




Figure 7.




Figure 8. Phase I area floodplain water depths for this week, one month ago, and two months ago. Note that the WDAT color-coding has been modified to accommodate greater water depths; these maps are not directly comparable to Kissimmee Basin WDAT maps published prior to January 16, 2012.




Figure 9. Mean water depth at stage recorders in the northern Phase I, southern Phase I, northern Phase II/III, and southern Phase II/III areas in relation to the S-65A discharge and S-65D headwater stage.




Figure 10. Mean daily dissolved oxygen, discharge, temperature and rainfall in the Phase I river channel.

Water Management Recommendations

Kissimmee Basin Adaptive Recommendations and Operational Actions

Recommendation Date	Recommendation	Purpose	Outcome	Source	Report Date
11/26/2018	No new recommendations.		N/A		11/27/2018
11/19/2018	No new recommendations.		N/A		11/20/2018
11/12/2018	No new recommendations.		N/A		11/13/2018
11/2/2018	Reduce S-65/S-65A discharge to approximately 250 cfs.	To conserve stage in Lake Kissimmee.	Implemented	SFWMD Water Mgt/KB Ops	11/6/2018
10/30/2018	No new recommendations.		N/A		10/30/2018
10/22/2018	Reduce S-65/S-65A discharge to approximately 300 cfs (minimum discharge) in one step of approximately 1100 cfs today.	Reduce rate of stage decline in lakes Kissimmee-Cypress-Hatchineha	Implemented	SFWMD Water Mgt/KB Ops	10/23/2018
10/16/2018	No new recommendations.		N/A		10/16/2018
10/9/2018	No new recommendations.		N/A		10/9/2018
10/2/2018	No new recommendations.		N/A		10/2/2018
9/25/2018	No new recommendations.		N/A		9/25/2018
9/18/2018	No new recommendations.		N/A		9/18/2018
9/11/2018	No new recommendations.		N/A		9/11/2018
9/4/2018	No new recommendations.		N/A		9/4/2018
8/28/2018	No new recommendations.		N/A		8/28/2018
8/21/2018	No new recommendations.		N/A		8/21/2018
8/14/2018	No new recommendations.		N/A		8/14/2018
8/7/2018	No new recommendations.		N/A		8/7/2018
7/23/2018-7/24/2018	Increase discharge from 1400 cfs to 3000 cfs, then 3200 cfs and 3500 cfs.	For flood control in Lake Kissimmee.	Implemented	SFWMD Water Mgt/KB Ops	7/31/2018
7/19/2018	Follow Revised (X2) 2018 Wet Season Discharge Plan to the extent possible, including 50 foot stage threshold and 0.5 foot flood control buffer.	To the extent possible, maintain sufficient discharge to keep areas under snail kites nests in Pool D hydrated until nests fledge, while avoiding large increases in discharge that might flood the nests.	N/A	KB Ops	7/24/2018
7/13/2018	Maintain at least 1400 cfs at S-65A while Lake Kissimmee stage is above 50 feet. (See revised 2018 discharge plan).	To the extent possible, maintain sufficient discharge to keep areas under snail nest kites in Pool D hydrated until nests fledge.	N/A	KB Ops	7/17/2018
7/13/2018	Reduce S-65/S-65A discharge by 600 cfs/day until 1400 cfs is reached. (See revised 2018 discharge plan, below).	Reach 1400 cfs faster to help stabilize Lake Kissimmee stage.	Implemented	SFWMD Water Mgt/KB Ops	7/17/2018
7/9/2018	Increase S-65/S-65A discharge by 300 cfs if needed.	Stabilize Lake Kissimmee stage.	N/A	SFWMD Water Mgt/KB Ops	7/10/2018
7/8/2018	Increase S-65/S-65A discharge by 900 cfs today in 3 increments of 300 cfs each.	Stabilize Lake Kissimmee stage.	Implemented	KB Ops	7/10/2018
7/5/2018	Increase S-65/S-65A discharge by 300 cfs/day (double the prescribed rate of increase) Thursday through Sunday .	Stabilize Lake Kissimmee stage.	Implemented	SFWMD Water Mgt	7/10/2018
7/2/2018	Increase S-65/S-65A discharge by 150 cfs/day (double the prescribed rate of increase).	Stabilize Lake Kissimmee stage.	Implemented	SFWMD Water Mgt/KB Ops	7/10/2018
6/30/2018	Increase S-65/S-65A discharge as slowly as feasible	Slow stage ascension in Kissimmee-Cypress-Hatchineha	Implemented	KB Ops/SFWMD Water Mgt	7/3/2018
6/28/2018	Continue to reduce discharge at S-65/S-65A as slowly as feasible.	Prevent stage decline in Kissimmee-Cypress-Hatchineha.	Implemented	KB Ops/SFWMD Water Mgt	7/3/2018
6/21/2018	Reduce discharge at S-65/S-65A as slowly as feasible.	Prevent stage decline in Kissimmee-Cypress-Hatchineha.	Implemented	KB Ops/SFWMD Water Mgt	6/26/2018
6/15/2018	Reduce S-65A discharge by 150-300 cfs over the weekend.	Slow or stop DO decline in Kissimmee River.	Implemented	KB Ops	6/19/2018
6/12/2018	No new recommendations.		N/A		6/12/2018
6/5/2018	No new recommendations.		N/A		6/5/2018
5/29/2018	Begin implementation of the 2018 Wet Season Discharge Plan for S-65/S-65A on June 1 (see figure).	Provide variable flow from S-65/S-65A to balance Kissimmee River and Headwaters Lakes objectives including Kissimmee River floodplain inundation, moderated rates of change in discharge, and constrained rate of stage rise in the lakes.	Planned	KB Ops/SFWMD Water Mgt/FWC/FWS	5/29/2018




Figure 11. The 2018 Wet Season Discharge Plan for S-65/S-65A.




Figure 12. Interim operations schedule for S-65. The discharge schedule shown to the right has not been used in recent years.




Figure 13. The Kissimmee Basin.

LAKE OKEECHOBEE

According to the USACE web site, Lake Okeechobee stage is at 13.22 feet NGVD for the period ending at midnight on November 26, 2018. This value is based on the use of four interior lake stations (L001, L005, L006 and LZ40) and four perimeter stations (S-308, S-352, S-4 and S-133). The Lake is now 0.61 feet lower than it was a month ago and 3.04 feet lower than a year ago when runoff from Hurricane Irma caused extreme high lake stages (Figure 1). The Lake is in the Base-Flow sub-band (Figure 2). The November 26 lake stage was the lowest for this time of year since 2010 (Figure 3). According to RAINДАР, 0.15 inches of rain fell over the Lake during the week November 20, 2018 – November 26, 2018. The northern portion of the watershed received more rainfall, between 0.25 and 1.5 inches, while much of the remainder of the watershed received similar to less, between 0 and 0.25 inches (Figure 4). Average daily inflows (minus rainfall) to the Lake were similar to the previous week, going from 367 cfs to 345 cfs.

Total outflows (minus evapotranspiration) decreased slightly from the previous week, going from 2,299 average daily cfs the previous week to 2,241 cfs this past week (Table 1). The decreases in outflows were primarily in discharges south through the S350 structures which went from 1,167 cfs the previous week to 1,126 cfs this past week. Outflows via the S-77 decreased slightly from 957 cfs the previous week to 920 average daily cfs this past week while outflows through the L8 at Canal Point increased slightly from 180 cfs the previous week to 195 cfs this past week. Flows through the S-308 have been passive flows through the navigation lock and/or structure and averaged 0 cfs. The corrected average daily evapotranspiration value for the week based on the L006 weather platform solar radiation decreased from 0.07 inches last week to 0.06 inches this week.

Total lake inflows and outflows for the past week are detailed in Table 1, as well as the approximate change in lake stage from each major structure's total flows over the period. Figure 5 shows the combined average daily cfs for inflows and outflows for the Lake over the past eight weeks. These data are provisional and are subject to change.

Water quality samples from 17 nearshore and pelagic stations on November 6 – November 7 showed continued decreases in average total phosphorus (TP) and a leveling off of turbidity values from January peaks (due to Hurricane Irma sediment disruptions) (Figure 6). Average monthly TP and turbidity values are close to pre-hurricane Irma levels.

Quarterly sampling of submersed aquatic vegetation (SAV) revealed some recovery of plants in sheltered areas since the August 2018 sampling, however, coverage remains very sparse (Figure 7). The November 2018 sampling found four of nine sites in Fisheating Bay with sparse SAV compared to only one site with SAV in August 2018. Similarly, two of nine sites around King's Bar had sparse SAV coverage in November, while only one site had SAV in August. No recovery has occurred in the southern or western nearshore regions. For comparison, in August 2017 (pre-Hurricane Irma) five of the nine sites around King's Bar and seven of the nine sites in Fisheating Bay had SAV, ten of which had moderate to dense coverage. Additionally, six of the fifteen sites in the southern region had SAV prior to Irma.

The most recent satellite imagery (November 21) using the cyanobacteria monitoring product derived from NOAA's analysis of EUMETSAT's OLCI satellite sensor showed a decrease in algal bloom potential in the south region but the southwest shore remains at medium risk (Figure 8).

Water Management Recommendations

Lake Okeechobee stage is 13.22 feet NGVD, falling 0.13 feet from the previous week and 0.61 feet over the past 30 days. Lake stages are now the lowest they have been for this time of year since 2010 and are 1.28 feet below the bottom of the preferred ecological envelope, which varies seasonally from 12.5 – 15.5 feet NGVD. However, given potential for heavy rainfall associated with El Niño conditions

this winter and the poor condition of SAV and EAV in the nearshore zone, these lower lake stages are ideal. Recovery of vegetation in the nearshore zone from Hurricane Irma impacts and 2016 El Niño-associated rainfall will require lake stages in the lower portion of the ecological envelope or lower for extended periods, so efforts to prepare for such an event will help speed the rebound of this important community.

Table 1. Average daily inflows and outflows and the approximate depth equivalents on Lake Okeechobee for various structures.

INFLOWS	Previous Week Avg Daily cfs	Avg Daily Inflow cfs	Equivalent Depth Week Total (in)	OUTFLOWS	Previous Week Avg Daily cfs	Avg Daily Outflow cfs	Equivalent Depth Week Total (in)
S65E & S65EX1	336	327	0.1	S77	957	920	0.4
S71 & 72	0	0	0.0	S308	-5	0	0.0
S84 & 84X	0	0	0.0	S351	551	470	0.2
Fisheating Creek	31	18	0.0	S352	401	470	0.2
S154	0	0	0.0	S354	215	186	0.1
S191	0	0	0.0	L8 Outflow	180	195	0.1
S133 P	0	0	0.0	ET	1073	1000	0.4
S127 P	0	0	0.0	Total	3372	3241	1.4
S129 P	0	0	0.0				
S131 P	0	0	0.0				
S135 P	0	0	0.0				
S2 P	0	0	0.0				
S3 P	0	0	0.0				
S4 P	0	0	0.0				
L8 Backflow							
Rainfall	794	357	0.2				
Total	1161	702	0.3				

PROVISIONAL DATA




Figure 1. Water depth estimates on Lake Okeechobee based on the South Florida Water Depth Assessment Tool.

Lake Okeechobee Water Level History and Projected Stages




Figure 2. Recent Lake Okeechobee stage and releases, with projected stages based on a dynamic position analysis.

Lake Okeechobee Water Level Comparison




Figure 3. Annual stage hydrographs for Lake Okeechobee from 2010 – 2018.

SFWMD PROVISIONAL RAINDAR 7-DAY BASIN RAINFALL ESTIMATES

FROM: 0615 EST, 11/20/2018 THROUGH: 0615 EST, 11/27/2018




Figure 4. Rainfall estimates by basin.




Figure 5. Major inflows and outflows of Lake Okeechobee, including the S350 structures designated as South. The L8 canal flows through culvert 10A are included as outflows when positive, and as inflows when backflowing into the lake. All inflows and outflows are shown as positive and negative, respectively, for visual purposes.




Figure 6. Water quality sampling locations on Lake Okeechobee in the nearshore and pelagic regions (right) and average turbidity (top graph) and total phosphorus (bottom graph) monthly values for each region from January 2015 – November 2018. Note: The FEBIN and FEBOUT stations were not included due to intermittent sampling.




Figure 7. Submerged Aquatic Vegetation (SAV) sampled in May 2018, August 2018, and November 2018 at the 44 sentinel grid cells.




Figure 8. Potential for cyanobacterial blooms on Lake Okeechobee based on NOAA's harmful algal bloom monitoring system derived from Copernicus Sentinel-3 OLCI data from EUMETSAT. Gray indicates cloud cover. All data are experimental and unvalidated at this point in product development.

Estuaries

St. Lucie Estuary:

Last week total inflow to the St. Lucie Estuary averaged approximately 157 cfs (Figures 1 and 2) and last month inflow averaged about 260 cfs. Last week's provisional averaged inflows from the tidal basin and the structures are shown in Table 1.

Table 1. Weekly average inflows (data are provisional).

Location	Flow (cfs)
Tidal Basin Inflow	95
S-80	0
S-308	0
S-49 on C-24	0
S-97 on C-23	0
Gordy Rd. structure on Ten Mile Creek	62

Over the past week, salinity increased throughout the estuary (Table 2, Figures 3 and 4). The seven-day moving average salinity of the water column (an average of the surface and bottom salinity) at the US1 Bridge is 24.0. Salinity conditions in the middle estuary are within the good range for adult eastern oysters (Figure 3).

Table 2. Seven-day average salinity at three monitoring sites in the St. Lucie Estuary. Current average is in bold face type, previous average in parentheses. The envelope reflects the preferred salinity range for adult eastern oysters (*Crassostrea virginica*) in the middle estuary.

Sampling Site	Surface	Bottom	Envelope
HR1 (North Fork)	21.3 (18.3)	23.8 (21.4)	NA ¹
US1 Bridge	24.3 (22.2)	24.4 (22.8)	10.0-26.0
A1A Bridge	30.3 (28.2)	31.2 (29.6)	NA ¹

¹Envelope not applicable and ²Not Reporting.

Caloosahatchee Estuary:

Last week total inflow to the Caloosahatchee Estuary averaged approximately 1,272 cfs (Figures 5 and 6) and last month inflow averaged about 1,262 cfs. Last week's provisional averaged inflows from the structures are shown in Table 3.

Table 3. Weekly average inflows (data is provisional).

Location	Flow (cfs)
S-77	897
S-78	685
S-79	1,150
Tidal Basin Inflow	122

Over the past week in the estuary, surface salinity remained about the same to Val I75 and increased downstream (Table 4, Figures 7 & 8). The seven-day average salinity values are estimated to be within the good range for adult eastern oysters at Cape Coral and at Shell Point (Figure 9). Salinity values were not available at Sanibel. The 30-day moving average surface salinity is 1.9 at Val I-75 and 7.1 at Ft. Myers. Salinity conditions between Val I-75 and Ft. Myers are good for tape grass.

Table 4. Seven-day average salinity at six monitoring stations in the Caloosahatchee Estuary. Current average is in bold face type, previous average in parentheses. The envelope reflects the preferred salinity range for tape grass (*Vallisneria americana*) at Val I-75 and for adult eastern oysters (*Crassostrea virginica*) elsewhere.

Sampling Site	Surface	Bottom	Envelope
S-79 (Franklin Lock)	1.8 (2.0)	1.9 (1.9)	NA ¹
Val I75	2.6 (2.1)	4.2 (3.9)	0.0-5.0 ²
Ft. Myers Yacht Basin	7.8 (6.8)	9.7 (11.0)	NA
Cape Coral	15.4 (14.6)	17.8 (17.4)	10.0-30.0
Shell Point	25.6 (24.1)	25.5 (23.8)	10.0-30.0
Sanibel	NR ³ (NR)	NR (NR)	10.0-30.0

¹Envelope not applicable, ²Envelope is based on a 30-day average, and ³Not Reporting.

Forecast of surface salinity (Table 5 and Figure 10) at Val I-75 for the next two weeks using the autoregression model (Qiu and Wan, 2013) coupled with a linear reservoir model for the tidal basin predicts daily salinity ranging from 1.7 to 4.4 at the end of the next two weeks for pulse release at S-79 ranging from 0 to 650 cfs and Tidal Basin inflows of 165 cfs.

Table 5. Predicted salinity at Val I-75 at the end of forecast period

Scenario	Q79 (cfs)	TB runoff (cfs)	Daily salinity	30 day mean
A	0	165	4.4	2.8
B	300	165	3.9	2.7
C	375	165	3.3	2.6
D	450	165	2.9	2.4
E	650	165	1.7	2.1

Red tide

The Florida Fish and Wildlife Research Institute reported on November 21, 2018, that *Karenia brevis*, the Florida red tide dinoflagellate, was observed at background to high concentrations in 28 samples collected from or offshore of Lee County. *Karenia brevis*, the Florida red tide dinoflagellate, was observed at background to very low concentrations in 15 samples collected from Palm Beach County and in two samples collected offshore of Broward County. A fish kill was reported in Palm Beach County. Respiratory irritation was reported in Palm Beach County and in Lee County.

Water Management Recommendations

Lake stage is in the Base Flow sub-band of 2008 LORS. Tributary hydrological conditions are dry. The 2008 LORS recommends up to 450 cfs at S-79 and up to 200 cfs at S-80. Given the current estuarine conditions, there are no ecological benefits associated with freshwater releases from Lake Okeechobee.




Figure 1. Basins, water control structures, and salinity monitoring for the St. Lucie Estuary.




Figure 2. Total daily inflows from Lake Okeechobee and runoff from the C-44, C-23, C-24, Ten Mile Creek, and tidal basins into the St. Lucie Estuary.




Figure 3. Seven-day mean salinity of the water column at the US1 Bridge.




Figure 4. Daily mean salinity at the A1A, US1 and estimated HR1 stations.




Figure 5. Basins, water control structures, and salinity monitoring for the Caloosahatchee Estuary.




Figure 6. Total daily inflows from Lake Okeechobee, runoff from the C-43 basin, and tributaries in the tidal basin into the Caloosahatchee River Estuary.




Figure 7. Daily mean flows at S-79 and salinity at upper estuary monitoring stations.




Figure 8. Daily mean flows at S-79 and salinity at lower estuary stations.




Figure 9. Seven-day mean salinity at Cape Coral, Shell Point, and Sanibel monitoring stations.





Figure 10. Forecasted Val I-75 surface salinity assuming no pulse release at S79.


EVERGLADES

At the gauge locations monitored for this report stages within the WCAs fell 0.03 feet on average over the last week. The most extreme individual gauge changes within the WCAs ranged from +0.07 feet (WCA-2B) to -0.14 feet (WCA-2A). Pan evaporation was estimated at 1.12 inches this week.



Everglades Region	Rainfall (Inches)	Stage Change (feet)
WCA-1	0.26	-0.02
WCA-2A	0.07	-0.14
WCA-2B	0.02	+0.07
WCA-3A	0.34	-0.02
WCA-3B	0.28	-0.07
ENP	0.22	-0.14



Regulation Schedules: Gauge 1-8C is 0.41 feet below the Zone A2 regulation line. The three-gauge average remains 0.22 below the canal stage. S11B Headwater stage is 0.58 feet above the Zone A regulation line and is receding parallel to that line. WCA-3A three-gauge average stage is 0.44 feet below the Zone D regulation line and trends away from the rising regulation line. WCA-3A stage at gauge 62 (northwest corner) is 1.12 feet below the Lower Schedule.






Water Depths and Changes: The WDAT tool for spatial interpolation of depth monthly snapshots indicate drying conditions. Regions with depths from 0.5 feet to 0.0 feet have increased across most of WCA-3A North. In the extreme northeast region of WCA-3A and northern WCA-1 the model also indicates expanding regions with water depths significantly below ground. WDAT difference output indicates that water levels dropped across all of the Everglades over the last month, yet the rate of change is moderate. The S-150 structure continues to discharge into northeastern WCA-3A over the last month, now near 300 cubic feet per second. In the “1 Year” inset we see the comparison between current depth conditions and post Hurricane Irma’s (9/10/17) impact on water depths.




Taylor Slough Water Levels: An average of 0.2 inches of rain fell on Taylor Slough and Florida Bay this past week allowing stages to continue decreasing by an average of 0.03 feet last week. Water depths averaged 0.91 feet across Taylor Slough with northern Taylor Slough being 3 inches higher than the historical average for this time of year, but southwestern Taylor Slough is 0.3 inches lower than average.

Florida Bay Salinities: Salinities averaged an increase of 4 psu this past week and range from 22 psu in the north to 36 psu in the west. Conditions in the west-central Florida Bay nearshore area are up to 11 psu higher than their historical averages for this time of year which is undesirable at the start the dry season.





Florida Bay MFL: Mangrove zone daily average salinity remained at 0.5 psu for most of the week but jumped to 2.4 on Sunday. The 30-day moving average stayed at 0.5 psu. The weekly cumulative flow from the five creeks denoted by yellow stars on the map totaled about -6,850 acre-feet last week as most of the week experienced upstream flows. The 365-day moving sum of flow from the five creeks has been dropping rapidly with an additional decrease of 12,000 acre-feet over the last week to end at 233,103 acre-feet (less than the long-term average of 257,628 acre-feet). Creek flow is provisional data from the USGS and is highly variable.



Water Management Recommendations

Water management that protects peat soils, especially in WCA-3A North, as the dry season becomes established has increasing ecological benefit as unusually dry conditions pervade the WCA-3A basin. Any water not available to protect the peat soils in WCA-3A North, would be ecologically beneficial to Holeyland and Rotenburger WMA. According to the WDAT modeling, depths in the northern portion of WCA-1 that are below ground level have expanded significantly over the last month. This historically dry area would continue to benefit from hydration as the 3-gauge average stage is now 0.63 feet below the Zone A2 regulation line but has remained steady over the last several weeks. Incremental change in the rate of structure flows (i.e., when changing flow rates from 0 cfs to 1,000 cfs, make 500 cfs adjustment per week) to the WCAs is more ecologically sensitive than abrupt rate changes. More specific recommendations appear in the summary table below. The red text represents new or modified information or recommendations.

SFWMD Everglades Ecological Recommendations, November 27th, 2018 (red is new)

Area	Weekly change	Recommendation	Reasons
WCA-1	Stage decreased by 0.02'	Maintain depths at regulation schedule. Moderate recession rates to the extent possible.	Protect upstream/downstream habitat and wildlife.
WCA-2A	Stage decreased by 0.14'	Maintain depths at regulation schedule. Moderate recession rates to the extent possible.	Protect upstream/downstream habitat and wildlife.
WCA-2B	Stage increased by 0.07'	Maintain depths at regulation schedule. Moderate recession rates to the extent possible.	Protect upstream/downstream habitat and wildlife.
WCA-3A NE	Stage decreased by 0.06'	Maintain depths at regulation schedule. Moderate recession rates to the extent possible.	Protect habitat including <u>peat soil</u> development, tree islands and wildlife. Hydration provided to this area has high ecological value due to unseasonably dry conditions.
WCA-3A NW	Stage decreased by 0.01'	Maintain depths at regulation schedule. Moderate recession rates to the extent possible.	
Central WCA-3A S	Stage remained unchanged	Maintain depths at regulation schedule. Moderate recession rates to the extent possible.	Protect habitat including peat soil development, <u>tree islands</u> and wildlife.
Southern WCA-3A S	Stage decreased by 0.02'		
WCA-3B	Stage decreased by 0.07'	Maintain depths at temporary regulation schedule. Moderate recession rates to the extent possible.	Protect upstream/downstream habitat and wildlife.
ENP-SRS	Stage decreased by 0.14'	Make discharges to the Park according to the 2012 WCP rainfall plan.	Protect upstream/downstream habitat and wildlife.
Taylor Slough	Stage changes ranged from -0.05' to -0.00'	Move water southward as possible	When available, provide freshwater buffer for downstream conditions. Decrease potential for high phosphorus input to ENP.
FB- Salinity	Salinity changes ranged -2.5 to +17.2 psu	Move water southward as possible	When available, provide freshwater to maintain low salinity buffer and promote water movement.