

Disclaimer: Information contained in the report addresses environmental conditions only and is not the official South Florida Water Management District operations recommendation or decision.

M E M O R A N D U M

TO: John Mitnik, Chief, Operations, Engineering and Construction Bureau
Paul Linton, Chief, Operations Section

FROM: SFWMD Staff Environmental Advisory Team

DATE: May 7, 2019

SUBJECT: Weekly Environmental Conditions for Systems Operations

Summary

Weather Conditions and Forecast

A weak cold front entering the far northern portion of the District this morning should slowly sag southward to a position not far to the north of Lake Okeechobee by this evening. To the south of the frontal boundary a continued rich supply of moisture over a deep layer, enhanced atmospheric instability, and daytime heating should result in yet another day of active rains across parts of the District this afternoon to evening. However, the spatial distribution of today's rains should have some noticeable differences compared that observed yesterday. A change in the direction of low-level winds from the west to the northeast in response to high pressure over the western Atlantic building across the front should mean a progression of the east coast sea breeze farther inland by afternoon, which would tend to lessen the amount of rain along and near the east coast compared to the last couple of days. How much of a decrease over this region is unknown now due to a significant disagreement in the models with regard to the likely amount of rain over this region, and an update may be necessary by early afternoon to properly modify the forecast based on new data and trends. The west and east coast sea breezes are forecast to meet up over the interior of the peninsula later this afternoon and result in a large concentration of rains from the south-central and possibly southwestern interior to around or west of Lake Okeechobee. The weaker steering flow across the area would suggest that some significant rainfall totals over this region are possible, even though the favorable large-scale dynamics from yesterday would be lacking. Overall, the total District rainfall today is predicted to be lower than yesterday's, with the focus of this afternoon's activity shifted perhaps significantly to the west.

Kissimmee

Tuesday morning stages were 55.6 feet NGVD (0.4 feet below schedule) in East Lake Toho, 52.5 feet NGVD (0.5 feet below schedule) in Toho, and 49.4 feet NGVD (0.4 feet below schedule) in Kissimmee-Cypress-Hatchineha; headwater stages were 46.3 feet NGVD at S-65A and 25.9 feet NGVD at S-65D. Tuesday morning discharges were 1216 cfs at S-65, 975 cfs at S-65A, 452 cfs at S-65D and 404 cfs at S-65E. Dissolved oxygen concentration in the Kissimmee River averaged 5.4 mg/L for the week. Kissimmee River mean floodplain depth on Sunday was 0.11 feet. Recommendations (5/6/2019): Due to the rainfall, increase S-65A to 1000 cfs today 5/6 in two increments, and increase flow at S-65 accordingly. We will reassess the rise in KCH stage 5/7 AM. Purpose is to try to keep S65-A discharge at or below 1000 cfs while holding the reversal in KCH to less than about 0.4 ft.

Lake Okeechobee

Lake Okeechobee stage is 11.23 feet NGVD, decreasing 0.01 feet from the previous week and experiencing a reversal due to recent rains. The Lake dropped into the Beneficial Use sub-band on March 7, 2019 and is now 0.37 feet above the Water Shortage sub-band. The lake remains below the bottom of the ecological envelope (currently 1.26 feet below), which varies seasonally from 12.5 – 15.5

feet NGVD. Given the continued potential for above average rainfall (associated with a weak El Niño and the upcoming wet season) over the next few months, and the poor condition of SAV and EAV in the nearshore zone, these lower lake stages are ideal for vegetation recovery. However, low stages will reduce habitat for fish and wildlife in the near-term and encourage spread of invasive vegetation in the upper marsh. Wading bird foraging numbers have decreased as suitable habitat is limited to small ponded areas in the marsh and along the outer marsh-nearshore interface. Estimated algal bloom potential using satellite imagery suggests medium bloom risk in the north of the lake and along the western shore, particularly within Fish-eating Bay and along the edge of Indian Prairie.

Estuaries

Total inflow to the St. Lucie Estuary averaged 556 cfs over the past week with no inflow coming from Lake Okeechobee. Over the past week, the seven-day average salinities were little changed in the estuary but there was a dip at the end of the week due to rainfall event. The seven-day average salinity at the US1 Bridge is within the good range for adult eastern oysters. Total inflow to the Caloosahatchee Estuary averaged 1,227 cfs over the past week with 334 cfs coming from the Lake. Over the past week, salinity increased in the estuary. The 30-day moving average surface salinity is 0.6 at Val I-75 and 5.6 at Ft. Myers. Salinity conditions between Val I-75 and Ft. Myers are good for tape grass. Salinity conditions are in the good range for adult eastern at Cape Coral and Shell Point and in the fair range at Sanibel.

Stormwater Treatment Areas

Over the past week, approximately 4,900 acre-feet of Lake water was delivered to the STAs to maintain target stages. The total amount of Lake releases sent to the STAs/FEBs in WY2020 (since May 1, 2019) is approximately 4,400 acre-feet. The total amount of inflows to the STAs in WY2020 is approximately 10,200 acre-feet. Most STA cells are at or near target depths except STA-5/6 cells which are below target. STA-1W Northern Flow-way is offline for STA-1W Expansion project construction activities, STA-1E Western Flow-way is offline for levee repairs in the West Distribution Cell, and STA-5/6 Flow-ways 2 and 3 are offline for the Restoration Strategies project to grade non-effective treatment areas. Operational restrictions are in place in STA-5/6 Flow-ways 1 and 4 to facilitate the Restoration Strategies grading project in Flow-ways 2 and 3. The nest of an Endangered Species Act (ESA) protected species has been observed in STA-1E and the nests of Migratory Bird Treaty Act (MBTA) protected species have been observed in STA-5/6. It is recommended that no Lake releases be sent to the STAs this week.

Everglades

A reversal in stage conditions occurred across the Everglades over the last week, most dramatically in portions of WCA-3A. Ecological areas of concern are the foraging conditions in WCA-2A and WCA-3A South and peat soils / fire risk in northern WCA-3A. Water depths in Taylor Slough and the ENP panhandle continue to decrease as is typical for this time of year. Salinities in Florida Bay and the mangrove zone increased over the week. Over the last several weeks the number of white ibis nesting in WCA-3A's Alley North Colony decrease by approximately 6000 nests, now that island contains approximately 2,000 nests; an increase in the number of wading birds nesting in WCA-1 was noted. Large flocks of wading birds are foraging in western WCA-2A.

Supporting Information

KISSIMMEE BASIN

Kissimmee Basin Rainfall

The Upper Kissimmee Basin received 1.45 inches of rainfall in the past week and the Lower Basin received 0.91 inches (SFWMD Daily Rainfall Report 5/6/2018).

Upper Kissimmee Basin

Stages and departures in the Kissimmee Chain of Lakes (KCOL) are shown in **Table 1**. KCOL stage hydrographs with respective regulation schedules and rainfall are shown in Figures 1-7.

Table 1. Average discharge (cfs) for the preceding seven days, one-day stage (feet NGVD), and departures from KCOL flood regulation (R) or temporary schedules (T, A, or S). Provisional, real-time data are from SFWMD.
Report Date: 5/7/2019

Water Body	Structure	7-day Average Discharge (cfs) ¹	Stage Monitoring Site ²	Lake Stage (feet)	Schedule Type ³	Schedule Stage (feet)	Daily Departure (feet)						
							5/5/19	4/28/19	4/21/19	4/14/19	4/7/19	3/31/19	3/24/19
Lakes Hart and Mary Jane	S-62	14	LKMJ	60.0	R	60.0	0.0	-0.2	-0.2	-0.3	-0.4	-0.3	-0.4
Lakes Myrtle, Preston, and Joel	S-57	0	S-57	60.3	R	60.3	0.0	-0.1	-0.1	0.0	0.0	0.0	0.0
Alligator Chain	S-60	70	ALLI	62.8	R	62.7	0.1	0.0	0.0	-0.1	0.0	0.0	-0.1
Lake Gentry	S-63	105	LKGT	60.3	R	60.2	0.1	0.0	0.0	0.0	0.0	0.0	0.0
East Lake Toho	S-59	51	TOHOE	55.7	R	56.1	-0.4	-0.7	-0.8	-1.0	-1.1	-1.2	-1.4
Lake Toho	S-61	219	TOHOW, S-61	52.6	R	53.1	-0.5	-0.7	-0.8	-1.0	-1.1	-1.2	-1.4
Lakes Kissimmee, Cypress, and Hatchineha	S-65	428	KUB011, LKISSB	49.4	R	49.9	-0.5	-0.6	-0.7	-0.9	-0.9	-1.1	-0.9

¹ Seven-day average of weighted daily means through midnight.

² Names of in-lake monitoring sites and structures used to determine lake stage; if more than one site is listed, an average is reported.

³ A = projected ascension line, R = USACE regulation schedule, S = temporary recession target line, T = temporary schedule, N/A = not applicable or data not available.
DATA ARE PROVISIONAL

Lower Kissimmee Basin

Discharges at Lower Basin structures are shown in Table 2. SFWDAT depth maps for the Phase I restoration area are shown in Figure 8. Kissimmee River floodplain stages at selected stations are shown in Figure 9.

Table 2. One-day and seven-day averages of discharge at S-65x structures, of dissolved oxygen concentration in the Phase I area river channel, and water depth in the Phase I area floodplain. Data are provisional real-time
Report Date: 5/7/2019

Metric	Location	1-Day Average		Average for the Preceding 7-Days ¹								3/3/19
		5/5/2019	5/5/19	4/28/19	4/21/19	4/14/19	4/7/19	3/31/19	3/24/19	3/17/19	3/10/19	
Discharge (cfs)	S-65	552	428	438	525	710	434	452	833	529	513	1,368
Discharge (cfs)	S-65A ²	415	314	314	400	559	334	353	699	420	409	1,190
Discharge (cfs)	S-65D ²	445	403	466	584	703	367	563	859	505	1,103	2,310
Headwater Stage (feet NGVD)	S-65D ²	25.76	25.81	25.76	25.78	25.77	25.73	25.76	25.77	25.78	25.72	25.76
Discharge (cfs)	S-65E ²	403	351	441	563	679	330	539	855	497	1,026	2,167
Discharge (cfs)	S-67	104	68	107	110	106	0	9	162	0	51	30
DO (mg/L) ³	Phase I river channel	5.1	5.4	5.6	6.0	6.3	6.9	7.4	6.7	5.9	5.6	4.1
Mean depth (feet) ⁴	Phase I floodplain	0.11	0.10	0.12	0.16	0.18	0.16	0.21	0.34	0.29	0.43	0.86

¹ Seven-day average of weighted daily means through Sunday midnight.

² S-65A discharge combines S-65A with auxiliary structures; S-65D discharge combines discharge at S-65D, S-65DX1, and S-65DX2; S-65D stage averages stage at S-65D and S-65DX1; S-65E discharge combines S-65E and S-65EX1.

³ DO is the average for sondes at PC62 and PC33.

⁴ 1-day spatial average from South Florida Water Depth Assessment Tool (SFWDAT).

DATA ARE PROVISIONAL; N/A indicates that data were not available.

KCOL Hydrographs (through Sunday midnight)




Figure 1.




Figure 2.




Figure 3.




Figure 4.




Figure 5.




Figure 6.




Figure 7.




Figure 8. Phase I area floodplain water depths for this week, one month ago, and two months ago. Note that the WDAT color-coding has been modified to accommodate greater water depths; these maps are not directly comparable to Kissimmee Basin WDAT maps published prior to January 16, 2012.





Figure 9. Mean water depth at stage recorders in the northern Phase I, southern Phase I, northern Phase II/III, and southern Phase II/III areas in relation to the S-65A discharge and S-65D headwater stage.




Report Date: 5/7/2019; data are through: 5/5/2019.

Figure 10. Mean daily dissolved oxygen, discharge, temperature and rainfall in the Phase I river channel.

Water Management Recommendations

Recommendation Date	Recommendation	Purpose	Outcome	Source	Report Date
5/6/2019	Due to the rainfall, increase S65-A to 1000 cfs today in two increments and increase flow at S-65 accordingly. We will reassess the rise in KCH stage tomorrow 5/7.	Short-term goals: try to keep S65-A discharge at or below 1000 cfs for KR fish sampling this week and next, while keeping the reversal in KCH less than about 0.4 ft.	Implemented	KB Ops	5/7/2019
4/29/2019	No new recommendations.		N/A		4/30/2019
4/23/2019	No new recommendations.		N/A		4/23/2019
4/15/2019	No new recommendations.		N/A		4/16/2019
4/8/2019	No new recommendations.		N/A		4/9/2019
4/1/2019	No new recommendations.		N/A		4/2/2019
3/25/2019	No new recommendations.		N/A		3/26/2019
3/18/2019	No new recommendations.		N/A		3/19/2019
3/11/2019	No new recommendations.		N/A		3/12/2019
3/4/2019	No new recommendations.		N/A		3/5/2019
2/26/2019	No new recommendations.		N/A		2/26/2019
2/19/2019	No new recommendations.		N/A		2/19/2019
2/10/2019	Increase discharge at S-65 by 600 cfs.	To compensate for increased inflow and rain forecast for Tuesday.	Implemented	KB Ops/SFWMD Water Mgt	2/12/2019
2/4/2019	Increase discharge at S-65/S-65A to begin reducing KCH stage to reach 50.75 ft on 2/15/2019.	Reduce to the stage at which the seasonal recession will begin.	Implemented	KB Ops/SFWMD Water Mgt	2/5/2019
1/26/2019	Increase S65A discharge by a total of 350 cfs today, which will put S65A at 1,400 cfs. Continue to increase discharge as needed.	Moderate or stop the rise in Lake KCH preemptively before forecast rainfall and provide capacity at S65A for S65A basin runoff.	Implemented	SFWMD Water Mgt/KB Ops	1/29/2019
1/22/2019	No new recommendations.		N/A		1/22/2019
	Begin recessions on Lake Toho and East Lake Toho on Jan 15, with a continuous recession to the regulation dry season low (52.0 ft on Toho; 55.0 ft on East Lake) on May 31. The lines are represented graphically in the Dry Season Operations slides.				
1/15/2019	Tentatively plan on a recession in Kissimmee-Cypress-Hatchineha starting on February 15 with a continuous recession to the dry season low (49 ft) on May 31. A provisional diagram is included in the Dry Season Operations slides; however, starting stage may change depending on conditions.	Slow recession rates in East Toho, Toho, and KCH to benefit fish and wildlife; as possible limit flow volume at S-65D to facilitate KRR construction.	N/A	KB Ops	1/15/2019
	Discharge and reversal guidelines are provided in the Dry Season Operations slides.				
1/4/2019	Discontinue 54 foot stage reduction target in Lake Toho.	Lake Kissimmee has already risen by ~1.5 ft.	Implemented	SFWMD Water Mgt/KB Ops	1/8/2019
12/14/2018	Manage S-61 discharge to reduce stage in Lake Toho to 54 ft over the next 7-9 days.	Move water to KCH to reduce the rate of stage decline in KCH; reduce the head difference between S-61 headwater and tailwater.	N/A	SFWMD Water Mgt/KB Ops	12/18/2018
12/10/2018	Reduce S-65A discharge to 180 cfs.	Reduce rate of stage decline in lakes Kissimmee-Cypress-Hatchineha	N/A	SFWMD Water Mgt/KB Ops	12/11/2018
12/3/2018	No new recommendations.		N/A		12/4/2018
11/26/2018	No new recommendations.		N/A		11/27/2018
11/19/2018	No new recommendations.		N/A		11/20/2018
11/12/2018	No new recommendations.		N/A		11/13/2018
11/2/2018	Reduce S-65/S-65A discharge to approximately 250 cfs.	To conserve stage in Lake Kissimmee.	Implemented	SFWMD Water Mgt/KB Ops	11/6/2018
10/30/2018	No new recommendations.		N/A		10/30/2018
10/22/2018	Reduce S-65/S-65A discharge to approximately 300 cfs (minimum discharge) in one step of approximately 1100 cfs today.	Reduce rate of stage decline in lakes Kissimmee-Cypress-Hatchineha	Implemented	SFWMD Water Mgt/KB Ops	10/23/2018
10/16/2018	No new recommendations.		N/A		10/16/2018
10/9/2018	No new recommendations.		N/A		10/9/2018
10/2/2018	No new recommendations.		N/A		10/2/2018
9/25/2018	No new recommendations.		N/A		9/25/2018
9/18/2018	No new recommendations.		N/A		9/18/2018
9/11/2018	No new recommendations.		N/A		9/11/2018
9/4/2018	No new recommendations.		N/A		9/4/2018
8/28/2018	No new recommendations.		N/A		8/28/2018
8/21/2018	No new recommendations.		N/A		8/21/2018
8/14/2018	No new recommendations.		N/A		8/14/2018
8/7/2018	No new recommendations.		N/A		8/7/2018

Dry Season Operations Slide 1 - 2018-2019 (NOTE revised discharge table)



Other Considerations

- KCH starting stage may vary; the maximum is 50.75 ft NGVD on Feb 15.
- Maintain S65/S65A discharge of at least 300 cfs.
- If outlook is for extreme dry conditions meet with KB staff to discuss modifications to this plan.

Version 1: January 14 2019

Discharge Rate of Change Limits for S65/S65A (revised 1/14/19).

Q (cfs)	Maximum rate of INCREASE (cfs/day)	Maximum rate of DECREASE (cfs/day)
0-300	100	-50
301-650	150	-75
651-1400	300	-150
1401-3000	600	-300
>3000	1000	-1000

Figure 11A. Slide 1 of the 2018-2019 Dry Season Operations Plan for S-59, S-61, and S-65/S-65A.

Dry Season Operations Slide 2 - 2018-2019

East Lake (ELT) and Toho (WLT)

- **East Toho and Toho Recessions:**
 - Make releases to begin recessions on Jan 15 with lake stage approximately 0.4 ft below winter pool and continue to follow straight line recessions through May 31st to the extent practical
- **East Toho and Toho Stage Reversals :**
 - Adjust discharge to bring stage back to the recession line within about a week
 - Pre-storm releases may be used to lower stage below the recession line and create storage of about half of the forecast rain volume
 - If stage cannot be brought back to the recession line within about a week, the recession line may need to be reset following discussion with partner agencies
 - In general, the water released from ELT and WLT basins will be released to KHC (to the extent that hydraulic capacity is available) without consideration for Lake KHC stage. However, the priority of KCH is subject to change if more nesting occurs in KCH than Toho or East

Kissimmee-Cypress-Hatchineha (KCH)

- **KCH Recession:**
 - Begin recession on February 15 (subject to change) starting no higher than 50.75 feet
 - To the extent feasible considering discharge constraints, make releases to follow a straight-line recession through May 31
 - In general, use the available storage in Lake KCH to keep flow at S-65D below 1,000 cfs; when possible keep flow below 600 cfs
- **KCH Stage Reversals :**
 - To address reversals, in general increase flow by 100 cfs for every 0.1 foot of rise above the recession line (e.g. from 300 cfs at the line to 800 cfs at 0.5 feet above the line)

Figure 11B. Slide 2 of the 2018-2019 Dry Season Operations Plan for S-59, S-61, and S-65/S-65A.




Figure 12. Interim operations schedule for S-65. The discharge schedule shown to the right has not been used in recent years.




Figure 13. Kissimmee River Wading Bird and Waterfowl Surveys from November 2018 to March 2019.

Table 3. Upper Kissimmee Basin Snail Kite Survey Update

Survey 3: April 27-29, 2019

Area	KITES	TOTAL NESTS	SUCCESSFUL	ACTIVE NESTS
E TOHO	8	4	0	2
TOHO	124	48	1	18
KISSIMMEE	235	37	9	27
Grand Total	367	89	10	47




Figure 14. The Kissimmee Basin.

LAKE OKEECHOBEE

According to the USACE web site, Lake Okeechobee average daily lake stage is at 11.23 feet NGVD for May 07, 2019 decreasing 0.01 feet from the previous week and experiencing a reversal due to recent rains. This value is based on the use of four interior lake stations (L001, L005, L006 and LZ40) and three perimeter stations (S-308, S-4 and S-133). The Lake is now 0.57 feet lower than a month ago and 1.74 feet lower than a year ago when stages were still recovering from Hurricane Irma (Figure 1). The Lake dropped into the Beneficial Use sub-band on March 7, 2019 and is currently 0.37 feet above the Water Shortage sub-band (Figure 2). Even with the slight reversal, Lake stage is still the lowest for this time of year since 2011, which followed a very dry rainy season in 2010 (Figure 3). According to RAINDAR, during the week of April 29 to May 6, 2019, 1.83 inches of rain fell directly over the Lake. Rainfall across the rest of the watershed was highly variable with regions receiving as little as 0.1 inches of rain to almost 6 inches of rain (Figure 4).

Average daily inflows (minus rainfall) to the Lake this week decreased at all structures and were lower than last week at 434 cfs compared to 674 cfs, respectively. The inflows from the Kissimmee River decreased slightly, going from 427 cfs to 352 cfs. Inflows from Lake Istokpoga into the Kissimmee River (via the S-84 structures) also decreased from the previous week, going from 243 cfs to 79 cfs. Fisheating Creek outflow decreased from 4 cfs last week to 2 cfs this week (Table 1).

Total outflows (minus evapotranspiration) also decreased from the previous week, going from 3,431 average daily cfs to 1,785 cfs this past week mostly due to decreased outflows south through the S-350 structures (Table 1). Outflows south decreased from 2,558 cfs to 1,348 cfs. Outflows west via S-77 also decreased from the previous week going from 939 cfs to 418 cfs. Outflows east via S-308 and from the L-8 at Canal Point were minimal at 11 cfs and 5 cfs, respectively. The corrected average daily evapotranspiration value for the week based on the L006 and LZ40 weather platform solar radiations was 0.19 inches this week.

Total lake inflows and outflows for the past week are detailed in Table 1, as well as the approximate change in lake stage from each major structure's total flows over the period. Figure 5 shows the combined average daily cfs for inflows and outflows for the Lake over the past eight weeks. These data are provisional and are subject to change.

The most recent satellite imagery (May 03, 2019) using NOAA's cyanobacteria monitoring product derived from EUMETSAT's Sentinel OLCI sensor data showed bloom potential is low for most of the Lake but is increasing and in the medium range along the north and western shores, particularly in Fisheating Bay and along Indian Prairie (Figure 6). The color scheme that classifies algal densities in the image has changed, so direct comparison between the latest image and earlier images is more difficult.

The most recent wading bird survey (May 03, 2019) reported almost 8,000 foraging wading birds on the lake, a considerable decrease from the April 26, 2019 survey (Figure 7). Suitable foraging habitat is limited to small ponded areas of the marsh and along the outer marsh-lake interface (Figure 8, yellow/green areas). Current water levels remain too shallow to support much wading bird breeding effort on the lake this nesting season.

Water Management Recommendations

Lake Okeechobee stage is 11.23 feet NGVD, decreasing 0.01 feet from the previous week. The Lake dropped into the Beneficial Use sub-band on March 7, 2019 and is now 0.37 feet above the Water Shortage sub-band. The lake remains below the bottom of the ecological envelope (currently 1.26 feet below), which varies seasonally from 12.5 – 15.5 feet NGVD. Given the continued potential for above average rainfall (associated with a weak El Niño and the upcoming wet season) over the next few months, and the poor condition of SAV and EAV in the nearshore zone, these lower lake stages are ideal for vegetation recovery. However, low stages will reduce habitat for fish and wildlife in the near-term and encourage spread of invasive vegetation in the upper marsh. Estimated algal bloom potential using satellite imagery suggests medium bloom risk in the north of the lake and along the western shore, particularly within Fisheating Bay and along the edge of Indian Prairie.

Table 1. Average daily inflows and outflows and the approximate depth equivalents on Lake Okeechobee for various structures.

INFLOWS	Previous week Avg Daily CFS	Avg Daily Flow cfs	Equivalent Depth Week Total (in)	OUTFLOWS	Previous week Avg Daily CFS	Avg Daily Flow cfs	Equivalent Depth Week Total (in)
S65E & S65EX1	427	352	0.2	S77	939	418	0.2
S71 & 72	0	0	0.0	S308	-78	11	0.0
S84 & 84X	243	79	0.0	S351	1117	551	0.3
Fisheating Creek	4	2	0.0	S352	782	629	0.3
S154	0	0	0.0	S354	659	169	0.1
S191	0	0	0.0	L8 Outflow	12	7	0.0
S133 P	0	0	0.0	ET	2293	2768	1.3
S127 P	0	0	0.0	Total	5723	4554	2.2
S129 P	0	0	0.0				
S131 P	0	0	0.0				
S135 P	0	0	0.0				
S2 P	0	0	0.0				
S3 P	0	0	0.0				
S4 P	0	0	0.0				
L8 Backflow							
Rainfall	0	3773	1.8				
Total	674	4207	2.0				

Provisional Data




Figure 1. Water depth estimates on Lake Okeechobee based on the South Florida Water Depth Assessment Tool.

Lake Okeechobee Water Level History and Projected Stages




Figure 2. Recent Lake Okeechobee stage and releases, with projected stages based on a dynamic position analysis.

Lake Okeechobee Water Level Comparison




Figure 3. Select annual stage hydrographs for Lake Okeechobee from 2010 – 2019.

SFWMD PROVISIONAL RAINDAR 7-DAY BASIN RAINFALL ESTIMATES
FROM: 0515 EST, 04/30/2019 THROUGH: 0515 EST, 05/07/2019




Figure 4. Rainfall estimates by basin.




Figure 5. Major inflows (orange) and outflows (blue) of Lake Okeechobee, including the S-350 structures designated as South (green). The L-8 Canal flows through Culvert 10A are included as outflows when positive, and as inflows when backflowing into the lake. All inflows and outflows are shown as positive and negative, respectively, for visual purposes.




Figure 6. Potential for cyanobacterial blooms on Lake Okeechobee based on NOAA's harmful algal bloom monitoring system derived from Copernicus Sentinel-3 OLCI data from EUMETSAT. **Note** new color scale on larger image. Gray indicates cloud cover. All data are experimental and unvalidated at this point in product development.

Wading Bird Foraging Locations May 3rd, 2019






Figure 7. Locations of foraging flocks of wading birds observed during a monitoring flight on May 03, 2019 are shown in yellow, with circle sizes representing the size of the flocks. Previous survey totals from this season and from 2018 are compared in the bar graph.




Figure 8. Wading bird habitat suitability index for Lake Okeechobee based on the South Florida Water Depth Assessment Tool.

ESTUARIES

St. Lucie Estuary:

Last week total inflow to the St. Lucie Estuary averaged approximately 556 cfs (Figures 1 and 2) and last month inflow averaged about 273 cfs. Last week's provisional averaged inflows from the tidal basin and the structures are shown in Table 1.

Table 1. Weekly average inflows (data are provisional).

Location	Flow (cfs)
Tidal Basin Inflow	486
S-80	0
S-308	11
S-49 on C-24	0
S-97 on C-23	0
Gordy Rd. structure on Ten Mile Creek	70

Over the past week, salinity remained about the same throughout the estuary (Table 2, Figures 3 and 4). The seven-day moving average of the water column (an average of the surface and bottom salinity) was within the Envelope (10-26). Salinity conditions in the middle estuary are within the good range for adult eastern oysters (Figure 3).

Table 2. Seven-day average salinity at three monitoring sites in the St. Lucie Estuary. Current average is in bold face type, previous average in parentheses. The envelope reflects the preferred salinity range for adult eastern oysters (*Crassostrea virginica*) in the middle estuary.

Sampling Site	Surface	Bottom	Envelope
HR1 (North Fork)	17.6 (18.1)	19.9 (20.3)	NA ¹
US1 Bridge	22.2 ² (22.2)	22.9 ² (22.9)	10.0-26.0
A1A Bridge	29.9 (29.3)	31.4 (31.5)	NA ¹

¹Envelope not applicable and ²Four day average

Caloosahatchee Estuary:

Last week total inflow to the Caloosahatchee Estuary averaged approximately 1,227 cfs (Figures 5 and 6) and last month inflow averaged about 1,165 cfs. Last week's provisional averaged inflows from the structures are shown in Table 3.

Table 3. Weekly average inflows (data is provisional).

Location	Flow (cfs)
S-77	418
S-78	644
S-79	1045
Tidal Basin Inflow	182

Over the past week, salinity increased throughout the estuary (Table 4, Figures 7 & 8). The seven-day average salinity values are estimated to be within the good range for adult eastern oysters at Cape Coral and at Shell Point and in the fair range at Sanibel (Figure 9). The 30-day moving average surface salinity is 0.6 at Val I-75 and 5.6 at Ft. Myers. Salinity conditions between Val I-75 and Ft. Myers are good for tape grass.

Table 4. Seven-day average salinity at six monitoring stations in the Caloosahatchee Estuary. Current average is in bold face type, previous average in parentheses. The envelope reflects the preferred salinity range for tape grass (*Vallisneria americana*) at Val I-75 and for adult eastern oysters (*Crassostrea virginica*) elsewhere.

Sampling Site	Surface	Bottom	Envelope
S-79 (Franklin Lock)	1.2 (0.4)	1.2 (0.4)	NA ¹
Val I75	1.5 (0.4)	1.9 (0.7)	0.0-5.0 ²
Ft. Myers Yacht Basin	7.9 (5.2)	8.4 (8.4)	NA
Cape Coral	14.5 (13.6)	16.9 (17.0)	10.0-30.0
Shell Point	28.7 (26.7)	28.7 (27.0)	10.0-30.0
Sanibel	32.2 (30.8)	33.0 (31.6)	10.0-30.0

¹Envelope not applicable and ²Envelope is based on a 30-day average.

Forecast of surface salinity (Table 5 and Figure 10) at Val I-75 for the next two weeks using the autoregression model (Qiu and Wan, 2013) coupled with a linear reservoir model for the tidal basin predicts daily salinity ranging from 2.7 to 5.5 at the end of the next two weeks for pulse release at S-79 ranging from 0 to 800 cfs and Tidal Basin inflows of 170 cfs.

Table 5. Predicted salinity at Val I-75 at the end of forecast period

Scenario	Q79 (cfs)	TB runoff (cfs)	Daily salinity	30 day mean
A	0	170	5.5	2.2
B	300	170	4.4	1.9
C	450	170	3.7	1.8
D	650	170	3.2	1.7
E	800	170	2.7	1.4

Red tide

The Florida Fish and Wildlife Research Institute reported on May 3, 2019, that *Karenia brevis*, the Florida red tide dinoflagellate, was observed at background concentrations in one samples collected from Lee and was not observed in samples collected from Palm Beach county (no samples from St. Lucie, Martin, Broward or Miami-Dade counties).

Water Management Recommendations

Lake stage is in the Beneficial Use sub-band of 2008 LORS. Tributary hydrological conditions are normal. The 2008 LORS recommends no release at S-79 and S-80. Given the current estuarine conditions, there are no ecological benefits to the upper estuary associated with freshwater releases from Lake Okeechobee, but some benefits may accrue to areas further downstream.




Figure 1. Basins, water control structures, and salinity monitoring for the St. Lucie Estuary.




Figure 2. Total daily inflows from Lake Okeechobee and runoff from the C-44, C-23, C-24, Ten Mile Creek, and tidal basins into the St. Lucie Estuary.




Figure 3. Seven-day mean salinity of the water column at the US1 Bridge.




Figure 4. Daily mean salinity at the A1A, US1 and estimated HR1 stations.




Figure 5. Basins, water control structures, and salinity monitoring for the Caloosahatchee Estuary.




Figure 6. Total daily inflows from Lake Okeechobee, runoff from the C-43 basin, and tributaries in the tidal basin into the Caloosahatchee River Estuary.




Figure 7. Daily mean flows at S-79 and salinity at upper estuary monitoring stations.




Figure 8. Daily mean flows at S-79 and salinity at lower estuary stations.




Figure 9. Seven-day mean salinity at Cape Coral, Shell Point, and Sanibel monitoring stations.





Figure 10. Forecasted Val I-75 surface salinity assuming no pulse release at S79.

EVERGLADES


At the gauges monitored for this report the stages in the Everglades ascended on average 0.13 feet last week. All the basins experienced a reversal except ENP. The most extreme/moderate individual gauge changes ranged from +0.36 feet (NE WCA-3A) to -0.07 feet (ENP). Pan evaporation was estimated at 1.87 inches this week.

Everglades Region	Rainfall (Inches)	Stage Change (feet)			
WCA-1	1.83	+0.19			
WCA-2A	1.21	+0.01			
WCA-2B	2.84	+0.15			
WCA-3A	1.95	+0.22			
WCA-3B	2.01	+0.07			
ENP	0.17	-0.07			


	Good	Recession rate for wading bird foraging
	Fair	
	Poor	




Regulation Schedules: WCA1: Gauge 1-8C is 0.12 feet below the Zone A1 regulation line and following the seasonal recession. WCA2A: S11B Headwater stage is now 0.23 feet below the Zone A regulation line. WCA-3A: The Three Gauge Average stage is 0.12 feet above Zone E1 regulation line. WCA-3A at gauge 62 (Northwest corner) is 0.78 feet below the Upper Schedule.



Water Depths and Changes: The WDAT tool for spatial interpolation of depth monthly snapshots indicate stages in WCA-3A North are mostly below ground. Conditions in WCA-1 and WCA-2A look typical for this time of year. WDAT difference output indicates that water levels fell gradually across the majority of the Everglades during the last month except in most northern regions of WCA-3A and eastern WCA-2A. In the “1 Year” inset we see the difference between current depth conditions and those a year ago. Currently the depths are significantly greater across WCA-3A than they were a year ago, but lower in WCA-1 and WCA-2A. Conditions in the Lostman’s slough region remain significantly wetter than they were a year ago.

**SFWDAT Water Depth Monthly Snapshots**

South Florida Water Depth Assessment Tool (SFWDAT)

**SFWDAT Everglades Difference Maps (Present - Past)**


South Florida Water Depth Assessment Tool (SFWDAT)



Wildlife Update bullets:

- lost WHIB nests in Alley North in NE WCA-3A currently approximately 2000 nests
- some foraging flocks in southern WCA-3A, not much nesting
- approximately 800 WHIB in LOX (Col99); Boat ramp colony and LOX West becoming active
- large foraging flocks in southwestern 2A
- 8k WHIB nesting in ENP coastal colonies
- Despite recent rains conditions look favorable for continued CSSS nesting


Taylor Slough Water Levels: Stages in Taylor Slough and the ENP panhandle are decreasing as is typical for this time of year. The weekly recession averaged 0.18 feet this past week to leave the marsh area averaging a depth of 0.04 feet by Sunday. Craighead Pond (CP) is the only station that have above ground water at this point and that station is below sea-level. This is typical for the end of the dry season.

Florida Bay Salinities: Salinity in Florida Bay averaged a 2.1 psu increase from last week with the maximum change at an individual station being 4 psu in the northeast. Daily average salinities ranged from 35 psu in the northeast to 45 psu in the western nearshore area and are approximately 5 psu above average for this time of year.





Florida Bay MFL: Salinity in the mangrove zone increased 10 psu over the week to end at 20 psu. The 30-day moving average decreased 2.3 psu to 18.4 psu. The weekly cumulative flow from the five creeks denoted by yellow stars on the map totaled about -1,000 acre-feet with negative flows occurring 4 of the last 7 days. At this time of year, there is very little gravity driven downstream flow. The 365-day moving sum of flow from the five creeks (tracked as part of the Florida Bay MFL criteria) has decreased about 8,800 acre-feet to 215,999 acre-feet (less than the long-term average of 257,628 acre-feet but above the 25th percentile). Creek flow is provisional data from the USGS and is highly variable



Water Management Recommendations

The majority of WCA-3A North stages have gone below ground. Protecting peat soils in that sensitive region always has ecological benefit. Wading bird nesting is currently very uncertain in the WCAs. Reductions in the nesting effort at the Alley North colony may be off- set by new nests being seen initiated in WCA-1. Starting this late in the season these nesting efforts may only be successful if decent foraging conditions (recession rates near .05 to .09 feet per week) can be maintained. Large wading bird flocks have been noted foraging in western WCA-2A. Moderating any reversals and maintaining near optimal recession rates in WCA-2A and WCA-3A South will have ecological benefit throughout what remains of the wading bird nesting season. Discharges in WCA-2A should be protective of marsh stage recession rates. More specific recommendations appear in the summary table below. The red text represents new or modified information or recommendations.

SFWMD Everglades Ecological Recommendations, May 7th, 2019 (red is new)

Area	Weekly change	Recommendation	Reasons
WCA-1	Stage increased by 0.19'	Maintain depths at regulation schedule. Moderate reversals when possible. Manage recession rates not to exceed the recommended max rate for optimal wading bird foraging of -0.09 ft per week.	Protect upstream/downstream habitat and wildlife.
WCA-2A	Stage increased by 0.01'	Maintain depths at regulation schedule. Moderate reversals when possible. Manage recession rates not to exceed the recommended max rate for optimal wading bird foraging of -0.09 ft per week.	Protect conditions that provide wading bird foraging habitat later into the nesting season.
WCA-2B	Stage increased by 0.15'	Maintain depths at regulation schedule. Maintain recession rates to the extent possible.	Protect upstream/downstream habitat and wildlife.
WCA-3A NE	Stage increased by 0.17'	Maintain depths at regulation schedule.	Protect habitat including peat soil development and wildlife. Protect conditions that provide wading bird foraging habitat later into the nesting season.
WCA-3A NW	Stage increased by 0.36'	Maintain depths at regulation schedule.	
Central WCA-3A S	Stage increased by 0.34'	Maintain depths at regulation schedule. Moderate reversals when possible. Manage recession rates not to exceed the recommended max rate for optimal wading bird foraging of -0.09 ft per week.	Protect tree islands, upstream/downstream habitat and wildlife. Protect conditions that provide wading bird foraging habitat later into the nesting season.
Southern WCA-3A S	Stage decreased by 0.01'		
WCA-3B	Stage increased by 0.07'	Maintain depths at regulation schedule. Moderate recession rates to the extent possible.	Protect upstream/downstream habitat and wildlife.
ENP-SRS	Stage decreased by 0.07'	Make discharges to the Park according to the 2012 WCP rainfall plan.	Protect upstream/downstream habitat and wildlife.
Taylor Slough	Stage changes ranged from -0.12' to -0.33	Move water southward as possible	When available, provide freshwater buffer for downstream conditions. Decrease potential for high phosphorus input to ENP.
FB- Salinity	Salinity changes ranged +0.2 to +4.0 psu	Move water southward as possible	When available, provide freshwater to maintain low salinity buffer and promote water movement.