

MEMORANDUM

TO: John Mitnik, Division Director, Operations, Engineering, and Construction

THROUGH: Dean Powell, Bureau Chief, Water Supply

FROM: SFWMD Staff Water Supply Advisory Team

DATE: May 24th, 2016


SUBJECT: Water Supply Report

District-wide Conditions

Surface and groundwater levels showed mixed trends throughout the District over the last week. The majority of United States Geological Survey (USGS) real-time wells in the Kissimmee Basin (KB) within the District boundaries were in the median percentile range or higher for this time of year. Most of the surface and groundwater stations across the KB recorded increases in water levels during the last seven days. Stages in the Upper East Coast (UEC) canals C-23, C-24, and C-25 were at 21.79, 17.97, and 18.30 feet, respectively, well above the 14 feet NGVD agricultural cutoff level. Approximately half of UEC surficial aquifer wells are in the median percentile range for this time of year. The rest are in the upper 10th to 30th percentile range or higher. Surface and groundwater levels increased in about half of the stations in the Biscayne aquifer. Approximately two thirds of the USGS Biscayne aquifer monitor wells are in their upper 10th to 30th percentile range or higher at this time. Most of the remainder is at median levels.

In the Lower West Coast (LWC), groundwater levels increased in most of the monitor wells over the last seven days. The majority of the wells in the Surficial aquifer are in their upper 10th to 30th percentile range or higher for this time of year. About one third of the Lower Tamiami aquifer wells are at median levels, with most of the remaining sites in the upper 10th to 30th percentile range. Approximately eighty percent of Sandstone aquifer monitor wells are at median levels for this time of year. The remainder is in the upper 10th to 30th percentile range. About three quarters of the Mid-Hawthorn aquifer monitor wells are at median levels or higher for this time of year, with the remainder in the lowest 10th percentile range. **Figure 1** is a USGS map showing conditions on May 23rd, 2016, from a 7-day running average of daily recorded water levels compared to the statistical distribution of daily water levels for the period of record for selected sites in southern Florida.

PROVISIONAL DRAFT -- Subject to Revision



- | | |
|---|--|
| <ul style="list-style-type: none"> Rivers and canals Roads and highways County boundaries Telemetry site No telemetry, monthly download | <p>Water level compared to historical data, without trend analysis:</p> <ul style="list-style-type: none"> Insufficient information available to compute water-level statistics In lowest 10 percent of past water elevations Within lowest 10 to 30 percent of past water elevations Within 20 percent of the median of past water elevations Within highest 10 to 30 percent of past water elevations In highest 10 percent of past water elevations |
|---|--|

**Water levels at selected sites in South Florida,
Based on PROVISIONAL DATA, as of May 23, 2016.**

**Figure 1. Current Water-level Conditions in South Florida (source: USGS,
http://www.sflorida.er.usgs.gov/ddn_data/index_ndt.html)**

Water Supply Technical Input to LORS2008

The Palmer Index for Lake Okeechobee (LOK) Tributary Conditions is -0.41, classified as “normal,” and is in the “low” risk category. The LOK stage for the next two months is projected to be in the Low Flow Sub-Band, and the risk to water supply is categorized as “low.” The Climate Prediction Center’s (CPC) Precipitation Outlook is projected as “normal” for one month and “normal” for three months, leaving both the one month outlook and the three month outlook in the “low” risk category. The LOK Seasonal Net Inflow Forecast is in the “normal to extremely wet” range, with “low” risk to water supply. The Multi-Seasonal Net Inflow Forecast is projected as “wet,” with “low” risk to water supply. The stages in the Water Conservation Areas are all above line 1 and are in the “low” risk category. Groundwater levels in LEC Service Areas are in the “low” risk category. The Year-Round Irrigation Rule is in effect for the LEC Service Areas. **Figure 2** summarizes the water supply risk indicators.

LORS2008 Implementation on 5/23/2016 (ENSO Neutral Condition):

Water Supply Department Technical Input

Water Supply Outlook:

District wide, Raindar rainfall 3.98 inches for the week ending 5/24/2016. Lake stage on 5/23/2016 is 14.29 ft, up 0.62 ft from last week.

The updated May 2016 SFWMM Dynamic Position Analysis [percentile graph](#) and [tracking chart](#) for Lake Okeechobee show that the lake stage is in the Low Operational Sub-Band.

The LORS2008 tributary [indices](#) are classified as **Very Wet**. The PDSI indicates normal condition and the LONIN is Very Wet. The classification is based on the wetter of the two.

Water Supply Risk Evaluation

Area	Indicator	Value	Color Coded Scoring Scheme
LOK	Projected LOK Stage for the next two months	Low Sub-Band	L
	Palmer Index for LOK Tributary Conditions	-0.41 (Normal)	L
	CPC Precipitation Outlook	1 month: Normal	L
		3 months: Normal	L
	LOK Seasonal Net Inflow Forecast	2.69 ft (Normal to Extremely Wet)	L
	El Nino		
	LOK Multi-Seasonal Net Inflow Forecast	3.90 ft (Wet)	L
El Nino			
WCAs	WCA 1: Site 1-7, Site 1-8T, & Site 1-9 Average	Above Line 1 (16.14 ft)	L
	WCA 2A: Site 2-17 HW	Above Line1 (11.90 ft)	L
	WCA-3A: 3 Station Average (Site 63, 64 and 65)	Above Line 1 (9.55 ft)	L
LEC	Service Area 1	Year-Round Irrigation Rule in effect	L
	Service Area 2	Year-Round Irrigation Rule in effect	L
	Service Area 3	Year-Round Irrigation Rule in effect	L

Note: The water supply risk classification based on the Palmer index, as well as the LOK seasonal and multi-seasonal net inflow forecasts use slightly different classification intervals than those used by the 2008-LORS.

Figure 2. Water Supply Risk Indicators